

## MATH 449, HOMEWORK 5

DUE OCTOBER 15, 2014

### Part I. Theory

**Problem 1.** Prove the following norm inequalities.

- For all  $v \in \mathbb{R}^n$ ,  $\|v\|_2^2 \leq \|v\|_1 \|v\|_\infty$ .
- For any norm  $\|\cdot\|$  on  $\mathbb{R}^n$ , if  $\lambda$  is an eigenvalue of  $A \in \mathbb{R}^{n \times n}$ , then  $\|A\| \geq |\lambda|$  in the induced norm.

**Problem 2.** Consider the function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined by

$$f(x_1, x_2) = \begin{pmatrix} x_1^3 - 3x_1x_2^2 - 1 \\ 3x_1^2x_2 - x_2^3 \end{pmatrix}.$$

Show that  $f(x_1, x_2) = 0$  has three solutions:  $(1, 0)$ ,  $(-\frac{1}{2}, \frac{\sqrt{3}}{2})$ , and  $(-\frac{1}{2}, -\frac{\sqrt{3}}{2})$ .  
Hint: Start by solving the second equation for  $x_2$  in terms of  $x_1$ , and then substitute this into the first equation.

*Remark.* The function in Problem 2 can be identified with the complex-valued function  $z \mapsto z^3 - 1$ , whose roots are the three complex cube roots of 1 (or “roots of unity”). Indeed, if  $z = x_1 + ix_2$ , then we have

$$\begin{aligned} (x_1 + ix_2)^3 - 1 &= x_1^3 + 3x_1^2(ix_2) + 3x_1(ix_2)^2 + (ix_2)^3 - 1 \\ &= x_1^3 + i3x_1^2x_2 - 3x_1x_2^2 - ix_2^3 - 1 \\ &= (x_1^3 - 3x_1x_2^2 - 1) + i(3x_1^2x_2 - x_2^3) \end{aligned}$$

whose real and imaginary parts are precisely the two components of  $f$ .

### Part II. Programming

**Instructions.** For the programming portion of this assignment, you will be running and modifying the code in the provided file `hw5.py`. Hand in a printed copy of the modified file, as well as a printout of the IPython terminal session(s) containing the commands and output you used to get your answers.

**Problem 3.**

- Write a function `hilbert(n)` which returns the  $n \times n$  Hilbert matrix. Print the output of `hilbert(5)`.

- b. Let  $A$  be the  $20 \times 20$  Hilbert matrix,  $x = (1, \dots, 20)$ , and  $b = Ax$ . Find  $\delta x$  by comparing  $x$  with the numerical solution `solve(A,b)` (which uses NumPy's built-in solver), and then find  $\delta b$  by comparing  $A(x + \delta x)$  with  $b$ . Using either one of the functions `norm` or `l2norm` for  $\|\cdot\|_2$ , compute and print the values of  $\|\delta b\|_2/\|b\|_2$  and  $\|\delta x\|_2/\|x\|_2$ .

**Problem 4.** Newton's method does not always converge to the closest root to the starting point—in fact, the behavior can be very complex. In  $\mathbb{R}^2$ , we can obtain fractal patterns called “Newton fractals” by coloring in the regions that converge to each root. The function `newtonFractal` in `hw5.py` does exactly this.

- a. The function  $f$  from Problem 2 is already coded in `hw5.py` as `f(x)`. Create a function `Jf(x)` which returns the Jacobian matrix  $J_f(x)$ . Produce the corresponding Newton fractal by running `newtonFractal(f, Jf)`, and print the resulting image.
- b. Recall, from the remark following Problem 2, that  $f$  gives the real and imaginary components of  $(x_1 + ix_2)^3 - 1$ . Now, create a new function `g(x)`, corresponding to the real and imaginary parts of  $(x_1 + ix_2)^4 - 1$ , and its Jacobian matrix function `Jg(x)`. Run `newtonFractal(g, Jg)` and print the resulting Newton fractal.




FIGURE 1. Newton fractal for  $(x_1 + ix_2)^5 - 1$ .