

# MATH 450, HOMEWORK 1

DUE JANUARY 29, 2016

## Part I. Theory

**Problem 1.** Given  $a \in \mathbb{R}$ , consider the scalar initial value problem

$$(1) \quad y' = ay, \quad y(0) = 1.$$

- Prove that  $f(t, y) = ay$  is Lipschitz. What is the constant  $\lambda$ ?
- Recall that the Picard iteration for (1) is defined recursively by

$$y_0(t) = 1, \quad y_{k+1}(t) = 1 + \int_0^t ay_k(s) ds.$$

Find the first three iterates  $y_1(t)$ ,  $y_2(t)$ , and  $y_3(t)$ .

- Find a general expression for  $y_k(t)$ , and use this to prove directly that  $\lim_{k \rightarrow \infty} y_k(t) = e^{at}$ .

**Problem 2.** Suppose Euler's method is applied to (1) for  $t \in [0, 1]$  by taking  $N$  time steps of size  $h = 1/N$ . Find a general expression for  $y_n$ , and use this to prove directly that  $y_N$  converges to  $y(1)$  as  $N \rightarrow \infty$ .

*Hint:* Use the identity  $e^a = \lim_{N \rightarrow \infty} (1 + a/N)^N$ .

## Part II. Programming

To get started, download and install the Anaconda software package from <http://continuum.io/downloads>. Next, start the Anaconda Launcher, and install and launch the Spyder application. (You can also launch *Spyder* by typing `spyder` in a command line terminal.) Finally, open the file `hw1.py` (available on the class web page) in Spyder, and click the green "play" button to run the code in the IPython console.

The file `hw1.py` contains a single function, `euler(f, t0, y0, h, N)`, which computes an approximate solution to the initial value problem

$$y' = f(t, y), \quad y(t_0) = y_0,$$

by performing  $N$  steps of Euler's method with time step size  $h$ . The function returns the array  $(y_0, y_1, \dots, y_N)$ .

**Problem 3.** Approximate  $e = 2.71828\dots$  by applying `euler` to (1) with  $a = 1$  on the interval  $t \in [0, 1]$ . Use  $h = 1, 0.1, 0.01, 0.001$ , and  $0.0001$ .

**Problem 4.** Reproduce the first plot at the top of p. 11 by (a) applying `euler` to the initial value problem

$$y' = -y + 2e^{-t} \cos 2t, \quad y(0) = 0,$$

on the interval  $t \in [0, 10]$  with  $h = \frac{1}{2}, \frac{1}{10}$ , and  $\frac{1}{50}$ ; and (b) plotting the log absolute error,  $\ln|y_n - y(t_n)|$ , where  $y(t) = e^{-t} \sin 2t$  is the exact solution.

(Don't worry about the formatting of the plot: axis limits, aspect ratio, line styles, etc. Just focus on getting the plot itself correct.)