

Some Remarks on Prill's Problem

N. Mohan Kumar

Abstract.

If $f : X \rightarrow Y$ is a non-constant map of smooth curves over \mathbb{C} and if there is a degree two map $\pi : X \rightarrow C$ where C is a smooth curve with genus less than that of Y , we show that for a general point $P \in Y$, $f^{-1}(P)$ does not move except possibly in one particular case. In particular, this implies that Prill's problem has an affirmative answer if X as above is hyperelliptic or if f is Galois.

2000 *Mathematics Subject Classification.* 14F05

Key words and phrases. Curves, divisors

§1. Introduction

Let $f : X \rightarrow Y$ be a finite morphism of non-singular irreducible projective curves over \mathbb{C} of degree d . Let G (resp. g) denote the genus of X (resp. Y). Further assume that $g \geq 2$. Then Prill's problem states that for a general point $P \in Y$, $f^{-1}(P)$ does not move. That is, $H^0(X, f^*\mathcal{O}_Y(P)) = \mathbb{C}$ (See Arbarello et. al. pp268 [ACGH85]). Since Prill's problem has an affirmative solution if f is cyclic (that is, f is Galois with Galois group cyclic), which will be shown in Proposition 2 and is well known, we will assume that $d \geq 3$, noting that any degree two map is cyclic. One of the consequences stated in [ACGH85] is that if f is as above and Galois and if Prill's problem is false for this f then $h^0(X, f^*\mathcal{O}_Y(P)) > 2$. We will write down a proof of this for completeness in Proposition 3.

Recently, it has been shown by Biswas and Butler in [BB05] that Prill's problem has an affirmative answer if X is hyperelliptic. Our theorem below is a generalization of theirs. Our methods are somewhat different from theirs and might be of independent interest.

Received November 22, 2005.

To Masayoshi Miyanishi who initiated me on the beautiful theory of Algebraic Surfaces .

I thank G. V. Ravindra for numerous discussions on the topic while I was preparing this paper.

Theorem 1. *Let $f : X \rightarrow Y$ be as above. Assume that one has a degree two morphism $\pi : X \rightarrow C$ where C is a non-singular curve with genus $\rho < g$. Then either Prill's problem has an affirmative answer for f or f is étale, $g = 2$ and $\rho = 1$.*

In particular, Prill's problem has an affirmative answer if either X is hyperelliptic or if f is Galois.

§2. Preliminaries

Here we collect some results on Prill's problem, which are mostly well known. We fix our notation $f : X \rightarrow Y$ to be a finite map of degree d with G (resp. g) denoting the genus of X (resp. Y). Also $g \geq 2$.

Proposition 2. *Let f be cyclic. Then Prill's problem has an affirmative answer.*

Proof. If f is cyclic, then $f_*\mathcal{O}_X$ is a direct sum of line bundles on Y , the eigenspaces for the cyclic group. Thus $f_*\mathcal{O}_X = \bigoplus_{i=1}^d L_i$ and clearly we may assume that $L_1 = \mathcal{O}_Y$, $\deg L_i \leq 0$ and $H^0(Y, L_i) = 0$ for $i > 1$. Thus it suffices to prove that if L_2, \dots, L_d are a finite set of line bundles on Y with $\deg L_i \leq 0$ with no sections, then for a general point $P \in Y$, $H^0(Y, L_i(P)) = 0$ for $i \geq 2$. Thus it suffices to prove that for a single such line bundle $L = L_i$, the set S of points P with $H^0(Y, L(P)) \neq 0$ is a finite set.

It is clear that if $\deg L < -1$, then S is empty. So, we may assume that $\deg L = 0$ or -1 . If it is -1 and if $P \in S$, we see that $L = \mathcal{O}_Y(-P)$. Then for any point $Q \neq P$, $L(Q)$ has no section since no two distinct points can be rationally equivalent. If $\deg L = 0$ and $P \in S$, then $L = \mathcal{O}_Y(Q - P)$ for some point Q . Since $H^0(L) = 0$, $Q \neq P$. If $P \neq R \in S$, we see that there exists a point R' such that $Q + R \sim P + R'$. This implies that Y is hyperelliptic and if σ is the hyperelliptic involution, S consists of at most two points, $P, \sigma(Q)$. Q.E.D.

The following is essentially the content of the exercise in [ACGH85].

Proposition 3. *Let f as above be Galois and assume that Prill's problem has a negative answer for f . Then for a general point $P \in Y$, $h^0(X, f^*\mathcal{O}_Y(P)) > 2$.*

Proof. On the contrary, assume that $W_P = H^0(X, f^*\mathcal{O}_Y(P)) = 2$ for a general point $P \in Y$. Let G be the Galois group. Thus G acts on W_P and if for a general point P , the group homomorphism $G \rightarrow \text{Aut } \mathbb{P}(W_P)$ is not injective, then by continuity, there is a normal subgroup $H \subset G$ which acts trivially on W_P for all P . If we consider the map $f' : X/H \rightarrow Y$, we immediately see that for a general point $P \in Y$, $h^0(X/H, f'^*(\mathcal{O}_Y(P))) =$

2. Since f' is Galois with Galois group G/H , we may replace X by X/H and thus assume to start with that the map $G \rightarrow \text{Aut } \mathbb{P}(W_P)$ is injective for general $P \in Y$ and the map f itself. But, the section corresponding to $f^{-1}(P)$ is fixed by G and thus $G \subset \text{Aut } \mathbb{A}^1$. We have an exact sequence of groups,

$$1 \rightarrow \mathbb{C} \rightarrow \text{Aut } \mathbb{A}^1 \rightarrow \mathbb{C}^* \rightarrow 1.$$

Since G is finite, this implies that G is a subgroup of \mathbb{C}^* and hence cyclic. Now by Proposition 2 we are done. Q.E.D.

The following has been proved in [BB05]. Our proof is somewhat different.

Proposition 4. *If Prill's problem is false for f , then f_*K_X is not generically globally generated. That is, the subsheaf of f_*K_X generated by $H^0(Y, f_*K_X)$ has rank less than d .*

Proof. Suffices to show that for a general point $P \in Y$ the natural map $H^0(f_*K_X) \rightarrow H^0(f_*K_{X|_P})$ is not onto. Since the latter is a vector space of dimension d and the former is a vector space of dimension G , suffices to show that the kernel $H^0(f_*K_X(-P))$ has dimension greater than $G - d$. By Serre duality, this is just the dimension of $H^1(X, f^*(P))$. By Riemann-Roch we have, $h^1(X, f^*(P)) = h^0(f^*(P)) - 1 + G - d$ and by hypothesis $h^0(f^*(P)) > 1$. Q.E.D.

Proposition 5. *Let $f : X \rightarrow Y$ be a finite map of non-singular curves. Assume that we have finite morphisms $\phi : Y \rightarrow \mathbb{P}^1$ and $\psi : Z \rightarrow \mathbb{P}^1$ where Z is a non-singular curve. Further assume that $Z' = Z \times_{\mathbb{P}^1} Y$ is irreducible and we have a morphism $\eta : Z' \rightarrow X$ such that the composite $Z' \rightarrow X \rightarrow Y$ is the natural projection $Z' \rightarrow Y$. Then Prill's problem has an affirmative answer for f .*

Proof. If Prill's problem is false for f , clearly it is false for $f' = f \circ \eta$, though Z' may be singular. If $p : Z' \rightarrow Z$ and $q : Z' \rightarrow Y$ denote the two projections, for any point $P \in Y$, we have, $p_*f'^*(\mathcal{O}_Y(P)) = \psi^*\phi_*(\mathcal{O}_Y(P))$. If we write $\phi_*(\mathcal{O}_Y(P))$ as a direct sum of line bundles $\oplus L_i$, $H^0(Y, \mathcal{O}_Y(P)) = \mathbb{C}$ implies that one of the $L_i = \mathcal{O}_{\mathbb{P}^1}$ and the others have negative degree. But then $\psi^*\phi_*(\mathcal{O}_Y(P))$ is a direct sum of one copy of \mathcal{O}_Z and the rest of negative degree. Thus $H^0(Z', f'^*(\mathcal{O}_Y(P))) = \mathbb{C}$. Q.E.D.

§3. Proof of Theorem 1

Proof. Write $\pi_*\mathcal{O}_X = \mathcal{O}_C \oplus L$ where L is a line bundle of degree $-m$ on C with $m > 0$. For a point $P \in Y$ we have $V_P = \pi_*f^*(\mathcal{O}_Y(P))$

a rank two vector bundle on C . Also we have the natural inclusion $\pi_*\mathcal{O}_X \rightarrow V_P$ using the natural inclusion of $\mathcal{O}_X \subset f^*(\mathcal{O}_Y(P))$. Since the cokernel of this map is a sky-scraper sheaf of length d , we see that $\deg V_P = -m + d$. We have $G = h^1(\pi_*\mathcal{O}_X) = \rho + h^1(L)$. By Riemann-Roch $h^1(L) = m + \rho - 1$ and thus $G = 2\rho + m - 1$. By Riemann-Hurwitz, we have

$$2\rho + m - 2 = G - 1 \geq d(g - 1) = (d - 2)(g - 1) + 2g - 2$$

and thus, $m \geq (d - 2)(g - 1) + 2(g - \rho)$. Since $g > \rho$ and $g \geq 2$, this implies $m \geq d$.

We will separate the cases when $m = d$ and $m > d$. We see from the above that if $m = d$, then $g = 2$ and $\rho = 1$, since we have assumed that $d \geq 3$. Also the above inequality from Riemann-Hurwitz must be an equality. That is f is etale.

So, now on we will assume that $m > d$. Then $\deg V_P < 0$ for any $P \in Y$. Let M be the saturation of \mathcal{O}_C in V_P . We have then an exact sequence $0 \rightarrow M \rightarrow V_P \rightarrow M' \rightarrow 0$ with M, M' line bundles on C and since $\deg M \geq 0$, $\deg M' < 0$. In particular the map $H^0(V_P) \rightarrow H^0(M') = 0$ is zero. So $H^0(M) = H^0(V_P)$ and if Prill's problem is false for f we have $h^0(V_P) > 1$ for a general $P \in Y$. This implies that the inclusion of \mathcal{O}_C in M is strict.

Consider the map $X \times Y \xrightarrow{(\pi, Id)=\phi} C \times Y$. Let $\Gamma \subset X \times Y$ be the graph of f . Also let $p : C \times Y \rightarrow C$ and $q : C \times Y \rightarrow Y$ be the natural projections. We have an exact sequence

$$0 \rightarrow \mathcal{O}_{X \times Y} \rightarrow \mathcal{O}_{X \times Y}(\Gamma) \rightarrow \Gamma_{|\Gamma} \rightarrow 0.$$

Let D be the image of Γ in $C \times Y$. Then we claim that the map $\Gamma \rightarrow D$ is birational. If not, since the composite $\Gamma \rightarrow D \xrightarrow{p} C$ is just π which has degree two, we see that $D \rightarrow C$ must be birational. But C is smooth and thus $D \rightarrow C$ must be an isomorphism. But, we have a morphism $D \xrightarrow{q} Y$ and thus we get a non-constant morphism from $C \rightarrow Y$. This is absurd since $\rho < g$. Taking direct images, we get an exact sequence,

$$0 \rightarrow \phi_*\mathcal{O}_{X \times Y} \rightarrow \phi_*\mathcal{O}_{X \times Y}(\Gamma) \rightarrow \phi_*\Gamma_{|\Gamma} \rightarrow 0.$$

Notice that $\phi_*\mathcal{O}_{X \times Y}(\Gamma) = E$ is a rank two vector bundle on $C \times Y$ since ϕ is a two to one map. Also $\phi_*\mathcal{O}_{X \times Y}$ is just the pull back of $\mathcal{O}_C \oplus L$ by p . Identifying the pull back of \mathcal{O}_C as $\mathcal{O}_{C \times Y}$ let us look at the inclusion of this sheaf in E and let F be the cokernel. I claim that F has torsion. If it has no torsion, then it is a line bundle outside

a finite set of points and thus restricting to a general point $P \in Y$, we get an exact sequence $0 \rightarrow \mathcal{O}_C \rightarrow E|_P = V_P \rightarrow F|_P \rightarrow 0$. Since $F|_P$ is assumed to be a line bundle, we see that \mathcal{O}_C is saturated in V_P , which we have seen is not the case. Thus we see that F has torsion. Taking the inverse image of the torsion subsheaf of F in E , we get an exact sequence, $0 \rightarrow A \rightarrow E \rightarrow E/A \rightarrow 0$ where $\mathcal{O}_{C \times Y} \subset A$ and this inclusion is strict and E/A is torsion free. It is clear that the composite $p^*L \rightarrow E \rightarrow E/A$ is an injection. Thus we get an inclusion $A \oplus p^*L \subset E$ and let B be its cokernel. We have a commutative diagram,

$$\begin{array}{ccccccccc} 0 & \rightarrow & \mathcal{O}_{C \times Y} \oplus p^*L & \rightarrow & E & \rightarrow & \phi_*\Gamma|_\Gamma & \rightarrow & 0 \\ & & \downarrow & & \parallel & & \downarrow & & \\ 0 & \rightarrow & A \oplus p^*L & \rightarrow & E & \rightarrow & B & \rightarrow & 0 \end{array}$$

Thus by snake lemma we get an exact sequence,

$$0 \rightarrow A' \rightarrow \phi_*\Gamma|_\Gamma \rightarrow B \rightarrow 0$$

where A' is the cokernel of $\mathcal{O}_{C \times Y} \subset A$. Since this inclusion is strict, we see that $A' \neq 0$. Since $\phi_*\Gamma|_\Gamma$ is torsion free as an \mathcal{O}_D -module we see that A' is supported on all of D . Since $\phi : \Gamma \rightarrow D$ is birational we see that $\phi_*\Gamma|_\Gamma$ is a line bundle on D for general points of D and thus A' is a line bundle on D at general points of D , since D is a smooth curve generically and these two sheaves are equal at general points of D . This implies that B is supported on a finite set of points of D . But, B is the quotient of a rank two vector bundle by another rank two vector bundle on a smooth surface and thus for homological reasons, either support of B is a divisor or empty. This implies that $B = 0$. So, we get $A \oplus p^*L = E$. Restricting to a general point $P \in Y$ and calling the restricted bundle A_P we see that $A_P \oplus L = V_P$. This implies that $\deg A_P = d$. But, then $\pi^*A_P \subset f^*\mathcal{O}_Y(P)$ and the first line bundle has degree $2d$ and the latter d . This is clearly impossible.

If f is Galois, then the only case left to prove is when $g = 2, \rho = 1$ and f is etale. Then as we saw, $\pi_*\mathcal{O}_X = \mathcal{O}_Y \oplus L$ with $\deg L = -d$. Thus $\deg V_P = 0$ and $h^0(V_P) > 1$ implies $V_P = \mathcal{O}_Y \oplus \mathcal{O}_Y$. Thus $h^0(V_P) = 2$ for a general point $P \in Y$. This contradicts Proposition 3. Q.E.D.

Corollary 6. *If $f : X \rightarrow Y$ has degree 3, then Prill's problem has an affirmative answer.*

Proof. From Theorem 1, we may assume that X is not hyperelliptic. Also note that by Riemann-Hurwitz, since $g \geq 2, G \geq 4$. The morphism f induces a morphism $Y \rightarrow J^3X$ where J^3X is the variety parametrizing line bundles of degree 3. Also, the image is contained in $W_3^1(X)$ if Prill's

problem had a negative answer for this f . Since X is not hyperelliptic, by Martens Theorem [Mar67] (also see pp 191-2 [ACGH85]) we have $\dim W_3^1(X) \leq 0$. Thus image of Y in $J^3 X$ is constant. In other words, for any two points $P, Q \in Y$, $f^*(P) \sim f^*(Q)$. This is impossible since $g > 0$. Q.E.D.

References

- [ACGH85] E. Arbarello, M. Cornalba, P. A. Griffiths, and J. Harris. *Geometry of Algebraic Curves*. Number 267 in Grundlehren I. Springer Verlag, New York, 1985.
- [BB05] Indranil Biswas and David C. Butler. On Prill's problem. *Comm. Algebra*, 33(1):323–330, 2005.
- [Mar67] Henrik H. Martens. On the varieties of special divisors on a curve. *J. Reine Angew. Math.*, 227:111–120, 1967.

Department of Mathematics, Washington University in St. Louis, St. Louis, Missouri, 63130, USA.

E-mail address: `kumar@wustl.edu`

URL: `http://www.math.wustl.edu/~kumar`