Math 132

Exponentials/Logarithms Overview – February 20, 2012

1. Definition

- (a) We defined $\ln x$ to be $\int_1^x \frac{1}{t} dt$.
- (b) e^x is the inverse function of $\ln x$. This means that $\ln e^x = e^{\ln x} = x$.
- (c) e is a real number (about 2.7).
- (d) $e^3 = e \cdot e \cdot e$, and $e^{0.5} = \sqrt{e}$. So $e^{3/2} = \sqrt{e^3}$. (Similarly for other rational powers.)

2. Useful values and limits.

$$\begin{array}{lll} e^0 &=& 1, & & \lim_{x \to \infty} e^x = \infty, & \lim_{x \to -\infty} e^x = 0 \\ \ln 1 &=& 0, & & \ln e = 1, & \lim_{x \to 0+} \ln x = -\infty, & & \lim_{x \to \infty} \ln x = \infty. \end{array}$$

Note: $\ln x$ is not defined for $x \leq 0$!

3. Change of base

(a)
$$2^x = (e^{\ln 2})^x = e^{x \cdot \ln 2}$$

(b)
$$\log_2 x = \frac{\ln x}{\ln 2}$$

In both formulas, 2 can be replaced by any positive number.

4. Derivatives and integrals

$$\frac{d}{dx}e^{x} = e^{x}, \qquad \frac{d}{dx}\ln x = \frac{1}{x}$$

$$\int e^{x} dx = e^{x} + C \qquad \int \frac{1}{x} dx = \ln|x| + C$$

5. Derivatives in other bases:

We consider e^x and $\ln x$ rather than, say, 2^x and $\log_2 x$ because their derivatives and integrals have these natural forms. Compare with the derivatives of 2^x and $\log_2 x$:

$$\frac{d}{dx}2^x = \frac{d}{dx}e^{x \cdot \ln 2} = \ln 2 \cdot e^{x \cdot \ln 2} = \ln 2 \cdot 2^x$$

$$\frac{d}{dx}\log_2 x = \frac{d}{dx}\frac{\ln x}{\ln 2} = \frac{1}{x \cdot \ln 2}$$

Integrals in other bases are handled similarly.

6. Fundamental identities:

$$e^{x+y}=e^x e^y$$
 (i.e., e^x "turns + into ·")
 $e^{xy}=(e^x)^y$
 $\ln xy=\ln x+\ln y$ (i.e., $\ln x$ "turns ·into +")
 $\ln x^y=y\ln x$

7. Graphs.

(The graph of $\ln x$ is that of e^x flipped over the line y=x!)



