

## 614 Chapter 10 Vectors and the Geometry of Space

19. (a)  $x = 3$

(b)  $y = -1$

(c)  $z = -2$

20. (a)  $x = 3$

(b)  $y = -1$

(c)  $z = 2$

21. (a)  $z = 1$

(b)  $x = 3$

(c)  $y = -1$

22. (a)  $x^2 + y^2 = 4, z = 0$

(b)  $y^2 + z^2 = 4, x = 0$

(c)  $x^2 + z^2 = 4, y = 0$

23. (a)  $x^2 + (y - 2)^2 = 4, z = 0$

(b)  $(y - 2)^2 + z^2 = 4, x = 0$

(c)  $x^2 + z^2 = 4, y = 2$

24. (a)  $(x + 3)^2 + (y - 4)^2 = 1, z = 1$

(b)  $(y - 4)^2 + (z - 1)^2 = 1, x = -3$

(c)  $(x + 3)^2 + (z - 1)^2 = 1, y = 4$

25. (a)  $y = 3, z = -1$

(b)  $x = 1, z = -1$

(c)  $x = 1, y = 3$

26.  $\sqrt{x^2 + y^2 + z^2} = \sqrt{x^2 + (y - 2)^2 + z^2} \Rightarrow x^2 + y^2 + z^2 = x^2 + (y - 2)^2 + z^2 \Rightarrow y^2 = y^2 - 4y + 4 \Rightarrow y = 1$

27.  $x^2 + y^2 + z^2 = 25, z = 3 \Rightarrow x^2 + y^2 = 16$  in the plane  $z = 3$

28.  $x^2 + y^2 + (z - 1)^2 = 4$  and  $x^2 + y^2 + (z + 1)^2 = 4 \Rightarrow x^2 + y^2 + (z - 1)^2 = x^2 + y^2 + (z + 1)^2 \Rightarrow z = 0, x^2 + y^2 = 3$

29.  $0 \leq z \leq 1$

30.  $0 \leq x \leq 2, 0 \leq y \leq 2, 0 \leq z \leq 2$

31.  $z \leq 0$

32.  $z = \sqrt{1 - x^2 - y^2}$

33. (a)  $(x - 1)^2 + (y - 1)^2 + (z - 1)^2 < 1$

(b)  $(x - 1)^2 + (y - 1)^2 + (z - 1)^2 > 1$

34.  $1 \leq x^2 + y^2 + z^2 \leq 4$

35.  $|P_1P_2| = \sqrt{(3 - 1)^2 + (3 - 1)^2 + (0 - 1)^2} = \sqrt{9} = 3$

36.  $|P_1P_2| = \sqrt{(2 + 1)^2 + (5 - 1)^2 + (0 - 5)^2} = \sqrt{50} = 5\sqrt{2}$

37.  $|P_1P_2| = \sqrt{(4 - 1)^2 + (-2 - 4)^2 + (7 - 5)^2} = \sqrt{49} = 7$

38.  $|P_1P_2| = \sqrt{(2 - 3)^2 + (3 - 4)^2 + (4 - 5)^2} = \sqrt{3}$

39.  $|P_1P_2| = \sqrt{(2 - 0)^2 + (-2 - 0)^2 + (-2 - 0)^2} = \sqrt{3 \cdot 4} = 2\sqrt{3}$

40.  $|P_1P_2| = \sqrt{(0 - 5)^2 + (0 - 3)^2 + (0 + 2)^2} = \sqrt{38}$

41. center  $(-2, 0, 2)$ , radius  $2\sqrt{2}$

42. center  $(-\frac{1}{2}, -\frac{1}{2}, -\frac{1}{2})$ , radius  $\frac{\sqrt{21}}{2}$

43. center  $(\sqrt{2}, \sqrt{2}, -\sqrt{2})$ , radius  $\sqrt{2}$

44. center  $(0, -\frac{1}{3}, \frac{1}{3})$ , radius  $\frac{\sqrt{29}}{3}$

45.  $(x-1)^2 + (y-2)^2 + (z-3)^2 = 14$

46.  $x^2 + (y+1)^2 + (z-5)^2 = 4$

47.  $(x+2)^2 + y^2 + z^2 = 3$

48.  $x^2 + (y+7)^2 + z^2 = 49$

49.  $x^2 + y^2 + z^2 + 4x - 4z = 0 \Rightarrow (x^2 + 4x + 4) + y^2 + (z^2 - 4z + 4) = 4 + 4$

$$\Rightarrow (x+2)^2 + (y-0)^2 + (z-2)^2 = (\sqrt{8})^2 \Rightarrow \text{the center is at } (-2, 0, 2) \text{ and the radius is } \sqrt{8}$$

50.  $x^2 + y^2 + z^2 - 6y + 8z = 0 \Rightarrow x^2 + (y^2 - 6y + 9) + (z^2 + 8z + 16) = 9 + 16 \Rightarrow (x-0)^2 + (y-3)^2 + (z+4)^2 = 5^2$   
 $\Rightarrow \text{the center is at } (0, 3, -4) \text{ and the radius is } 5$

51.  $2x^2 + 2y^2 + 2z^2 + x + y + z = 9 \Rightarrow x^2 + \frac{1}{2}x + y^2 + \frac{1}{2}y + z^2 + \frac{1}{2}z = \frac{9}{2}$

$$\Rightarrow (x^2 + \frac{1}{2}x + \frac{1}{16}) + (y^2 + \frac{1}{2}y + \frac{1}{16}) + (z^2 + \frac{1}{2}z + \frac{1}{16}) = \frac{9}{2} + \frac{3}{16} \Rightarrow (x + \frac{1}{4})^2 + (y + \frac{1}{4})^2 + (z + \frac{1}{4})^2 = (\frac{5\sqrt{3}}{4})^2$$

$$\Rightarrow \text{the center is at } (-\frac{1}{4}, -\frac{1}{4}, -\frac{1}{4}) \text{ and the radius is } \frac{5\sqrt{3}}{4}$$

52.  $3x^2 + 3y^2 + 3z^2 + 2y - 2z = 9 \Rightarrow x^2 + y^2 + \frac{2}{3}y + z^2 - \frac{2}{3}z = 3 \Rightarrow x^2 + (y^2 + \frac{2}{3}y + \frac{1}{9}) + (z^2 - \frac{2}{3}z + \frac{1}{9}) = 3 + \frac{2}{9}$

$$\Rightarrow (x-0)^2 + (y + \frac{1}{3})^2 + (z - \frac{1}{3})^2 = (\frac{\sqrt{29}}{3})^2 \Rightarrow \text{the center is at } (0, -\frac{1}{3}, \frac{1}{3}) \text{ and the radius is } \frac{\sqrt{29}}{3}$$

53. (a) the distance between  $(x, y, z)$  and  $(x, 0, 0)$  is  $\sqrt{y^2 + z^2}$

(b) the distance between  $(x, y, z)$  and  $(0, y, 0)$  is  $\sqrt{x^2 + z^2}$

(c) the distance between  $(x, y, z)$  and  $(0, 0, z)$  is  $\sqrt{x^2 + y^2}$

54. (a) the distance between  $(x, y, z)$  and  $(x, y, 0)$  is  $z$

(b) the distance between  $(x, y, z)$  and  $(0, y, z)$  is  $x$

(c) the distance between  $(x, y, z)$  and  $(x, 0, z)$  is  $y$

55.  $|AB| = \sqrt{(1 - (-1))^2 + (-1 - 2)^2 + (3 - 1)^2} = \sqrt{4 + 9 + 4} = \sqrt{17}$

$$|BC| = \sqrt{(3 - 1)^2 + (4 - (-1))^2 + (5 - 3)^2} = \sqrt{4 + 25 + 4} = \sqrt{33}$$

$$|CA| = \sqrt{(-1 - 3)^2 + (2 - 4)^2 + (1 - 5)^2} = \sqrt{16 + 4 + 16} = \sqrt{36} = 6$$

Thus the perimeter of triangle ABC is  $\sqrt{17} + \sqrt{33} + 6$ .

56.  $|PA| = \sqrt{(2 - 3)^2 + (-1 - 1)^2 + (3 - 2)^2} = \sqrt{1 + 4 + 1} = \sqrt{6}$

$$|PB| = \sqrt{(4 - 3)^2 + (3 - 1)^2 + (1 - 2)^2} = \sqrt{1 + 4 + 1} = \sqrt{6}$$

Thus P is equidistant from A and B.

## 10.2 VECTORS

1. (a)  $\langle 3(3), 3(-2) \rangle = \langle 9, -6 \rangle$

(b)  $\sqrt{9^2 + (-6)^2} = \sqrt{117} = 3\sqrt{13}$

2. (a)  $\langle -2(-2), -2(5) \rangle = \langle 4, -10 \rangle$

(b)  $\sqrt{4^2 + (-10)^2} = \sqrt{116} = 2\sqrt{29}$

3. (a)  $\langle 3 + (-2), -2 + 5 \rangle = \langle 1, 3 \rangle$





(b)  $\sqrt{1^2 + 3^2} = \sqrt{10}$

4. (a)  $\langle 3 - (-2), -2 - 5 \rangle = \langle 5, -7 \rangle$





(b)  $\sqrt{5^2 + (-7)^2} = \sqrt{74}$

5. (a)  $2\mathbf{u} = \langle 2(3), 2(-2) \rangle = \langle 6, -4 \rangle$   
 $3\mathbf{v} = \langle 3(-2), 3(5) \rangle = \langle -6, 15 \rangle$   
 $2\mathbf{u} - 3\mathbf{v} = \langle 6 - (-6), -4 - 15 \rangle = \langle 12, -19 \rangle$   
 (b)  $\sqrt{12^2 + (-19)^2} = \sqrt{505}$
6. (a)  $-2\mathbf{u} = \langle -2(3), -2(-2) \rangle = \langle -6, 4 \rangle$   
 $5\mathbf{v} = \langle 5(-2), 5(5) \rangle = \langle -10, 25 \rangle$   
 $-2\mathbf{u} + 5\mathbf{v} = \langle -6 + (-10), 4 + 25 \rangle = \langle -16, 29 \rangle$   
 (b)  $\sqrt{(-16)^2 + 29^2} = \sqrt{1097}$
7. (a)  $\frac{3}{5}\mathbf{u} = \langle \frac{3}{5}(3), \frac{3}{5}(-2) \rangle = \langle \frac{9}{5}, -\frac{6}{5} \rangle$   
 $\frac{4}{5}\mathbf{v} = \langle \frac{4}{5}(-2), \frac{4}{5}(5) \rangle = \langle -\frac{8}{5}, 4 \rangle$   
 $\frac{3}{5}\mathbf{u} + \frac{4}{5}\mathbf{v} = \langle \frac{9}{5} + (-\frac{8}{5}), -\frac{6}{5} + 4 \rangle = \langle \frac{1}{5}, \frac{14}{5} \rangle$   
 (b)  $\sqrt{(\frac{1}{5})^2 + (\frac{14}{5})^2} = \frac{\sqrt{197}}{5}$
8. (a)  $-\frac{5}{13}\mathbf{u} = \langle -\frac{5}{13}(3), -\frac{5}{13}(-2) \rangle = \langle -\frac{15}{13}, \frac{10}{13} \rangle$   
 $\frac{12}{13}\mathbf{v} = \langle \frac{12}{13}(-2), \frac{12}{13}(5) \rangle = \langle -\frac{24}{13}, \frac{60}{13} \rangle$   
 $-\frac{5}{13}\mathbf{u} + \frac{12}{13}\mathbf{v} = \langle -\frac{15}{13} + (-\frac{24}{13}), \frac{10}{13} + \frac{60}{13} \rangle = \langle -3, \frac{70}{13} \rangle$   
 (b)  $\sqrt{(-3)^2 + (\frac{70}{13})^2} = \frac{\sqrt{6421}}{13}$
9.  $\langle 2 - 1, -1 - 3 \rangle = \langle 1, -4 \rangle$
10.  $\langle \frac{2+(-4)}{2} - 0, \frac{-1+3}{2} - 0 \rangle = \langle -1, 1 \rangle$
11.  $\langle 0 - 2, 0 - 3 \rangle = \langle -2, -3 \rangle$
12.  $\vec{AB} = \langle 2 - 1, 0 - (-1) \rangle = \langle 1, 1 \rangle$ ,  $\vec{CD} = \langle -2 - (-1), 2 - 3 \rangle = \langle -1, -1 \rangle$ ,  $\vec{AB} + \vec{CD} = \langle 0, 0 \rangle$
13.  $\langle \cos \frac{2\pi}{3}, \sin \frac{2\pi}{3} \rangle = \langle -\frac{1}{2}, \frac{\sqrt{3}}{2} \rangle$
14.  $\langle \cos (-\frac{3\pi}{4}), \sin (-\frac{3\pi}{4}) \rangle = \langle -\frac{1}{\sqrt{2}}, -\frac{1}{\sqrt{2}} \rangle$
15. This is the unit vector which makes an angle of  $120^\circ + 90^\circ = 210^\circ$  with the positive x-axis;  
 $\langle \cos 210^\circ, \sin 210^\circ \rangle = \langle -\frac{\sqrt{3}}{2}, -\frac{1}{2} \rangle$
16.  $\langle \cos 135^\circ, \sin 135^\circ \rangle = \langle -\frac{1}{\sqrt{2}}, \frac{1}{\sqrt{2}} \rangle$
17.  $P_1\vec{P}_2 = (2 - 5)\mathbf{i} + (9 - 7)\mathbf{j} + (-2 - (-1))\mathbf{k} = -3\mathbf{i} + 2\mathbf{j} - \mathbf{k}$
18.  $P_1\vec{P}_2 = (-3 - 1)\mathbf{i} + (0 - 2)\mathbf{j} + (5 - 0)\mathbf{k} = -4\mathbf{i} - 2\mathbf{j} + 5\mathbf{k}$
19.  $\vec{AB} = (-10 - (-7))\mathbf{i} + (8 - (-8))\mathbf{j} + (1 - 1)\mathbf{k} = -3\mathbf{i} + 16\mathbf{j}$
20.  $\vec{AB} = (-1 - 1)\mathbf{i} + (4 - 0)\mathbf{j} + (5 - 3)\mathbf{k} = -2\mathbf{i} + 4\mathbf{j} + 2\mathbf{k}$
21.  $5\mathbf{u} - \mathbf{v} = 5\langle 1, 1, -1 \rangle - \langle 2, 0, 3 \rangle = \langle 5, 5, -5 \rangle - \langle 2, 0, 3 \rangle = \langle 5 - 2, 5 - 0, -5 - 3 \rangle = \langle 3, 5, -8 \rangle = 3\mathbf{i} + 5\mathbf{j} - 8\mathbf{k}$
22.  $-2\mathbf{u} + 3\mathbf{v} = -2\langle -1, 0, 2 \rangle + 3\langle 1, 1, 1 \rangle = \langle 2, 0, -4 \rangle + \langle 3, 3, 3 \rangle = \langle 5, 3, -1 \rangle = 5\mathbf{i} + 3\mathbf{j} - \mathbf{k}$

23. The vector  $v$  is horizontal and 1 in. long. The vectors  $u$  and  $w$  are  $\frac{11}{16}$  in. long.  $w$  is vertical and  $u$  makes a  $45^\circ$  angle with the horizontal. All vectors must be drawn to scale.



24. The angle between the vectors is  $120^\circ$  and vector  $u$  is horizontal. They are all 1 in. long. Draw to scale.



25. length  $= |2\mathbf{i} + \mathbf{j} - 2\mathbf{k}| = \sqrt{2^2 + 1^2 + (-2)^2} = 3$ , the direction is  $\frac{2}{3}\mathbf{i} + \frac{1}{3}\mathbf{j} - \frac{2}{3}\mathbf{k} \Rightarrow 2\mathbf{i} + \mathbf{j} - 2\mathbf{k} = 3\left(\frac{2}{3}\mathbf{i} + \frac{1}{3}\mathbf{j} - \frac{2}{3}\mathbf{k}\right)$

26. length  $= |9\mathbf{i} - 2\mathbf{j} + 6\mathbf{k}| = \sqrt{81 + 4 + 36} = 11$ , the direction is  $\frac{9}{11}\mathbf{i} - \frac{2}{11}\mathbf{j} + \frac{6}{11}\mathbf{k} \Rightarrow 9\mathbf{i} - 2\mathbf{j} + 6\mathbf{k} = 11\left(\frac{9}{11}\mathbf{i} - \frac{2}{11}\mathbf{j} + \frac{6}{11}\mathbf{k}\right)$

27. length  $= |5\mathbf{k}| = \sqrt{25} = 5$ , the direction is  $\mathbf{k} \Rightarrow 5\mathbf{k} = 5(\mathbf{k})$

28. length  $= \left|\frac{3}{5}\mathbf{i} + \frac{4}{5}\mathbf{k}\right| = \sqrt{\frac{9}{25} + \frac{16}{25}} = 1$ , the direction is  $\frac{3}{5}\mathbf{i} + \frac{4}{5}\mathbf{k} \Rightarrow \frac{3}{5}\mathbf{i} + \frac{4}{5}\mathbf{k} = 1\left(\frac{3}{5}\mathbf{i} + \frac{4}{5}\mathbf{k}\right)$

29. length  $= \left|\frac{1}{\sqrt{6}}\mathbf{i} - \frac{1}{\sqrt{6}}\mathbf{j} - \frac{1}{\sqrt{6}}\mathbf{k}\right| = \sqrt{3\left(\frac{1}{\sqrt{6}}\right)^2} = \sqrt{\frac{1}{2}}$ , the direction is  $\frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k}$   
 $\Rightarrow \frac{1}{\sqrt{6}}\mathbf{i} - \frac{1}{\sqrt{6}}\mathbf{j} - \frac{1}{\sqrt{6}}\mathbf{k} = \sqrt{\frac{1}{2}}\left(\frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k}\right)$

30. length =  $\left| \frac{1}{\sqrt{3}}\mathbf{i} + \frac{1}{\sqrt{3}}\mathbf{j} + \frac{1}{\sqrt{3}}\mathbf{k} \right| = \sqrt{3 \left( \frac{1}{\sqrt{3}} \right)^2} = 1$ , the direction is  $\frac{1}{\sqrt{3}}\mathbf{i} + \frac{1}{\sqrt{3}}\mathbf{j} + \frac{1}{\sqrt{3}}\mathbf{k}$   
 $\Rightarrow \frac{1}{\sqrt{3}}\mathbf{i} + \frac{1}{\sqrt{3}}\mathbf{j} + \frac{1}{\sqrt{3}}\mathbf{k} = 1 \left( \frac{1}{\sqrt{3}}\mathbf{i} + \frac{1}{\sqrt{3}}\mathbf{j} + \frac{1}{\sqrt{3}}\mathbf{k} \right)$
31. (a)  $2\mathbf{i}$  (b)  $-\sqrt{3}\mathbf{k}$  (c)  $\frac{3}{10}\mathbf{j} + \frac{2}{5}\mathbf{k}$  (d)  $6\mathbf{i} - 2\mathbf{j} + 3\mathbf{k}$
32. (a)  $-7\mathbf{j}$  (b)  $-\frac{3\sqrt{2}}{5}\mathbf{i} - \frac{4\sqrt{2}}{5}\mathbf{k}$  (c)  $\frac{1}{4}\mathbf{i} - \frac{1}{3}\mathbf{j} - \mathbf{k}$  (d)  $\frac{a}{\sqrt{2}}\mathbf{i} + \frac{a}{\sqrt{3}}\mathbf{j} - \frac{a}{\sqrt{6}}\mathbf{k}$
33.  $|\mathbf{v}| = \sqrt{12^2 + 5^2} = \sqrt{169} = 13$ ;  $\frac{\mathbf{v}}{|\mathbf{v}|} = \frac{1}{13}\mathbf{v} = \frac{1}{13}(12\mathbf{i} - 5\mathbf{k}) \Rightarrow$  the desired vector is  $\frac{7}{13}(12\mathbf{i} - 5\mathbf{k})$
34.  $|\mathbf{v}| = \sqrt{\frac{1}{4} + \frac{1}{4} + \frac{1}{4}} = \frac{\sqrt{3}}{2}$ ;  $\frac{\mathbf{v}}{|\mathbf{v}|} = \frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k} \Rightarrow$  the desired vector is  $-3 \left( \frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k} \right)$   
 $= -\sqrt{3}\mathbf{i} + \sqrt{3}\mathbf{j} + \sqrt{3}\mathbf{k}$
35. (a)  $3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k} = 5\sqrt{2} \left( \frac{3}{5\sqrt{2}}\mathbf{i} + \frac{4}{5\sqrt{2}}\mathbf{j} - \frac{1}{\sqrt{2}}\mathbf{k} \right) \Rightarrow$  the direction is  $\frac{3}{5\sqrt{2}}\mathbf{i} + \frac{4}{5\sqrt{2}}\mathbf{j} - \frac{1}{\sqrt{2}}\mathbf{k}$   
 (b) the midpoint is  $\left( \frac{1}{2}, 3, \frac{5}{2} \right)$
36. (a)  $3\mathbf{i} - 6\mathbf{j} + 2\mathbf{k} = 7 \left( \frac{3}{7}\mathbf{i} - \frac{6}{7}\mathbf{j} + \frac{2}{7}\mathbf{k} \right) \Rightarrow$  the direction is  $\frac{3}{7}\mathbf{i} - \frac{6}{7}\mathbf{j} + \frac{2}{7}\mathbf{k}$   
 (b) the midpoint is  $\left( \frac{5}{2}, 1, 6 \right)$
37. (a)  $-\mathbf{i} - \mathbf{j} - \mathbf{k} = \sqrt{3} \left( -\frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k} \right) \Rightarrow$  the direction is  $-\frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k}$   
 (b) the midpoint is  $\left( \frac{5}{2}, \frac{7}{2}, \frac{9}{2} \right)$
38. (a)  $2\mathbf{i} - 2\mathbf{j} - 2\mathbf{k} = 2\sqrt{3} \left( \frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k} \right) \Rightarrow$  the direction is  $\frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k}$   
 (b) the midpoint is  $(1, -1, -1)$
39.  $\vec{AB} = (5-a)\mathbf{i} + (1-b)\mathbf{j} + (3-c)\mathbf{k} = \mathbf{i} + 4\mathbf{j} - 2\mathbf{k} \Rightarrow 5-a=1, 1-b=4, \text{ and } 3-c=-2 \Rightarrow a=4, b=-3, \text{ and } c=5 \Rightarrow$  A is the point  $(4, -3, 5)$
40.  $\vec{AB} = (a+2)\mathbf{i} + (b+3)\mathbf{j} + (c-6)\mathbf{k} = -7\mathbf{i} + 3\mathbf{j} + 8\mathbf{k} \Rightarrow a+2=-7, b+3=3, \text{ and } c-6=8 \Rightarrow a=-9, b=0, \text{ and } c=14 \Rightarrow$  B is the point  $(-9, 0, 14)$
41.  $2\mathbf{i} + \mathbf{j} = a(\mathbf{i} + \mathbf{j}) + b(\mathbf{i} - \mathbf{j}) = (a+b)\mathbf{i} + (a-b)\mathbf{j} \Rightarrow a+b=2 \text{ and } a-b=1 \Rightarrow 2a=3 \Rightarrow a=\frac{3}{2}$  and  $b=a-1=\frac{1}{2}$
42.  $\mathbf{i} - 2\mathbf{j} = a(2\mathbf{i} + 3\mathbf{j}) + b(\mathbf{i} + \mathbf{j}) = (2a+b)\mathbf{i} + (3a+b)\mathbf{j} \Rightarrow 2a+b=1 \text{ and } 3a+b=-2 \Rightarrow a=-3$  and  $b=1-2a=7 \Rightarrow \mathbf{u}_1 = a(2\mathbf{i} + 3\mathbf{j}) = -6\mathbf{i} - 9\mathbf{j}$  and  $\mathbf{u}_2 = b(\mathbf{i} + \mathbf{j}) = 7\mathbf{i} + 7\mathbf{j}$
43. If  $|x|$  is the magnitude of the x-component, then  $\cos 30^\circ = \frac{|x|}{|F|} \Rightarrow |x| = |F| \cos 30^\circ = (10) \left( \frac{\sqrt{3}}{2} \right) = 5\sqrt{3}$  lb  
 $\Rightarrow F_x = 5\sqrt{3}\mathbf{i}$ ;  
 if  $|y|$  is the magnitude of the y-component, then  $\sin 30^\circ = \frac{|y|}{|F|} \Rightarrow |y| = |F| \sin 30^\circ = (10) \left( \frac{1}{2} \right) = 5$  lb  $\Rightarrow F_y = 5\mathbf{j}$ .

44. If  $|x|$  is the magnitude of the x-component, then  $\cos 45^\circ = \frac{|x|}{|F|} \Rightarrow |x| = |F| \cos 45^\circ = (12) \left(\frac{\sqrt{2}}{2}\right) = 6\sqrt{2}$  lb  
 $\Rightarrow \mathbf{F}_x = -6\sqrt{2} \mathbf{i}$  (the negative sign is indicated by the diagram)  
 if  $|y|$  is the magnitude of the y-component, then  $\sin 45^\circ = \frac{|y|}{|F|} \Rightarrow |y| = |F| \sin 45^\circ = (12) \left(\frac{\sqrt{2}}{2}\right) = 6\sqrt{2}$  lb  
 $\Rightarrow \mathbf{F}_y = -6\sqrt{2} \mathbf{j}$  (the negative sign is indicated by the diagram)

45.  $25^\circ$  west of north is  $90^\circ + 25^\circ = 115^\circ$  north of east.  $800 \langle \cos 115^\circ, \sin 115^\circ \rangle \approx \langle -338.095, 725.046 \rangle$

46.  $10^\circ$  east of south is  $270^\circ + 10^\circ = 280^\circ$  "north" of east.  $600 \langle \cos 280^\circ, \sin 280^\circ \rangle \approx \langle 104.189, -590.885 \rangle$

47. (a) The tree is located at the tip of the vector  $\vec{OP} = (5 \cos 60^\circ) \mathbf{i} + (5 \sin 60^\circ) \mathbf{j} = \frac{5}{2} \mathbf{i} + \frac{5\sqrt{3}}{2} \mathbf{j} \Rightarrow P = \left(\frac{5}{2}, \frac{5\sqrt{3}}{2}\right)$

(b) The telephone pole is located at the point Q, which is the tip of the vector  $\vec{OP} + \vec{PQ}$   
 $= \left(\frac{5}{2} \mathbf{i} + \frac{5\sqrt{3}}{2} \mathbf{j}\right) + (10 \cos 315^\circ) \mathbf{i} + (10 \sin 315^\circ) \mathbf{j} = \left(\frac{5}{2} + \frac{10\sqrt{2}}{2}\right) \mathbf{i} + \left(\frac{5\sqrt{3}}{2} - \frac{10\sqrt{2}}{2}\right) \mathbf{j}$   
 $\Rightarrow Q = \left(\frac{5+10\sqrt{2}}{2}, \frac{5\sqrt{3}-10\sqrt{2}}{2}\right)$

48. Let  $t = \frac{q}{p+q}$  and  $s = \frac{p}{p+q}$ . Choose T on  $\vec{OP}_1$  so that  $\vec{TQ}$  is parallel to  $\vec{OP}_2$ , so that  $\triangle TP_1Q$  is similar to  $\triangle OP_1P_2$ . Then  $\frac{|\vec{OT}|}{|\vec{OP}_1|} = t \Rightarrow \vec{OT} = t \vec{OP}_1$  so that  $T = (tx_1, ty_1, tz_1)$ .

Also,  $\frac{|\vec{TQ}|}{|\vec{OP}_2|} = s \Rightarrow \vec{TQ} = s \vec{OP}_2 = s \langle x_2, y_2, z_2 \rangle$ .


Letting  $Q = (x, y, z)$ , we have that

$$\vec{TQ} = \langle x - tx_1, y - ty_1, z - tz_1 \rangle = s \langle x_2, y_2, z_2 \rangle$$

Thus  $x = tx_1 + sx_2, y = ty_1 + sy_2, z = tz_1 + sz_2$ .

(Note that if Q is the midpoint, then  $\frac{p}{q} = 1$  and  $t = s = \frac{1}{2}$

so that  $x = \frac{1}{2}x_1 + \frac{1}{2}x_2 = \frac{x_1+x_2}{2}, y = \frac{y_1+y_2}{2}, z = \frac{z_1+z_2}{2}$  so that this result agrees with the midpoint formula.)



49. (a) the midpoint of AB is  $M \left(\frac{5}{2}, \frac{5}{2}, 0\right)$  and  $\vec{CM} = \left(\frac{5}{2} - 1\right) \mathbf{i} + \left(\frac{5}{2} - 1\right) \mathbf{j} + (0 - 3) \mathbf{k} = \frac{3}{2} \mathbf{i} + \frac{3}{2} \mathbf{j} - 3 \mathbf{k}$

(b) the desired vector is  $\left(\frac{2}{3}\right) \vec{CM} = \frac{2}{3} \left(\frac{3}{2} \mathbf{i} + \frac{3}{2} \mathbf{j} - 3 \mathbf{k}\right) = \mathbf{i} + \mathbf{j} - 2 \mathbf{k}$

(c) the vector whose sum is the vector from the origin to C and the result of part (b) will terminate at the center of mass  $\Rightarrow$  the terminal point of  $(\mathbf{i} + \mathbf{j} + 3\mathbf{k}) + (\mathbf{i} + \mathbf{j} - 2\mathbf{k}) = 2\mathbf{i} + 2\mathbf{j} + \mathbf{k}$  is the point  $(2, 2, 1)$ , which is the location of the center of mass

50. The midpoint of AB is  $M \left(\frac{3}{2}, 0, \frac{5}{2}\right)$  and  $\left(\frac{2}{3}\right) \vec{CM} = \frac{2}{3} \left[\left(\frac{3}{2} + 1\right) \mathbf{i} + (0 - 2) \mathbf{j} + \left(\frac{5}{2} + 1\right) \mathbf{k}\right] = \frac{2}{3} \left(\frac{5}{2} \mathbf{i} - 2 \mathbf{j} + \frac{7}{2} \mathbf{k}\right)$   
 $= \frac{5}{3} \mathbf{i} - \frac{4}{3} \mathbf{j} + \frac{7}{3} \mathbf{k}$ . The terminal point of  $\left(\frac{5}{3} \mathbf{i} - \frac{4}{3} \mathbf{j} + \frac{7}{3} \mathbf{k}\right) + \vec{OC} = \left(\frac{5}{3} \mathbf{i} - \frac{4}{3} \mathbf{j} + \frac{7}{3} \mathbf{k}\right) + (-\mathbf{i} + 2\mathbf{j} - \mathbf{k})$   
 $= \frac{2}{3} \mathbf{i} + \frac{2}{3} \mathbf{j} + \frac{4}{3} \mathbf{k}$  is the point  $\left(\frac{2}{3}, \frac{2}{3}, \frac{4}{3}\right)$  which is the location of the intersection of the medians.

51. Without loss of generality we identify the vertices of the quadrilateral such that  $A(0, 0, 0)$ ,  $B(x_b, 0, 0)$ ,

$C(x_c, y_c, 0)$  and  $D(x_d, y_d, z_d) \Rightarrow$  the midpoint of AB is  $M_{AB} \left(\frac{x_b}{2}, 0, 0\right)$ , the midpoint of BC is

$M_{BC} \left(\frac{x_b+x_c}{2}, \frac{y_c}{2}, 0\right)$ , the midpoint of CD is  $M_{CD} \left(\frac{x_c+x_d}{2}, \frac{y_c+y_d}{2}, \frac{z_d}{2}\right)$  and the midpoint of AD is

$M_{AD} \left(\frac{x_d}{2}, \frac{y_d}{2}, \frac{z_d}{2}\right) \Rightarrow$  the midpoint of  $M_{AB}M_{CD}$  is  $\left(\frac{\frac{x_b}{2} + \frac{x_c+x_d}{2}}{2}, \frac{y_c+y_d}{4}, \frac{z_d}{4}\right)$  which is the same as the midpoint

of  $M_{AD}M_{BC} = \left(\frac{\frac{x_b+x_c}{2} + \frac{x_d}{2}}{2}, \frac{y_c+y_d}{4}, \frac{z_d}{4}\right)$ .

52. Let  $V_1, V_2, V_3, \dots, V_n$  be the vertices of a regular  $n$ -sided polygon and  $\mathbf{v}_i$  denote the vector from the center to  $V_i$  for  $i = 1, 2, 3, \dots, n$ . If  $\mathbf{S} = \sum_{i=1}^n \mathbf{v}_i$  and the polygon is rotated through an angle of  $\frac{i(2\pi)}{n}$  where  $i = 1, 2, 3, \dots, n$ , then  $\mathbf{S}$  would remain the same. Since the vector  $\mathbf{S}$  does not change with these rotations we conclude that  $\mathbf{S} = \mathbf{0}$ .
53. Without loss of generality we can coordinatize the vertices of the triangle such that  $A(0, 0)$ ,  $B(b, 0)$  and  $C(x_c, y_c) \Rightarrow$   $a$  is located at  $(\frac{b+x_c}{2}, \frac{y_c}{2})$ ,  $b$  is at  $(\frac{x_c}{2}, \frac{y_c}{2})$  and  $c$  is at  $(\frac{b}{2}, 0)$ . Therefore,  $\vec{Aa} = (\frac{b}{2} + \frac{x_c}{2})\mathbf{i} + (\frac{y_c}{2})\mathbf{j}$ ,  $\vec{Bb} = (\frac{x_c}{2} - b)\mathbf{i} + (\frac{y_c}{2})\mathbf{j}$ , and  $\vec{Cc} = (\frac{b}{2} - x_c)\mathbf{i} + (-y_c)\mathbf{j} \Rightarrow \vec{Aa} + \vec{Bb} + \vec{Cc} = \mathbf{0}$ .
54. Let  $\mathbf{u}$  be any unit vector in the plane. If  $\mathbf{u}$  is positioned so that its initial point is at the origin and terminal point is at  $(x, y)$ , then  $\mathbf{u}$  makes an angle  $\theta$  with  $\mathbf{i}$ , measured in the counter-clockwise direction. Since  $|\mathbf{u}| = 1$ , we have that  $x = \cos \theta$  and  $y = \sin \theta$ . Thus  $\mathbf{u} = \cos \theta \mathbf{i} + \sin \theta \mathbf{j}$ . Since  $\mathbf{u}$  was assumed to be any unit vector in the plane, this holds for every unit vector in the plane.

### 10.3 THE DOT PRODUCT

**NOTE:** In Exercises 1-8 below we calculate  $\text{proj}_{\mathbf{u}} \mathbf{v}$  as the vector  $(\frac{|\mathbf{u}| \cos \theta}{|\mathbf{v}|}) \mathbf{v}$ , so the scalar multiplier of  $\mathbf{v}$  is the number in column 5 divided by the number in column 2.

	$\mathbf{v} \cdot \mathbf{u}$	$ \mathbf{v} $	$ \mathbf{u} $	$\cos \theta$	$ \mathbf{u}  \cos \theta$	$\text{proj}_{\mathbf{u}} \mathbf{v}$
1.	-25	5	5	-1	-5	$-2\mathbf{i} + 4\mathbf{j} - \sqrt{5}\mathbf{k}$
2.	3	1	13	$\frac{3}{13}$	3	$3(\frac{3}{5}\mathbf{i} + \frac{4}{5}\mathbf{k})$
3.	25	15	5	$\frac{1}{3}$	$\frac{5}{3}$	$\frac{1}{9}(10\mathbf{i} + 11\mathbf{j} - 2\mathbf{k})$
4.	13	15	3	$\frac{13}{45}$	$\frac{13}{15}$	$\frac{13}{225}(2\mathbf{i} + 10\mathbf{j} - 11\mathbf{k})$
5.	2	$\sqrt{34}$	$\sqrt{3}$	$\frac{2}{\sqrt{3}\sqrt{34}}$	$\frac{2}{\sqrt{34}}$	$\frac{1}{17}(5\mathbf{j} - 3\mathbf{k})$
6.	$\sqrt{3} - \sqrt{2}$	$\sqrt{2}$	3	$\frac{\sqrt{3} - \sqrt{2}}{3\sqrt{2}}$	$\frac{\sqrt{3} - \sqrt{2}}{\sqrt{2}}$	$\frac{\sqrt{3} - \sqrt{2}}{2}(-\mathbf{i} + \mathbf{j})$
7.	$10 + \sqrt{17}$	$\sqrt{26}$	$\sqrt{21}$	$\frac{10 + \sqrt{17}}{\sqrt{546}}$	$\frac{10 + \sqrt{17}}{\sqrt{26}}$	$\frac{10 + \sqrt{17}}{\sqrt{26}}(5\mathbf{i} + \mathbf{j})$
8.	$\frac{1}{6}$	$\frac{\sqrt{30}}{6}$	$\frac{\sqrt{30}}{6}$	$\frac{1}{5}$	$\frac{1}{\sqrt{30}}$	$\frac{1}{5}\langle \frac{1}{\sqrt{2}}, \frac{1}{\sqrt{3}} \rangle$

$$9. \theta = \cos^{-1} \left( \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}| |\mathbf{v}|} \right) = \cos^{-1} \left( \frac{(2)(1) + (1)(2) + (0)(-1)}{\sqrt{2^2 + 1^2 + 0^2} \sqrt{1^2 + 2^2 + (-1)^2}} \right) = \cos^{-1} \left( \frac{4}{\sqrt{5}\sqrt{6}} \right) = \cos^{-1} \left( \frac{4}{\sqrt{30}} \right) \approx 0.75 \text{ rad}$$

$$10. \theta = \cos^{-1} \left( \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}| |\mathbf{v}|} \right) = \cos^{-1} \left( \frac{(2)(3) + (-2)(0) + (1)(4)}{\sqrt{2^2 + (-2)^2 + 1^2} \sqrt{3^2 + 0^2 + 4^2}} \right) = \cos^{-1} \left( \frac{10}{\sqrt{9}\sqrt{25}} \right) = \cos^{-1} \left( \frac{2}{3} \right) \approx 0.84 \text{ rad}$$

$$11. \theta = \cos^{-1} \left( \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}| |\mathbf{v}|} \right) = \cos^{-1} \left( \frac{(\sqrt{3})(\sqrt{3}) + (-7)(1) + (0)(-2)}{\sqrt{(\sqrt{3})^2 + (-7)^2 + 0^2} \sqrt{(\sqrt{3})^2 + (1)^2 + (-2)^2}} \right) = \cos^{-1} \left( \frac{3-7}{\sqrt{32}\sqrt{8}} \right) \\ = \cos^{-1} \left( \frac{-1}{\sqrt{26}} \right) \approx 1.77 \text{ rad}$$

12.  $\theta = \cos^{-1} \left( \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}| |\mathbf{v}|} \right) = \cos^{-1} \left( \frac{(1)(-1) + (\sqrt{2})(1) + (-\sqrt{2})(1)}{\sqrt{(1)^2 + (\sqrt{2})^2 + (-\sqrt{2})^2} \sqrt{(-1)^2 + (1)^2 + (1)^2}} \right) = \cos^{-1} \left( \frac{-1}{\sqrt{5} \sqrt{3}} \right)$   
 $= \cos^{-1} \left( \frac{-1}{\sqrt{15}} \right) \approx 1.83 \text{ rad}$
13.  $\vec{AB} = \langle 3, 1 \rangle$ ,  $\vec{BC} = \langle -1, -3 \rangle$ , and  $\vec{AC} = \langle 2, -2 \rangle$ .  $\vec{BA} = \langle -3, -1 \rangle$ ,  $\vec{CB} = \langle 1, 3 \rangle$ ,  $\vec{CA} = \langle -2, 2 \rangle$ .  
 $|\vec{AB}| = |\vec{BA}| = \sqrt{10}$ ,  $|\vec{BC}| = |\vec{CB}| = \sqrt{10}$ ,  $|\vec{AC}| = |\vec{CA}| = 2\sqrt{2}$ ,  
 Angle at A =  $\cos^{-1} \left( \frac{\vec{AB} \cdot \vec{AC}}{|\vec{AB}| |\vec{AC}|} \right) = \cos^{-1} \left( \frac{3(2) + 1(-2)}{(\sqrt{10})(2\sqrt{2})} \right) = \cos^{-1} \left( \frac{1}{\sqrt{5}} \right) \approx 63.435^\circ$   
 Angle at B =  $\cos^{-1} \left( \frac{\vec{BC} \cdot \vec{BA}}{|\vec{BC}| |\vec{BA}|} \right) = \cos^{-1} \left( \frac{(-1)(-3) + (-3)(-1)}{(\sqrt{10})(\sqrt{10})} \right) = \cos^{-1} \left( \frac{3}{5} \right) \approx 53.130^\circ$ , and  
 Angle at C =  $\cos^{-1} \left( \frac{\vec{CB} \cdot \vec{CA}}{|\vec{CB}| |\vec{CA}|} \right) = \cos^{-1} \left( \frac{1(-2) + 3(2)}{(\sqrt{10})(2\sqrt{2})} \right) = \cos^{-1} \left( \frac{1}{\sqrt{5}} \right) \approx 63.435^\circ$
14.  $\vec{AC} = \langle 2, 4 \rangle$  and  $\vec{BD} = \langle 4, -2 \rangle$ .  $\vec{AC} \cdot \vec{BD} = 2(4) + 4(-2) = 0$ , so the angle measures are all  $90^\circ$ .
15. (a)  $\cos \alpha = \frac{\mathbf{i} \cdot \mathbf{v}}{|\mathbf{i}| |\mathbf{v}|} = \frac{a}{|\mathbf{v}|}$ ,  $\cos \beta = \frac{\mathbf{j} \cdot \mathbf{v}}{|\mathbf{j}| |\mathbf{v}|} = \frac{b}{|\mathbf{v}|}$ ,  $\cos \gamma = \frac{\mathbf{k} \cdot \mathbf{v}}{|\mathbf{k}| |\mathbf{v}|} = \frac{c}{|\mathbf{v}|}$  and  
 $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = \left( \frac{a}{|\mathbf{v}|} \right)^2 + \left( \frac{b}{|\mathbf{v}|} \right)^2 + \left( \frac{c}{|\mathbf{v}|} \right)^2 = \frac{a^2 + b^2 + c^2}{|\mathbf{v}|^2} = \frac{|\mathbf{v}|^2}{|\mathbf{v}|^2} = 1$   
 (b)  $|\mathbf{v}| = 1 \Rightarrow \cos \alpha = \frac{a}{|\mathbf{v}|} = a$ ,  $\cos \beta = \frac{b}{|\mathbf{v}|} = b$  and  $\cos \gamma = \frac{c}{|\mathbf{v}|} = c$  are the direction cosines of  $\mathbf{v}$
16.  $\mathbf{u} = 10\mathbf{i} + 2\mathbf{k}$  is parallel to the pipe in the north direction and  $\mathbf{v} = 10\mathbf{j} + \mathbf{k}$  is parallel to the pipe in the east direction. The angle between the two pipes is  $\theta = \cos^{-1} \left( \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{u}| |\mathbf{v}|} \right) = \cos^{-1} \left( \frac{2}{\sqrt{104} \sqrt{101}} \right) \approx 1.55 \text{ rad} \approx 88.88^\circ$ .
17. The sum of two vectors of equal length is *always* orthogonal to their difference, as we can see from the equation  
 $(\mathbf{v}_1 + \mathbf{v}_2) \cdot (\mathbf{v}_1 - \mathbf{v}_2) = \mathbf{v}_1 \cdot \mathbf{v}_1 + \mathbf{v}_2 \cdot \mathbf{v}_1 - \mathbf{v}_1 \cdot \mathbf{v}_2 - \mathbf{v}_2 \cdot \mathbf{v}_2 = |\mathbf{v}_1|^2 - |\mathbf{v}_2|^2 = 0$
18.  $\vec{CA} \cdot \vec{CB} = (-\mathbf{v} + (-\mathbf{u})) \cdot (-\mathbf{v} + \mathbf{u}) = \mathbf{v} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{v} - \mathbf{u} \cdot \mathbf{u} = |\mathbf{v}|^2 - |\mathbf{u}|^2 = 0$  because  $|\mathbf{u}| = |\mathbf{v}|$  since both equal the radius of the circle. Therefore,  $\vec{CA}$  and  $\vec{CB}$  are orthogonal.
19. Let  $\mathbf{u}$  and  $\mathbf{v}$  be the sides of a rhombus  $\Rightarrow$  the diagonals are  $\mathbf{d}_1 = \mathbf{u} + \mathbf{v}$  and  $\mathbf{d}_2 = -\mathbf{u} + \mathbf{v}$   
 $\Rightarrow \mathbf{d}_1 \cdot \mathbf{d}_2 = (\mathbf{u} + \mathbf{v}) \cdot (-\mathbf{u} + \mathbf{v}) = -\mathbf{u} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{u} + \mathbf{v} \cdot \mathbf{v} = |\mathbf{v}|^2 - |\mathbf{u}|^2 = 0$  because  $|\mathbf{u}| = |\mathbf{v}|$ , since a rhombus has equal sides.
20. Suppose the diagonals of a rectangle are perpendicular, and let  $\mathbf{u}$  and  $\mathbf{v}$  be the sides of a rectangle  $\Rightarrow$  the diagonals are  $\mathbf{d}_1 = \mathbf{u} + \mathbf{v}$  and  $\mathbf{d}_2 = -\mathbf{u} + \mathbf{v}$ . Since the diagonals are perpendicular we have  $\mathbf{d}_1 \cdot \mathbf{d}_2 = 0$   
 $\Leftrightarrow (\mathbf{u} + \mathbf{v}) \cdot (-\mathbf{u} + \mathbf{v}) = -\mathbf{u} \cdot \mathbf{u} + \mathbf{u} \cdot \mathbf{v} - \mathbf{v} \cdot \mathbf{u} + \mathbf{v} \cdot \mathbf{v} = 0 \Leftrightarrow |\mathbf{v}|^2 - |\mathbf{u}|^2 = 0 \Leftrightarrow (|\mathbf{v}| + |\mathbf{u}|)(|\mathbf{v}| - |\mathbf{u}|) = 0$   
 $\Leftrightarrow (|\mathbf{v}| + |\mathbf{u}|) = 0$  which is not possible, or  $(|\mathbf{v}| - |\mathbf{u}|) = 0$  which is equivalent to  $|\mathbf{v}| = |\mathbf{u}| \Rightarrow$  the rectangle is a square.
21. Clearly the diagonals of a rectangle are equal in length. What is not as obvious is the statement that equal diagonals happen only in a rectangle. We show this is true by letting the adjacent sides of a parallelogram be the vectors  $(v_1\mathbf{i} + v_2\mathbf{j})$  and  $(u_1\mathbf{i} + u_2\mathbf{j})$ . The equal diagonals of the parallelogram are  $\mathbf{d}_1 = (v_1\mathbf{i} + v_2\mathbf{j}) + (u_1\mathbf{i} + u_2\mathbf{j})$  and  $\mathbf{d}_2 = (v_1\mathbf{i} + v_2\mathbf{j}) - (u_1\mathbf{i} + u_2\mathbf{j})$ . Hence  $|\mathbf{d}_1| = |\mathbf{d}_2| \Rightarrow |(v_1\mathbf{i} + v_2\mathbf{j}) + (u_1\mathbf{i} + u_2\mathbf{j})| = |(v_1\mathbf{i} + v_2\mathbf{j}) - (u_1\mathbf{i} + u_2\mathbf{j})|$   
 $\Rightarrow |(v_1 + u_1)\mathbf{i} + (v_2 + u_2)\mathbf{j}| = |(v_1 - u_1)\mathbf{i} + (v_2 - u_2)\mathbf{j}| \Rightarrow \sqrt{(v_1 + u_1)^2 + (v_2 + u_2)^2} = \sqrt{(v_1 - u_1)^2 + (v_2 - u_2)^2}$   
 $\Rightarrow v_1^2 + 2v_1u_1 + u_1^2 + v_2^2 + 2v_2u_2 + u_2^2 = v_1^2 - 2v_1u_1 + u_1^2 + v_2^2 - 2v_2u_2 + u_2^2 \Rightarrow 2(v_1u_1 + v_2u_2)$

622 Chapter 10 Vectors and the Geometry of Space

$= -2(v_1u_1 + v_2u_2) \Rightarrow v_1u_1 + v_2u_2 = 0 \Rightarrow (v_1\mathbf{i} + v_2\mathbf{j}) \cdot (u_1\mathbf{i} + u_2\mathbf{j}) = 0 \Rightarrow$  the vectors  $(v_1\mathbf{i} + v_2\mathbf{j})$  and  $(u_1\mathbf{i} + u_2\mathbf{j})$  are perpendicular and the parallelogram must be a rectangle.

22. If  $|\mathbf{u}| = |\mathbf{v}|$  and  $\mathbf{u} + \mathbf{v}$  is the indicated diagonal, then  $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{u} = \mathbf{u} \cdot \mathbf{u} + \mathbf{v} \cdot \mathbf{u} = |\mathbf{u}|^2 + \mathbf{v} \cdot \mathbf{u} = \mathbf{u} \cdot \mathbf{v} + |\mathbf{v}|^2 = \mathbf{u} \cdot \mathbf{v} + \mathbf{v} \cdot \mathbf{v} = (\mathbf{u} + \mathbf{v}) \cdot \mathbf{v} \Rightarrow$  the angle  $\cos^{-1} \left( \frac{(\mathbf{u} + \mathbf{v}) \cdot \mathbf{u}}{|\mathbf{u} + \mathbf{v}| |\mathbf{u}|} \right)$  between the diagonal and  $\mathbf{u}$  and the angle  $\cos^{-1} \left( \frac{(\mathbf{u} + \mathbf{v}) \cdot \mathbf{v}}{|\mathbf{u} + \mathbf{v}| |\mathbf{v}|} \right)$  between the diagonal and  $\mathbf{v}$  are equal because the inverse cosine function is one-to-one. Therefore, the diagonal bisects the angle between  $\mathbf{u}$  and  $\mathbf{v}$ .


23. horizontal component:  $1200 \cos(8^\circ) \approx 1188$  ft/s; vertical component:  $1200 \sin(8^\circ) \approx 167$  ft/s

24.  $|\mathbf{w}| \cos(33^\circ - 15^\circ) = 2.5$  lb, so  $|\mathbf{w}| = \frac{2.5 \text{ lb}}{\cos 18^\circ}$ . Then  $\mathbf{w} = \frac{2.5 \text{ lb}}{\cos 18^\circ} \langle \cos 33^\circ, \sin 33^\circ \rangle \approx \langle 2.205, 1.432 \rangle$

25. (a) Since  $|\cos \theta| \leq 1$ , we have  $|\mathbf{u} \cdot \mathbf{v}| = |\mathbf{u}| |\mathbf{v}| |\cos \theta| \leq |\mathbf{u}| |\mathbf{v}| (1) = |\mathbf{u}| |\mathbf{v}|$ .

(b) We have equality precisely when  $|\cos \theta| = 1$  or when one or both of  $\mathbf{u}$  and  $\mathbf{v}$  is  $\mathbf{0}$ . In the case of nonzero vectors, we have equality when  $\theta = 0$  or  $\pi$ , i.e., when the vectors are parallel.

26.  $(x\mathbf{i} + y\mathbf{j}) \cdot \mathbf{v} = |x\mathbf{i} + y\mathbf{j}| |\mathbf{v}| \cos \theta \leq 0$  when  $\frac{\pi}{2} \leq \theta \leq \pi$ . This means  $(x, y)$  has to be a point whose position vector makes an angle with  $\mathbf{v}$  that is a right angle or bigger.




27.  $\mathbf{v} \cdot \mathbf{u}_1 = (a\mathbf{u}_1 + b\mathbf{u}_2) \cdot \mathbf{u}_1 = a\mathbf{u}_1 \cdot \mathbf{u}_1 + b\mathbf{u}_2 \cdot \mathbf{u}_1 = a|\mathbf{u}_1|^2 + b(\mathbf{u}_2 \cdot \mathbf{u}_1) = a(1)^2 + b(0) = a$

28. No,  $v_1$  need not equal  $v_2$ . For example,  $\mathbf{i} + \mathbf{j} \neq \mathbf{i} + 2\mathbf{j}$  but  $\mathbf{i} \cdot (\mathbf{i} + \mathbf{j}) = \mathbf{i} \cdot \mathbf{i} + \mathbf{i} \cdot \mathbf{j} = 1 + 0 = 1$  and  $\mathbf{i} \cdot (\mathbf{i} + 2\mathbf{j}) = \mathbf{i} \cdot \mathbf{i} + 2\mathbf{i} \cdot \mathbf{j} = 1 + 2 \cdot 0 = 1$ .

29.  $P(x_1, y_1) = P(x_1, \frac{c}{b} - \frac{a}{b}x_1)$  and  $Q(x_2, y_2) = Q(x_2, \frac{c}{b} - \frac{a}{b}x_2)$  are any two points  $P$  and  $Q$  on the line with  $b \neq 0$   
 $\Rightarrow \vec{PQ} = (x_2 - x_1)\mathbf{i} + \frac{a}{b}(x_1 - x_2)\mathbf{j} \Rightarrow \vec{PQ} \cdot \mathbf{v} = [(x_2 - x_1)\mathbf{i} + \frac{a}{b}(x_1 - x_2)\mathbf{j}] \cdot (a\mathbf{i} + b\mathbf{j}) = a(x_2 - x_1) + b(\frac{a}{b})(x_1 - x_2) = 0 \Rightarrow \mathbf{v}$  is perpendicular to  $\vec{PQ}$  for  $b \neq 0$ . If  $b = 0$ , then  $\mathbf{v} = a\mathbf{i}$  is perpendicular to the vertical line  $ax = c$ .  
 Alternatively, the slope of  $\mathbf{v}$  is  $\frac{b}{a}$  and the slope of the line  $ax + by = c$  is  $-\frac{a}{b}$ , so the slopes are negative reciprocals  
 $\Rightarrow$  the vector  $\mathbf{v}$  and the line are perpendicular.

30. The slope of  $\mathbf{v}$  is  $\frac{b}{a}$  and the slope of  $bx - ay = c$  is  $\frac{b}{a}$ , provided that  $a \neq 0$ . If  $a = 0$ , then  $\mathbf{v} = b\mathbf{j}$  is parallel to the vertical line  $bx = c$ . In either case, the vector  $\mathbf{v}$  is parallel to the line  $ax - by = c$ .


31.  $\mathbf{v} = \mathbf{i} + 2\mathbf{j}$  is perpendicular to the line  $x + 2y = c$ ;  
 $P(2, 1)$  on the line  $\Rightarrow 2 + 2 = c \Rightarrow x + 2y = 4$




$$8. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \frac{3}{2} & -\frac{1}{2} & 1 \\ 1 & 1 & 2 \end{vmatrix} = -2\mathbf{i} - 2\mathbf{j} + 2\mathbf{k} \Rightarrow \text{length} = 2\sqrt{3} \text{ and the direction is } -\frac{1}{\sqrt{3}}\mathbf{i} - \frac{1}{\sqrt{3}}\mathbf{j} + \frac{1}{\sqrt{3}}\mathbf{k}$$

$$\mathbf{v} \times \mathbf{u} = -(\mathbf{u} \times \mathbf{v}) = -(-2\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}) \Rightarrow \text{length} = 2\sqrt{3} \text{ and the direction is } \frac{1}{\sqrt{3}}\mathbf{i} + \frac{1}{\sqrt{3}}\mathbf{j} - \frac{1}{\sqrt{3}}\mathbf{k}$$


$$9. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{vmatrix} = \mathbf{k}$$




$$10. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & -1 \\ 0 & 1 & 0 \end{vmatrix} = \mathbf{i} + \mathbf{k}$$




$$11. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & -1 \\ 0 & 1 & 1 \end{vmatrix} = \mathbf{i} - \mathbf{j} + \mathbf{k}$$




$$12. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -1 & 0 \\ 1 & 2 & 0 \end{vmatrix} = 5\mathbf{k}$$



$$13. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 0 \\ 1 & -1 & 0 \end{vmatrix} = -2\mathbf{k}$$



$$14. \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & 2 \\ 1 & 0 & 0 \end{vmatrix} = 2\mathbf{j} - \mathbf{k}$$



$$15. (a) \vec{PQ} \times \vec{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & -3 \\ -1 & 3 & -1 \end{vmatrix} = 8\mathbf{i} + 4\mathbf{j} + 4\mathbf{k} \Rightarrow \text{Area} = \frac{1}{2} |\vec{PQ} \times \vec{PR}| = \frac{1}{2} \sqrt{64 + 16 + 16} = 2\sqrt{6}$$

$$(b) \mathbf{u} = \pm \frac{\vec{PQ} \times \vec{PR}}{|\vec{PQ} \times \vec{PR}|} = \pm \frac{1}{\sqrt{6}} (2\mathbf{i} + \mathbf{j} + \mathbf{k})$$

$$16. (a) \vec{PQ} \times \vec{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 2 \\ 2 & -2 & 0 \end{vmatrix} = 4\mathbf{i} + 4\mathbf{j} - 2\mathbf{k} \Rightarrow \text{Area} = \frac{1}{2} |\vec{PQ} \times \vec{PR}| = \frac{1}{2} \sqrt{16 + 16 + 4} = 3$$

$$(b) \mathbf{u} = \pm \frac{\vec{PQ} \times \vec{PR}}{|\vec{PQ} \times \vec{PR}|} = \pm \frac{1}{3} (2\mathbf{i} + 2\mathbf{j} - \mathbf{k})$$

$$17. (a) \vec{PQ} \times \vec{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{vmatrix} = -\mathbf{i} + \mathbf{j} \Rightarrow \text{Area} = \frac{1}{2} |\vec{PQ} \times \vec{PR}| = \frac{1}{2} \sqrt{1 + 1} = \frac{\sqrt{2}}{2}$$

$$(b) \mathbf{u} = \pm \frac{\vec{PQ} \times \vec{PR}}{|\vec{PQ} \times \vec{PR}|} = \pm \frac{1}{\sqrt{2}} (-\mathbf{i} + \mathbf{j}) = \pm \frac{1}{\sqrt{2}} (\mathbf{i} - \mathbf{j})$$

$$18. (a) \vec{PQ} \times \vec{PR} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & -1 & -1 \\ 1 & 0 & -2 \end{vmatrix} = 2\mathbf{i} + 3\mathbf{j} + \mathbf{k} \Rightarrow \text{Area} = \frac{1}{2} |\vec{PQ} \times \vec{PR}| = \frac{1}{2} \sqrt{4 + 9 + 1} = \frac{\sqrt{14}}{2}$$

$$(b) \mathbf{u} = \pm \frac{\vec{PQ} \times \vec{PR}}{|\vec{PQ} \times \vec{PR}|} = \pm \frac{1}{\sqrt{14}} (2\mathbf{i} + 3\mathbf{j} + \mathbf{k})$$

$$19. \text{ If } \mathbf{u} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}, \mathbf{v} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}, \text{ and } \mathbf{w} = c_1\mathbf{i} + c_2\mathbf{j} + c_3\mathbf{k}, \text{ then } \mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix},$$

$$\mathbf{v} \cdot (\mathbf{w} \times \mathbf{u}) = \begin{vmatrix} b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \\ a_1 & a_2 & a_3 \end{vmatrix} \text{ and } \mathbf{w} \cdot (\mathbf{u} \times \mathbf{v}) = \begin{vmatrix} c_1 & c_2 & c_3 \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} \text{ which all have the same value, since the}$$

interchanging of two pair of rows in a determinant does not change its value  $\Rightarrow$  the volume is

$$|(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}| = \text{abs} \begin{vmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{vmatrix} = 8$$

$$20. |(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}| = \text{abs} \begin{vmatrix} 1 & -1 & 1 \\ 2 & 1 & -2 \\ -1 & 2 & -1 \end{vmatrix} = 4 \text{ (for details about verification, see Exercise 19)}$$

$$21. |(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}| = \text{abs} \begin{vmatrix} 2 & 1 & 0 \\ 2 & -1 & 1 \\ 1 & 0 & 2 \end{vmatrix} = |-7| = 7 \text{ (for details about verification, see Exercise 19)}$$

$$22. |(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}| = \text{abs} \begin{vmatrix} 1 & 1 & -2 \\ -1 & 0 & -1 \\ 2 & 4 & -2 \end{vmatrix} = 8 \text{ (for details about verification, see Exercise 19)}$$

$$23. (a) \mathbf{u} \cdot \mathbf{v} = -6, \mathbf{u} \cdot \mathbf{w} = -81, \mathbf{v} \cdot \mathbf{w} = 18 \Rightarrow \text{none}$$

$$(b) \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 5 & -1 & 1 \\ 0 & 1 & -5 \end{vmatrix} \neq \mathbf{0}, \mathbf{u} \times \mathbf{w} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 5 & -1 & 1 \\ -15 & 3 & -3 \end{vmatrix} = \mathbf{0}, \mathbf{v} \times \mathbf{w} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 1 & -5 \\ -15 & 3 & -3 \end{vmatrix} \neq \mathbf{0}$$

$\Rightarrow \mathbf{u}$  and  $\mathbf{w}$  are parallel

$$24. (a) \mathbf{u} \cdot \mathbf{v} = 0, \mathbf{u} \times \mathbf{w} = \mathbf{0}, \mathbf{u} \cdot \mathbf{r} = -3\pi, \mathbf{v} \cdot \mathbf{w} = 0, \mathbf{v} \cdot \mathbf{r} = 0, \mathbf{w} \cdot \mathbf{r} = 0 \Rightarrow \mathbf{u} \perp \mathbf{v}, \mathbf{u} \perp \mathbf{w}, \mathbf{v} \perp \mathbf{w}, \mathbf{v} \perp \mathbf{r} \text{ and } \mathbf{w} \perp \mathbf{r}$$

$$(b) \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & -1 \\ -1 & 1 & 1 \end{vmatrix} \neq \mathbf{0}, \mathbf{u} \times \mathbf{w} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & -1 \\ 1 & 0 & 1 \end{vmatrix} \neq \mathbf{0}, \mathbf{u} \times \mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & -1 \\ -\frac{\pi}{2} & -\pi & \frac{\pi}{2} \end{vmatrix} = \mathbf{0}$$

$$\mathbf{v} \times \mathbf{w} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 1 & 1 \\ 1 & 0 & 1 \end{vmatrix} \neq \mathbf{0}, \mathbf{v} \times \mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 1 & 1 \\ -\frac{\pi}{2} & -\pi & \frac{\pi}{2} \end{vmatrix} \neq \mathbf{0}, \mathbf{w} \times \mathbf{r} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 0 & 1 \\ -\frac{\pi}{2} & -\pi & \frac{\pi}{2} \end{vmatrix} \neq \mathbf{0}$$

$\Rightarrow \mathbf{u}$  and  $\mathbf{r}$  are parallel

$$25. |\vec{PQ} \times \mathbf{F}| = |\vec{PQ}| |\mathbf{F}| \sin(60^\circ) = \frac{2}{3} \cdot 30 \cdot \frac{\sqrt{3}}{2} \text{ ft} \cdot \text{lb} = 10\sqrt{3} \text{ ft} \cdot \text{lb}$$

$$26. |\vec{PQ} \times \mathbf{F}| = |\vec{PQ}| |\mathbf{F}| \sin(135^\circ) = \frac{2}{3} \cdot 30 \cdot \frac{\sqrt{2}}{2} \text{ ft} \cdot \text{lb} = 10\sqrt{2} \text{ ft} \cdot \text{lb}$$

$$27. (a) \text{ true, } |\mathbf{u}| = \sqrt{a_1^2 + a_2^2 + a_3^2} = \sqrt{\mathbf{u} \cdot \mathbf{u}}$$

$$(b) \text{ not always true, } \mathbf{u} \cdot \mathbf{u} = |\mathbf{u}|^2$$

$$(c) \text{ true, } \mathbf{u} \times \mathbf{0} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ 0 & 0 & 0 \end{vmatrix} = 0\mathbf{i} + 0\mathbf{j} + 0\mathbf{k} = \mathbf{0} \text{ and } \mathbf{0} \times \mathbf{u} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & 0 \\ a_1 & a_2 & a_3 \end{vmatrix} = 0\mathbf{i} + 0\mathbf{j} + 0\mathbf{k} = \mathbf{0}$$

$$(d) \text{ true, } \mathbf{u} \times (-\mathbf{u}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ -a_1 & -a_2 & -a_3 \end{vmatrix} = (-a_2a_3 + a_2a_3)\mathbf{i} - (-a_1a_3 + a_1a_3)\mathbf{j} + (-a_1a_2 + a_1a_2)\mathbf{k} = \mathbf{0}$$

$$(e) \text{ not always true, } \mathbf{i} \times \mathbf{j} = \mathbf{k} \neq -\mathbf{k} = \mathbf{j} \times \mathbf{i} \text{ for example}$$

$$(f) \text{ true, distributive property of the cross product}$$

$$(g) \text{ true, } (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{v} = \mathbf{u} \cdot (\mathbf{v} \times \mathbf{v}) = \mathbf{u} \cdot \mathbf{0} = 0$$

$$(h) \text{ true, the volume of a parallelepiped with } \mathbf{u}, \mathbf{v}, \text{ and } \mathbf{w} \text{ along the three edges is } (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w} = (\mathbf{v} \times \mathbf{w}) \cdot \mathbf{u} = \mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}),$$

since the dot product is commutative.

$$28. (a) \text{ true, } \mathbf{u} \cdot \mathbf{v} = a_1b_1 + a_2b_2 + a_3b_3 = b_1a_1 + b_2a_2 + b_3a_3 = \mathbf{v} \cdot \mathbf{u}$$

$$(b) \text{ true, } \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = - \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ b_1 & b_2 & b_3 \\ a_1 & a_2 & a_3 \end{vmatrix} = -(\mathbf{v} \times \mathbf{u})$$

$$(c) \text{ true, } (-\mathbf{u}) \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a_1 & -a_2 & -a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = - \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = -(\mathbf{u} \times \mathbf{v})$$

$$(d) \text{ true, } (c\mathbf{u}) \cdot \mathbf{v} = (ca_1)b_1 + (ca_2)b_2 + (ca_3)b_3 = a_1(cb_1) + a_2(cb_2) + a_3(cb_3) = \mathbf{u} \cdot (c\mathbf{v}) = c(a_1b_1 + a_2b_2 + a_3b_3) = c(\mathbf{u} \cdot \mathbf{v})$$

$$(e) \text{ true, } c(\mathbf{u} \times \mathbf{v}) = c \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ ca_1 & ca_2 & ca_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = (c\mathbf{u}) \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ cb_1 & cb_2 & cb_3 \end{vmatrix} = \mathbf{u} \times (c\mathbf{v})$$

$$(f) \text{ true, } \mathbf{u} \cdot \mathbf{u} = a_1^2 + a_2^2 + a_3^2 = (\sqrt{a_1^2 + a_2^2 + a_3^2})^2 = |\mathbf{u}|^2$$

$$(g) \text{ true, } (\mathbf{u} \times \mathbf{u}) \cdot \mathbf{u} = \mathbf{0} \cdot \mathbf{u} = 0$$

$$(h) \text{ true, } \mathbf{u} \times \mathbf{v} \perp \mathbf{u} \text{ and } \mathbf{u} \times \mathbf{v} \perp \mathbf{v} \Rightarrow (\mathbf{u} \times \mathbf{v}) \cdot \mathbf{u} = \mathbf{v} \cdot (\mathbf{u} \times \mathbf{v}) = 0$$


$$29. (a) \text{ proj}_{\mathbf{v}} \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{|\mathbf{v}|} \right) \frac{\mathbf{v}}{|\mathbf{v}|} \quad (b) \pm(\mathbf{u} \times \mathbf{v}) \quad (c) \pm((\mathbf{u} \times \mathbf{v}) \times \mathbf{w}) \quad (d) |(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w}|$$

$$30. (a) (\mathbf{u} \times \mathbf{v}) \times (\mathbf{u} \times \mathbf{w})$$


$$(b) (\mathbf{u} + \mathbf{v}) \times (\mathbf{u} - \mathbf{v}) = (\mathbf{u} + \mathbf{v}) \times \mathbf{u} - (\mathbf{u} + \mathbf{v}) \times \mathbf{v} = \mathbf{u} \times \mathbf{u} + \mathbf{v} \times \mathbf{u} - \mathbf{u} \times \mathbf{v} - \mathbf{v} \times \mathbf{v} = \mathbf{0} + \mathbf{v} \times \mathbf{u} - \mathbf{u} \times \mathbf{v} - \mathbf{0} = 2(\mathbf{v} \times \mathbf{u}), \text{ or simply } \mathbf{u} \times \mathbf{v}$$

$$(c) |\mathbf{u}| \frac{\mathbf{v}}{|\mathbf{v}|} \quad (d) |\mathbf{u} \times \mathbf{w}|$$

19. The direction  $\vec{PQ} = -2\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$  and  $P(2, 0, 2)$   
 $\Rightarrow x = 2 - 2t, y = 2t, z = 2 - 2t$ , where  $0 \leq t \leq 1$



20. The direction  $\vec{PQ} = -\mathbf{i} + 3\mathbf{j} + \mathbf{k}$  and  $P(1, 0, -1)$   
 $\Rightarrow x = 1 - t, y = 3t, z = -1 + t$ , where  $0 \leq t \leq 1$



21.  $3(x - 0) + (-2)(y - 2) + (-1)(z + 1) = 0 \Rightarrow 3x - 2y - z = -3$

22.  $3(x - 1) + (1)(y + 1) + (1)(z - 3) = 0 \Rightarrow 3x + y + z = 5$

23.  $\vec{PQ} = \mathbf{i} - \mathbf{j} + 3\mathbf{k}, \vec{PS} = -\mathbf{i} - 3\mathbf{j} + 2\mathbf{k} \Rightarrow \vec{PQ} \times \vec{PS} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -1 & 3 \\ -1 & -3 & 2 \end{vmatrix} = 7\mathbf{i} - 5\mathbf{j} - 4\mathbf{k}$  is normal to the plane  
 $\Rightarrow 7(x - 2) + (-5)(y - 0) + (-4)(z - 2) = 0 \Rightarrow 7x - 5y - 4z = 6$

24.  $\vec{PQ} = -\mathbf{i} + \mathbf{j} + 2\mathbf{k}, \vec{PS} = -3\mathbf{i} + 2\mathbf{j} + 3\mathbf{k} \Rightarrow \vec{PQ} \times \vec{PS} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 1 & 2 \\ -3 & 2 & 3 \end{vmatrix} = -\mathbf{i} - 3\mathbf{j} + \mathbf{k}$  is normal to the plane  
 $\Rightarrow (-1)(x - 1) + (-3)(y - 5) + (1)(z - 7) = 0 \Rightarrow x + 3y - z = 9$

25.  $\mathbf{n} = \mathbf{i} + 3\mathbf{j} + 4\mathbf{k}, P(2, 4, 5) = (1)(x - 2) + (3)(y - 4) + (4)(z - 5) = 0 \Rightarrow x + 3y + 4z = 34$

26.  $\mathbf{n} = \mathbf{i} - 2\mathbf{j} + \mathbf{k}, P(1, -2, 1) = (1)(x - 1) + (-2)(y + 2) + (1)(z - 1) = 0 \Rightarrow x - 2y + z = 6$

27.  $\begin{cases} x = 2t + 1 = s + 2 \\ y = 3t + 2 = 2s + 4 \end{cases} \Rightarrow \begin{cases} 2t - s = 1 \\ 3t - 2s = 2 \end{cases} \Rightarrow \begin{cases} 4t - 2s = 2 \\ 3t - 2s = 2 \end{cases} \Rightarrow t = 0 \text{ and } s = -1; \text{ then } z = 4t + 3 = -4s - 1$   
 $\Rightarrow 4(0) + 3 = (-4)(-1) - 1$  is satisfied  $\Rightarrow$  the lines do intersect when  $t = 0$  and  $s = -1 \Rightarrow$  the point of intersection is  $x = 1, y = 2$ , and  $z = 3$  or  $P(1, 2, 3)$ . A vector normal to the plane determined by these lines is

$\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 3 & 4 \\ 1 & 2 & -4 \end{vmatrix} = -20\mathbf{i} + 12\mathbf{j} + \mathbf{k}$ , where  $\mathbf{n}_1$  and  $\mathbf{n}_2$  are directions of the lines  $\Rightarrow$  the plane

containing the lines is represented by  $(-20)(x - 1) + (12)(y - 2) + (1)(z - 3) = 0 \Rightarrow -20x + 12y + z = 7$ .

28.  $\begin{cases} x = t = 2s + 2 \\ y = -t + 2 = s + 3 \end{cases} \Rightarrow \begin{cases} t - 2s = 2 \\ -t - s = 1 \end{cases} \Rightarrow s = -1 \text{ and } t = 0; \text{ then } z = t + 1 = 5s + 6 \Rightarrow 0 + 1 = 5(-1) + 6$   
 is satisfied  $\Rightarrow$  the lines do intersect when  $s = -1$  and  $t = 0 \Rightarrow$  the point of intersection is  $x = 0, y = 2$  and  $z = 1$

or  $P(0, 2, 1)$ . A vector normal to the plane determined by these lines is  $\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -1 & 1 \\ 2 & 1 & 5 \end{vmatrix} = -6\mathbf{i} - 3\mathbf{j} + 3\mathbf{k}$ ,

where  $\mathbf{n}_1$  and  $\mathbf{n}_2$  are directions of the lines  $\Rightarrow$  the plane containing the lines is represented by  
 $(-6)(x-0) + (-3)(y-2) + (3)(z-1) = 0 \Rightarrow 6x + 3y - 3z = 3.$

29. The cross product of  $\mathbf{i} + \mathbf{j} - \mathbf{k}$  and  $-4\mathbf{i} + 2\mathbf{j} - 2\mathbf{k}$  has the same direction as the normal to the plane

$$\Rightarrow \mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & -1 \\ -4 & 2 & -2 \end{vmatrix} = 6\mathbf{j} + 6\mathbf{k}. \text{ Select a point on either line, such as } P(-1, 2, 1). \text{ Since the lines are given}$$

to intersect, the desired plane is  $0(x+1) + 6(y-2) + 6(z-1) = 0 \Rightarrow 6y + 6z = 18 \Rightarrow y + z = 3.$

30. The cross product of  $\mathbf{i} - 3\mathbf{j} - \mathbf{k}$  and  $\mathbf{i} + \mathbf{j} + \mathbf{k}$  has the same direction as the normal to the plane

$$\mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & -3 & -1 \\ 1 & 1 & 1 \end{vmatrix} = -2\mathbf{i} - 2\mathbf{j} + 4\mathbf{k}. \text{ Select a point on either line, such as } P(0, 3, -2). \text{ Since the lines are}$$

given to intersect, the desired plane is  $(-2)(x-0) + (-2)(y-3) + (4)(z+2) = 0 \Rightarrow -2x - 2y + 4z = -14$   
 $\Rightarrow x + y - 2z = 7.$

31.  $\mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 1 & -1 \\ 1 & 2 & 1 \end{vmatrix} = 3\mathbf{i} - 3\mathbf{j} + 3\mathbf{k}$  is a vector in the direction of the line of intersection of the planes

$\Rightarrow 3(x-2) + (-3)(y-1) + 3(z+1) = 0 \Rightarrow 3x - 3y + 3z = 0 \Rightarrow x - y + z = 0$  is the desired plane containing  $P_0(2, 1, -1)$

32. A vector normal to the desired plane is  $\vec{P_1P_2} \times \mathbf{n} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 0 & -2 \\ 4 & -1 & 2 \end{vmatrix} = -2\mathbf{i} - 12\mathbf{j} - 2\mathbf{k}$ ; choosing  $P_1(1, 2, 3)$  as a point on

the plane  $\Rightarrow (-2)(x-1) + (-12)(y-2) + (-2)(z-3) = 0 \Rightarrow -2x - 12y - 2z = -32 \Rightarrow x + 6y + z = 16$  is the desired plane

33.  $S(0, 0, 12)$ ,  $P(0, 0, 0)$  and  $\mathbf{v} = 4\mathbf{i} - 2\mathbf{j} + 2\mathbf{k} \Rightarrow \vec{PS} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 0 & 0 & 12 \\ 4 & -2 & 2 \end{vmatrix} = 24\mathbf{i} + 48\mathbf{j} = 24(\mathbf{i} + 2\mathbf{j})$

$\Rightarrow d = \frac{|\vec{PS} \times \mathbf{v}|}{|\mathbf{v}|} = \frac{24\sqrt{1+4}}{\sqrt{16+4+4}} = \frac{24\sqrt{5}}{\sqrt{24}} = \sqrt{5 \cdot 24} = 2\sqrt{30}$  is the distance from  $S$  to the line

34.  $S(0, 0, 0)$ ,  $P(5, 5, -3)$  and  $\mathbf{v} = 3\mathbf{i} + 4\mathbf{j} - 5\mathbf{k} \Rightarrow \vec{PS} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -5 & -5 & 3 \\ 3 & 4 & -5 \end{vmatrix} = 13\mathbf{i} - 16\mathbf{j} - 5\mathbf{k}$

$\Rightarrow d = \frac{|\vec{PS} \times \mathbf{v}|}{|\mathbf{v}|} = \frac{\sqrt{169+256+25}}{\sqrt{9+16+25}} = \frac{\sqrt{450}}{\sqrt{50}} = \sqrt{9} = 3$  is the distance from  $S$  to the line

35.  $S(2, 1, 3)$ ,  $P(2, 1, 3)$  and  $\mathbf{v} = 2\mathbf{i} + 6\mathbf{j} \Rightarrow \vec{PS} \times \mathbf{v} = \mathbf{0} \Rightarrow d = \frac{|\vec{PS} \times \mathbf{v}|}{|\mathbf{v}|} = \frac{0}{\sqrt{40}} = 0$  is the distance from  $S$  to the line  
 (i.e., the point  $S$  lies on the line)

36.  $S(2, 1, -1)$ ,  $P(0, 1, 0)$  and  $\mathbf{v} = 2\mathbf{i} + 2\mathbf{j} + 2\mathbf{k} \Rightarrow \vec{PS} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 0 & -1 \\ 2 & 2 & 2 \end{vmatrix} = 2\mathbf{i} - 6\mathbf{j} + 4\mathbf{k}$

$\Rightarrow d = \frac{|\vec{PS} \times \mathbf{v}|}{|\mathbf{v}|} = \frac{\sqrt{4+36+16}}{\sqrt{4+4+4}} = \frac{\sqrt{56}}{\sqrt{12}} = \sqrt{\frac{14}{3}}$  is the distance from  $S$  to the line