

Algebraic Geometry, Math 539, Homework 1

1. Let X be a smooth projective surface and $C \subset X$ a smooth curve. Using the exact sequence (which you should know),

$$0 \rightarrow \mathcal{O}_C(-C) \rightarrow \Omega_X^1|_C \rightarrow \Omega_C^1 \rightarrow 0,$$

show that $K_C = \Omega_C^1 = K_X \otimes \mathcal{O}_C(C)$, where as usual $K_X = \wedge^2 \Omega_X^1$, the canonical line bundle of X .

2. Let C be a smooth projective curve and let $\Delta \subset C \times C$ be the diagonal. Using the above, compute Δ^2 , the intersection number in terms of the genus of C .
3. Let X be a smooth projective surface and let $C, D \subset X$ be two curves. For a point $P \in X$, assume that P is isolated in $C \cap D$. Define $I(P, C, D) = \ell(\mathcal{O}_{X,P}/\mathcal{O}(-C) + \mathcal{O}(-D))$, the local intersection multiplicity where as usual ℓ denotes the length. If $C \cap D$ is a finite set of points, show that $(C \cdot D) = \sum_{P \in C \cap D} I(P, C, D)$.
4. Let notation be as above. Show that $I(P, C, D) = 1$ if and only if $P \in C \cap D$ and both C, D are smooth at P and the local equations of C, D at P form a local co-ordinate system for X .
5. Let $C, D \subset \mathbb{P}^2$ be two curves with no common components with $\deg C = d, \deg D = e$. Show that $(D \cdot C) = de$. This is known as Bezout's theorem, which started all of intersection theory.
6. Let $C, D \subset \mathbb{P}^2$ be smooth curves of degrees d, e respectively. Assume that $C \cap D$ has cardinality de and assume that they are defined by polynomials F, G . If $H = 0$ defines a curve E passing through all the points of $C \cap D$, show that $H = AF + BG$ for suitable polynomials A, B . (This is a slightly weak version of what is known as Max Noether's Theorem).
7. Using the above, deduce the following: If $C, D \subset \mathbb{P}^2$ are two smooth cubics meeting in nine distinct points and if E is a third cubic passing through eight of these nine points, then it also passes through the ninth.