

Lecture 9 : More on Determinants

In lecture 8, we defined $\det: \{ \text{"n} \times \text{n real matrices} \} \rightarrow \mathbb{R}$
 to be the unique function satisfying

- (i) linearity in each row (with the other rows held fixed)
- (ii) antisymmetry in the rows
- (iii) $\det I_n = 1$.

From these properties, we quickly deduced that

- if 2 rows are equal, then $\det A = 0$
- if A is upper/lower triangular, $\det A = \text{product of diagonal entries}$
 (in particular, if A is diagonal, i.e. $A = \begin{pmatrix} d_1 & & 0 \\ 0 & \ddots & 0 \\ & & d_n \end{pmatrix}$ then $\det A = d_1 \cdots d_n$)
- if $A_{ij}^{\hat{}} = (n-1) \times (n-1)$ matrix obtained by deleting i^{th} row + j^{th} column of A ,
 $C_{ij} = (-1)^{i+j} \det A_{ij}^{\hat{}} = (i,j)^{\text{th}}$ cofactor, then (for any i)
 $\det A = \sum_{j=1}^n a_{ij} C_{ij}$ (\in Laplace expansion along the i^{th} row).

Consequently,

- if A has a row of 0's, then $\det A = 0$.

Ex 1 /

$$\begin{vmatrix} 1 & 0 & 0 & 1 \\ 3 & 1 & 0 & 0 \\ 0 & 2 & 1 & 0 \\ -1 & 3 & 0 & 0 \end{vmatrix} = 1 \cdot \underbrace{\begin{vmatrix} 3 & 1 & 0 \\ 0 & 2 & 1 \\ -1 & 3 & 0 \end{vmatrix}}_{A_{11}^{\hat{}}} - 1 \cdot \underbrace{\begin{vmatrix} 1 & 3 & 1 \\ 0 & 2 & 1 \\ 1 & -1 & 3 \end{vmatrix}}_{A_{12}^{\hat{}}}$$

$$= 3 \begin{vmatrix} 2 & 1 \\ 3 & 0 \end{vmatrix} - 1 \cdot \begin{vmatrix} 0 & 1 \\ -1 & 0 \end{vmatrix} + 1 \cdot \begin{vmatrix} 3 & 1 \\ -1 & 3 \end{vmatrix} + 2 \cdot \begin{vmatrix} 1 & 3 \\ 1 & -1 \end{vmatrix}$$

$$= 3(-3) - 1 \cdot 1 + 1(9+1) + 2(-1-3) = -8.$$



Determinants and EROs (Elementary Row Operations)

Let E be an elementary matrix, so that $\tilde{A} = E \cdot A$ is one row operation applied to A .

Theorem 1: If E is a $\begin{cases} \text{replace} \\ \text{swap} \\ \text{scale by } \mu \end{cases}$ operation, $\det \tilde{A} = \begin{cases} \det A \\ -\det A \\ \mu \det A \end{cases}$.

Proof: Write $A = \begin{pmatrix} \vec{r}_1 \\ \vdots \\ \vec{r}_n \end{pmatrix}$.

$$\det \begin{pmatrix} \vec{r}_1 \\ \vdots \\ \vec{r}_j \\ \vec{r}_i + \alpha \vec{r}_j \\ \vdots \\ \vec{r}_n \end{pmatrix} = \det \underbrace{\begin{pmatrix} \vec{r}_1 \\ \vdots \\ \vec{r}_j \\ \vec{r}_i \\ \vdots \\ \vec{r}_n \end{pmatrix}}_A + \alpha \cdot \det \begin{pmatrix} \vec{r}_1 \\ \vdots \\ \vec{r}_j \\ \vec{r}_i \\ \vdots \\ \vec{r}_n \end{pmatrix} \stackrel{\text{eqn}}{=} \det A \quad \text{by linearity}$$

$$\det \begin{pmatrix} \vec{r}_1 \\ \vdots \\ \vec{r}_i \\ \vec{r}_j \\ \vdots \\ \vec{r}_n \end{pmatrix} = - \det \begin{pmatrix} \vec{r}_1 \\ \vdots \\ \vec{r}_j \\ \vec{r}_i \\ \vdots \\ \vec{r}_n \end{pmatrix} = -\det A \quad \text{by antisymmetry}$$

$$\det \begin{pmatrix} \vec{r}_1 \\ \vdots \\ \mu \vec{r}_i \\ \vdots \\ \vec{r}_n \end{pmatrix} = \mu \det \begin{pmatrix} \vec{r}_1 \\ \vdots \\ \vec{r}_i \\ \vdots \\ \vec{r}_n \end{pmatrix} = \mu \det A. \quad \text{by linearity}$$

□

Ex 2/ Find $\det \begin{pmatrix} 1 & -1 & 2 & -2 \\ -1 & 2 & 1 & 6 \\ 2 & 1 & 14 & 10 \\ -2 & 6 & 10 & 33 \end{pmatrix}$.

Idea: row-reduce to an upper triangular matrix by only replace operations (& swaps, if needed — not needed here).

$$\begin{vmatrix} 1 & -1 & 2 & -2 \\ 0 & 1 & 3 & 4 \\ 0 & 3 & 10 & 14 \\ 0 & 9 & 14 & 29 \end{vmatrix} \underset{\text{==}}{=} \begin{vmatrix} 1 & -1 & 2 & -2 \\ 0 & 1 & 3 & 4 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 2 & 13 \end{vmatrix} = \begin{vmatrix} 1 & -1 & 2 & -2 \\ 0 & 1 & 3 & 4 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 9 \end{vmatrix} = 1^3 \cdot 9 = 9.$$



Determinants and Invertibility

Theorem 2: $\det A \neq 0 \Leftrightarrow A$ invertible.

Proof: (\Leftarrow): A invertible $\Rightarrow A$ is obtained from \mathbb{I}_n by EROs
 $\Rightarrow \det A = \det \mathbb{I}_n \cdot (-1)^{\text{# swaps}}$
 Thm. 1 ~~(product of scaling factors)~~
 $\neq 0$.

(\Rightarrow): $\det A \neq 0 \Rightarrow \det(\text{rref } A) = \det A \cdot (-1)^{\frac{\#\text{rows}}{\text{# rows}}} \cdot (\text{product of scaling factors}) \neq 0$

$\Rightarrow \text{rref } A$ has no row of 0's

$\xrightarrow[\text{A square}]{} \text{rref } A = \mathbb{I}_n$

$\rightarrow A$ invertible.



Ex 3/ we know $\det \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} \neq \mathbb{I}_3$, so

$$\det \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix} = 0.$$

//

Ex 4/ Find all values of α they make

$$A = \begin{pmatrix} \alpha & 1 & 1 \\ \alpha & \alpha & 1 \\ 4 & \alpha & \alpha \end{pmatrix} \quad \text{non-invertible ("singular")}$$

Want $0 \Rightarrow$

$$\begin{vmatrix} \alpha & 1 & 1 \\ \alpha & \alpha & 1 \\ 4 & \alpha & \alpha \end{vmatrix} = \begin{vmatrix} \alpha & 1 & 1 \\ 0 & \alpha-1 & 0 \\ 4 & \alpha & \alpha \end{vmatrix} = (\alpha-1) \begin{vmatrix} \alpha & 1 \\ 4 & \alpha \end{vmatrix} = (\alpha-1)(\alpha^2-4) = (\alpha-1)(\alpha-2)(\alpha+2)$$

↑ expand in middle row

\Rightarrow values are $\alpha = 1, 2, -2.$

//

Determinants and Products

Consider the elementary matrices once more. What are their determinants?

$$\begin{vmatrix} 1 & & & \\ & \ddots & & \\ & & \alpha & \\ & & & \ddots & 1 \end{vmatrix} = 1, \quad \begin{vmatrix} 1 & & & \\ & \ddots & & \\ & & 0 & 1 & & 1 \\ & & & 0 & \ddots & \\ & & & & 0 & \dots & 1 \end{vmatrix} = -1, \quad \begin{vmatrix} 1 & & & \\ & \ddots & & \\ & & \mu & \\ & & & \ddots & 1 \\ & & & & \ddots & 1 \end{vmatrix} = \mu$$

Replace

(upper or lower triangular with 1's on diagonal)

Drop

(get In by swapping its i^{th} rows)

Scale

Looking at the statement of Theorem 1, we see it on

be rephrased as follows:

(*) If E is an elementary matrix, then
 $\det(EA) = \det(E) \cdot \det(A)$.
↑
[result of applying ERO
to A]

Theorem 3: Given $n \times n$ matrices $A \neq B$, $\det AB = (\det A)(\det B)$.

Proof: Case 1: $\det A = 0$. Then A isn't invertible.

If AB had an inverse C , then $\mathbb{I}_n = (AB)C = A(BC)$
 $\Rightarrow BC$ is inverse to A , a contradiction. So
 AB isn't invertible, and $\therefore \det AB = 0$.

Case 2: $\det A \neq 0$. Then A is invertible, and so
may be written as a product of elementary matrices:

$$A = E_N \cdots E_1 (\mathbb{I}_n)$$

 $\rightarrow AB = E_N \cdots E_1 B.$

By repeated application of (*),

$$\det A = \det E_N \cdots \det E_1 \det \mathbb{I}_n^{\rightarrow}$$

$$\& \det AB = \det E_N \cdots \det E_1 \det B, \text{ so } \det AB = \det A \cdot \det B. \quad \square$$

Corollary: If A is invertible, $\det(A^{-1}) = \frac{1}{\det A}$.

Ex 5 / $\det B^{-1}A^9B = (\det B)^{-1}(\det A)^9 \det B = (\det A)^9$. //

(What about $A+B$? In fact, we can't say anything about its determinant. It's certainly false that $\det A+B = \det A + \det B$; for example, take $A = \mathbb{I}_2$, $B = -\mathbb{I}_2$. Then $\det A + \det B = 1+1=2$, but $A+B$ is the zero matrix.)

Determinants and Column Vectors

I mentioned in Lecture 8 that if you define "det" to be the unique alternating multilinear normalized function on $\{\text{columns of } A\}$ then you get the same function as doing it via rows (as we've done). Since transposing takes rows to columns, this statement is equivalent to

Theorem 4: $\det A = \det A^T$.

Proof: First, $\det A \neq 0 \Leftrightarrow \det A^T \neq 0$, since they are both invertible (with $(A^T)^{-1} = (A^{-1})^T$) or both not.

So if they're invertible, we have $A = E_1 \cdots E_N$ and $A^T = E_N^T \cdots E_1^T$. By Thm. 3, it suffices to check $\det E_i = \det E_i^T$. But this is obvious: Swap & scale matrices are unchanged by transpose, and replace matrices have determinant 1.



Corollary 5:
(i) Elementary column operations have the same effect
on $\det A$ as ERO's.

(ii) Laplace expansion holds for columns: for each j ,
 $\det A = \sum_{i=1}^n a_{ij} C_{ij}$.

Determinants and Linear Independence

Recall from lecture 6 that the following conditions on an $n \times n$ (square) matrix A are equivalent:

- (C1) A is invertible for the corresponding L.T.
 T is invertible
- (C2) The columns of A span \mathbb{R}^n T is onto
- (C3) The columns of A are linearly independent T is 1-1

The same goes for rows since columns of A are rows of A^T .

Theorem 5: $\det A \neq 0 \iff$ rows are lin. ind. & span \mathbb{R}^n
 \iff columns are lin. ind. & span \mathbb{R}^n .

Ex 8 / For what values of α is $\begin{pmatrix} \alpha \\ \alpha \\ 4 \end{pmatrix}, \begin{pmatrix} 1 \\ \alpha \\ \alpha \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ \alpha \end{pmatrix}$
a linearly independent set?

We already solved this: They're independent \Leftrightarrow

$$\det \begin{pmatrix} \alpha & 1 & 1 \\ \alpha & -1 & 1 \\ 4 & 2 & 2 \end{pmatrix} \neq 0 \Leftrightarrow \alpha \neq 1, 2, -2.$$

