

# Ma 450: Mathematics for Multimedia

## Homework Assignment 2

Prof. Wickerhauser

Due Friday, February 17th, 2012

- (a) How many vertices are there in the unit cube in Euclidean  $N$ -space?  
(b) Fix a vertex in the  $N$ -cube. How many other vertices are connected to it by single edges?  
(c) How many edges are there?
- Let  $\mathbf{P}, \mathbf{Q}, \mathbf{S}$  be subspaces of  $\mathbf{R}^N$  with respective dimensions  $p, q, s$ . Suppose that  $\mathbf{S} = \mathbf{P} + \mathbf{Q}$ . (a) Prove that  $\max\{p, q\} \leq s \leq p + q$ . (b) Find an example that achieves the equality  $s = \max\{p, q\}$ . (c) Find an example that achieves the equality  $s = p + q$ .
- Prove Inequality 2.15 for every  $N$ .
- Prove that  $\|\mathbf{x} - \mathbf{y}\| \geq \left| \|\mathbf{x}\| - \|\mathbf{y}\| \right|$  for any vectors  $\mathbf{x}, \mathbf{y}$  in a normed vector space  $\mathbf{X}$ .
- Suppose that  $\mathbf{Y}$  is an  $m$ -dimensional subspace of an  $N$ -dimensional inner product space  $\mathbf{X}$ . Prove that  $\mathbf{Y}^\perp$  is at most  $N - m$  dimensional.
- Suppose that  $\mathbf{Y} = \text{span}\{\mathbf{y}_n : n = 1, \dots, N\}$  and  $\mathbf{Z} = \text{span}\{\mathbf{z}_m : m = 1, \dots, M\}$  are subspaces in an inner product space  $\mathbf{X}$ . Show that if  $\langle \mathbf{y}_n, \mathbf{z}_m \rangle = 0$  for all  $n, m$ , then  $\mathbf{Y} \perp \mathbf{Z}$ .
- Find an orthonormal basis for the subspace of  $\mathbf{E}^4$  spanned by the vectors  $\mathbf{x} = (1, 0, 0, 0)$ ,  $\mathbf{y} = (1, 0, 1, 0)$ , and  $\mathbf{z} = (1, 1, 1, 0)$ .

- Find the biorthogonal dual of the basis  $\left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} \right\}$  of  $\mathbf{E}^3$ .

- Prove that the inner product on  $\mathbf{Poly}$  given by

$$\langle p, q \rangle \stackrel{\text{def}}{=} \sum_k \bar{a}_k b_k,$$

is Hermitean symmetric, nondegenerate, and linear. Here  $p(x) = a_0 + a_1x + \dots + a_nx^n$ ,  $q(x) = b_0 + b_1x + \dots + b_mx^m$ , and the sum is over all nonzero terms  $\bar{a}_k b_k$ . Note that this inner product defines the derived norm in Equation 2.21.

- Compute  $\|T\|_{\text{op}}$  for  $T : \mathbf{Poly} \rightarrow \mathbf{Poly}$  defined by  $Tp(x) = xp(x)$ , with respect to the norm in Equation 2.21. Do the same for  $S : \mathbf{Poly} \rightarrow \mathbf{Poly}$  defined by  $Sp(x) = (1 + x)p(x)$ .
- Suppose that  $A$  is an *idempotent*  $N \times N$  matrix, namely  $A^k = Id$  for some integer  $k > 0$ . Prove that  $\|A\|_{\text{HS}} \geq 1$ .
- Can there be matrices  $A, B \in \mathbf{Mat}(N \times N)$  satisfying  $AB - BA = Id$ ?

13. For each  $i \in \{1, 2, \dots, N\}$ , prove that the transformation  $P_i : \mathbf{R}^N \rightarrow \mathbf{R}^N$  defined by

$$P_i(x_1, \dots, x_N) \stackrel{\text{def}}{=} x_i \mathbf{e}_i$$

is an orthogonal projection onto  $\text{span}\{\mathbf{e}_i\} \subset \mathbf{R}^N$ .

14. Let  $e_k(x) \stackrel{\text{def}}{=} \exp(2\pi i k x)$  for each integer  $k$  and all  $x \in [0, 1]$ . For integers  $-\infty < M \leq N < +\infty$  and any function  $f \in \mathbf{Lip}$ , prove that

$$\sum_{k=M}^N |\langle e_k, f \rangle|^2 \leq \|f\|^2.$$

for the norm and inner product for  $\mathbf{Lip}$  defined in Equation 2.28.

15. Determine whether the linear transformation  $T : \ell^2 \rightarrow \ell^2$  defined by

$$T(x_1, x_2, x_3, \dots) = (x_1, \frac{x_2 + x_3}{2}, \frac{x_4 + x_5 + x_6}{3}, \dots)$$

is bounded or unbounded.

16. Given a matrix  $A = \begin{pmatrix} x & y \\ z & w \end{pmatrix}$ , find a Givens rotation  $G$  such that  $GA$  is upper triangular.
17. Suppose that  $A$  and  $B$  are  $N \times N$  matrices satisfying the condition  $A(i, j) = B(i, j) = 0$  if  $i > j$ . Prove that their product satisfies the same condition. (This shows that the product of upper-triangular matrices is upper-triangular.)
18. Write a computer program that takes a list of edges in three-dimensional space, presented as pairs of coordinate triplets, and returns their plane coordinates as they would appear projected onto the  $xy$ -plane with perspective. Allow the user to specify the viewpoint  $(x_0, y_0, z_0)$ .