Online First version

Asymptotic Müntz–Szász theorems

by

JIM AGLER (La Jolla, CA) and JOHN E. MCCARTHY (St. Louis, MO)

Abstract. We define a monomial space to be a subspace of $L^{2}([0, 1])$ that can be approximated by spaces that are spanned by monomial functions. We describe the structure of monomial spaces.

1. Introduction. What sorts of subspaces in $L^2([0,1])$ can be limits of spans of monomials? Specifically, let

$$\mathbb{S} = \{ s \in \mathbb{C} \mid \operatorname{Re} s > -1/2 \},\$$

so that $s \in S$ if and only if $x^s \in L^2([0,1])$. For S a finite subset of S we let $\mathcal{M}(S)$ denote the span in $L^2([0,1])$ of the monomials whose exponents lie in S, i.e.,

$$\mathcal{M}(S) = \Big\{ \sum_{s \in S} a(s) x^s \ \Big| \ a : S \to \mathbb{C} \Big\}.$$

We refer to sets in $L^2([0,1])$ that have the form $\mathcal{M}(S)$ for some finite subset S of S as *finite monomial spaces*. We are interested in what the limits of such spaces are.

DEFINITION 1.1. If \mathcal{M} is a subspace of a Hilbert space \mathcal{H} and $\{\mathcal{M}_n\}$ is a sequence of closed subspaces, we say that $\{\mathcal{M}_n\}$ tends to \mathcal{M} and write $\mathcal{M}_n \to \mathcal{M}$ as $n \to \infty$ if

$$\mathcal{M} = \Big\{ f \in \mathcal{H} \ \Big| \ \lim_{n \to \infty} \operatorname{dist}(f, \mathcal{M}_n) = 0 \Big\}.$$

There are alternative ways to frame this definition; see Proposition 4.1.

DEFINITION 1.2. We say that a subspace \mathcal{M} of $L^2([0,1])$ is a monomial space if there exists a sequence $\{\mathcal{M}_n\}$ of finite monomial spaces such that $\mathcal{M}_n \to \mathcal{M}$.

²⁰²⁰ Mathematics Subject Classification: Primary 47A15; Secondary 41A10.

Key words and phrases: Hardy operator, Müntz spaces, monomial spaces, Beurling's theorem.

Received 13 July 2022; revised 2 February 2023. Published online *.

The goal of this paper is to study monomial spaces, which have a rich structure and are intimately related to the Müntz-Szasz theorem and generalizations thereof.

1.1. Monotone monomial spaces. In this subsection we recall several classical results that can be interpreted as facts about monomial spaces. In each example we consider there is a limit $\mathcal{M}_n \to \mathcal{M}$ of finite monomial spaces that is monotone, i.e.,

 $\mathcal{M}_i \subseteq \mathcal{M}_j$ whenever $i \leq j$.

For a detailed account of the results in this section, see [4] and [7].

EXAMPLE 1.3 (The Weierstrass approximation theorem). Let $S_n = \{0, 1, \dots, n\}$. The Weierstrass theorem, which implies that the polynomials are dense in $L^2([0, 1])$, also implies that

$$\mathcal{M}(S_n) \to \mathrm{L}^2([0,1]).$$

In particular, $L^2([0, 1])$ is a monomial space.

EXAMPLE 1.4 (Classical Müntz–Szász theorem [24, 28]). Fix a strictly increasing sequence of nonnegative integers s_0, s_1, s_2, \ldots , and let

$$S_n = \{s_0, s_1, \dots, s_n\}.$$

Then there exists a space \mathcal{M} such that

$$\mathcal{M}(S_n) \to \mathcal{M}.$$

Furthermore,

$$\mathcal{M} = \mathrm{L}^2([0,1])$$
 if and only if $\sum_{k=1}^{\infty} \frac{1}{s_k} = \infty.$

EXAMPLE 1.5 (Szász's theorem, real case (also known as full Müntz– Szász theorem in L²([0,1])) [29]). Fix a sequence of distinct real numbers s_0, s_1, s_2, \ldots in S and let again $S_n = \{s_0, s_1, \ldots, s_n\}$. Then there exists a space \mathcal{M} such that $\mathcal{M}(S_n) \to \mathcal{M}$. Furthermore,

$$\mathcal{M} = L^2([0,1])$$
 if and only if $\sum_{k=0}^{\infty} \frac{2s_k + 1}{(2s_k + 1)^2 + 1} = \infty.$

EXAMPLE 1.6 (Szász's theorem, complex case). Any of the proofs that the authors know of the previous example, including Szász's original proof, can be adapted to show that if s_0, s_1, s_2, \ldots is a sequence of distinct points in S and, as previously, $S_n = \{s_0, s_1, \ldots, s_n\}$, then there exists a space \mathcal{M} such that $\mathcal{M}(S_n) \to \mathcal{M}$, and where

$$\mathcal{M} = L^2([0,1])$$
 if and only if $\sum_{k=1}^{\infty} \frac{2 \operatorname{Re} s_k + 1}{|s_k + 1|^2} = \infty.$

What happens in the above examples when $\mathcal{M} \neq L^2([0,1])$?

EXAMPLE 1.7 (The Clarkson-Erdős theorem). With the setup of Example 1.4, assume that $\mathcal{M} \neq L^2([0, 1])$. Then Clarkson and Erdős proved in [12] that the elements of \mathcal{M} extend to be analytic on \mathbb{D} ! Furthermore, if $f \in \mathcal{M}$, then f has a power series representation of the form

$$f(z) = \sum_{k=0}^{\infty} a_k z^{s_k}, \quad z \in \mathbb{D}.$$

This result was generalized to arbitrary real powers in $(-1/2, \infty)$ by Erdélyi and Johnson [16], who showed that if $\mathcal{M} \neq L^2([0, 1])$, then every f in \mathcal{M} is analytic in $\mathbb{D} \setminus (-1, 0]$.

In honor of this remarkable theorem, we introduce the following definition.

DEFINITION 1.8. We say that \mathcal{M} is a *Clarkson–Erdős space* if there exist a sequence $\{s_0, s_1, \ldots\}$ in \mathbb{S} of distinct points such that

$$\mathcal{M}(\{s_0, s_1, \dots, s_n\}) \to \mathcal{M}$$

where $\mathcal{M} \neq L^2([0,1])$.

We want to allow for multiplicities. If an entry s is repeated in a sequence, this corresponds to multiplicity in the following way. The first occurrence of s in S_n gives the function x^s in \mathcal{M}_n . The second occurrence gives $\frac{\partial}{\partial s}x^s = x^s \log x$. If s occurs k times, then \mathcal{M}_n contains the functions $x^s, x^s \log x, \ldots, x^s (\log x)^{k-1}$. This leads to the following generalization of a Clarkson-Erdős space.

DEFINITION 1.9. We say that \mathcal{M} is an *Erdélyi–Johnson space* if there exist a sequence $\{s_0, s_1, \ldots\}$ in \mathbb{S} , with multiplicities allowed, such that

 $\mathcal{M}(\{s_0, s_1, \ldots, s_n\}) \to \mathcal{M}$

where $\mathcal{M} \neq L^2([0,1])$.

1.2. A nonmonotone monomial space. In the study of monomial spaces it is natural to consider the class of monomial operators, i.e., the class of bounded operators T acting on $L^2([0,1])$ that take monomials to monomials, i.e.,

(1.1)
$$\forall_{s\in\mathbb{S}} \exists_{\tau\in\mathbb{S}} \exists_{c\in\mathbb{C}} Tx^s = cx^{\tau}.$$

In [2] the authors studied the special case where it is assumed that there exists a fixed number m such that in (1.1), τ can be chosen to equal s + m for all s. We call these *flat monomial operators*. In the course of proving that flat monomial operators leave $L^2([a, 1])$ invariant for each $a \in [0, 1]$, the authors discovered the following example.

EXAMPLE 1.10. Fix $\rho \in [0, 1]$ and choose an increasing sequence of integers N_1, N_2, \ldots such that

$$\lim_{n \to \infty} \frac{n}{n+N_n} = \sqrt{\rho}.$$

If

$$S_n = \{n+1, n+2, \dots, n+N_n\},\$$

then

$$\mathcal{M}(S_n) \to \mathrm{L}^2([\rho, 1]).$$

In particular, $L^2([\rho, 1])$ is a monomial space for each $\rho \in [0, 1]$, where here and afterwards we identify $L^2([\rho, 1])$ with the subspace of $L^2([0, 1])$ consisting of functions that vanish a.e. on $[0, \rho]$.

1.3. Characterization of monomial spaces. The Hardy operator $H : L^2([0,1]) \to L^2([0,1])$ is defined by

(1.2)
$$Hf(x) = \frac{1}{x} \int_{0}^{x} f(t) dt.$$

This was introduced by Hardy [21], who proved it was bounded. As $Hx^s = \frac{1}{s+1}x^s$ for all $s \in S$, the Hardy operator leaves invariant every monomial space. The converse is true.

THEOREM 1.11. A closed subspace of $L^2([0,1])$ is a monomial space if and only if it is invariant for H.

A proof of 1.11 that uses real analysis techniques is given in [1].

COROLLARY 1.12. A bounded operator T on $L^2([0,1])$ is a monomial operator if and only if for every $\mathcal{M} \in Lat(H)$, the space $T\mathcal{M}$ is in Lat(H).

1.4. A decomposition theorem

DEFINITION 1.13. We say a space \mathcal{M} in $L^2([0,1])$ is a *singular space* if \mathcal{M} is a monomial space that does not contain any Clarkson–Erdős space.

THEOREM 1.14. Every monomial space \mathcal{M} has a unique decomposition,

$$\mathcal{M}=\overline{\mathcal{M}_0+\mathcal{M}_1},$$

where \mathcal{M}_0 is an Erdélyi–Johnson space and \mathcal{M}_1 is a singular space.

1.5. Atomic spaces. Unitary monomial operators can be characterized using a theorem of Bourdon and Narayan [8]. Their description is equivalent to the following reformulation from [3].

THEOREM 1.15. The operator

$$T: x^s \mapsto c(s) x^{\tau(s)}$$

4

is a unitary map from $L^2([0,1])$ to $L^2([0,1])$ if and only if τ is a holomorphic automorphism of S and c is given by

$$c(s) = c_0 \frac{1 + \overline{\tau(0)} + \tau(s)}{1 + s},$$

where c_0 is a constant satisfying

$$|c_0| = \frac{1}{\sqrt{1 + 2\operatorname{Re}\tau(0)}}$$

DEFINITION 1.16. We say a space \mathcal{M} in $L^2([0,1])$ is *atomic* if there exist $\rho \in (0,1)$ and a unitary monomial operator T such that $\mathcal{M} = TL^2([\rho,1])$.

We can describe all atomic spaces in the following way.

THEOREM 1.17. The functions

(1.3)
$$e_n(x) = \sum_{k=0}^n \binom{n}{k} \frac{(\ln x)^k}{k!}, \quad n \ge 0,$$

form an orthonormal basis for $L^2([0,1])$. Furthermore, the operator J defined on $L^2([0,1])$ by requiring

$$J(e_n) = \begin{cases} e_n & \text{if } n \text{ is even,} \\ -e_n & \text{if } n \text{ is odd,} \end{cases}$$

is a unitary monomial operator, corresponding to the choice

in Theorem 1.15.

The polynomials

$$p_n(t) = \sum_{k=0}^n \binom{n}{k} \frac{(t)^k}{k!}, \quad n \ge 0,$$

are the Laguerre polynomials. They are the orthogonal polynomials on $[0, \infty)$ for the measure $e^{-t}dt$. Under the change of variables $x = e^{-t}$ they become the functions e_n in (1.3). Their connection to the Hardy operator was shown in [10, 23].

If c is real, the multiplication operator $M_{x^{ic}}$ is also a unitary monomial operator. These two operators can be used to build the general atomic space:

(1) For any w > 0, define

$$\mathcal{A}_{1,w} := L^2([e^{-2w}, 1])$$

(2) Define

$$\mathcal{A}_{-1,w} := J\mathcal{A}_{1,u}$$

(3) For any $\tau \in \mathbb{T} \setminus \{1\}$, define

$$\mathcal{A}_{\tau,w} := M_{x^{ic}} \mathcal{A}_{-1,\wp}, \quad \text{where } 2ic = \frac{\tau+1}{\tau-1}, \ \wp = (1+4c^2)w.$$

(The reason for the strange scaling is to simplify the formulas in Section 5.) Atomic spaces are all of the form $\mathcal{A}_{\tau,w}$.

THEOREM 1.18. Every atomic space is equal to $\mathcal{A}_{\tau,w}$ for exactly one pair $(\tau, w) \in \mathbb{T} \times (0, \infty)$.

1.6. The structure of singular spaces. We say \mathcal{M} is finitely atomic if \mathcal{M} is a finite sum of atomic spaces. If

$$\mu = \sum_{k=1}^{n} w_k \delta_{\tau_k}$$

is a finitely atomic measure on \mathbb{T} , with distinct atoms τ_k , we define a finitely atomic space in $L^2([0,1])$ by the formula

$$\mathcal{M}(\mu) = \sum_{k=1}^{n} \mathcal{A}_{\tau_k, w_k}$$

THEOREM 1.19. The assignment $\mu \mapsto \mathcal{M}(\mu)$ extends by weak-* sequential continuity to a map from the positive singular Borel measures on \mathbb{T} into closed subspaces of $L^2([0,1])$. When extended,

 $\mu_n \to \mu \ weak^* \implies \mathcal{M}(\mu_n) \to \mathcal{M}(\mu).$

1.7. The main idea. It was proved by Brown, Halmos and Shields [10] that the Hardy operator is unitarily equivalent to $1 - S^*$, where S is the unilateral shift, via a unitary $U : L^2([0,1]) \to H^2$ which we call the Sarason transform, described in Section 2. It follows that the invariant subspaces of H^2 can be described by Beurling's theorem [6] in terms of model spaces, the invariant subspaces for the backward shift described for example in [19]. However, all the theorems above have been stated in terms that are *intrinsic* to $L^2([0,1])$. We believe that finding proofs that are also intrinsic to $L^2([0,1])$ will illuminate this space with a new light. So far, the authors have only succeeded in doing this for some of these results.

PROBLEM 1.20. Find real analysis proofs to Theorems 1.14 and 1.19.

2. The Sarason transform

2.1. The definition. We let H^2 denote the classical Hardy space of holomorphic functions on \mathbb{D} with square summable power series coefficients at the origin. We let k_{α} denote the Szegö kernel function for H^2 , i.e.,

$$k_{\alpha}(z) = \frac{1}{1 - \bar{\alpha}z}, \quad z \in \mathbb{D}.$$

As the monomials are linearly independent in $L^2([0,1])$, there is a welldefined map L, defined on polynomials in $L^2([0,1])$ into H^2 by the formula

$$L\left(\sum_{n=0}^{N} a_n x^n\right) = \sum_{n=0}^{N} a_n \frac{1}{n+1} k_{\frac{n}{n+1}}$$

Since

$$\langle L(x^i), L(x^j) \rangle_{\mathrm{H}^2} = \langle x^i, x^j \rangle_{\mathrm{L}^2([0,1])}$$

for all nonnegative integers i and j, it follows that

$$\langle L(p), L(q) \rangle_{\mathrm{H}^2} = \langle p, q \rangle_{\mathrm{L}^2([0,1])}$$

for all polynomials p and q, i.e., L is isometric. Hence, as the polynomials are dense in $L^2([0,1])$, L has a unique extension to an isometry U defined on all of $L^2([0,1])$. Finally, since

$$\left\{\frac{n}{n+1} \mid n \text{ is a nonnegative integer}\right\}$$

is a set of uniqueness for H^2 , it follows that the range of L is dense in H^2 , which implies that U is a unitary transformation from $\mathrm{L}^2([0,1])$ onto H^2 .

DEFINITION 2.1. We let U denote the unique unitary transformation from $L^2([0,1])$ onto H^2 that satisfies

$$U(x^n) = \frac{1}{n+1}k_{\frac{n}{n+1}}$$

for all nonnegative integers n.

We call U the Sarason transform, as it is similar to the transform from $L^2([0,\infty))$ onto H^2 used in [27].

2.2. Moments in $L^2([0,1])$ and interpolation in H^2 . As the monomials are dense in $L^2([0,1])$, a function $f \in L^2([0,1])$ is uniquely determined by its moment sequence

$$\int_{0}^{1} x^{n} f(x) \, dx, \qquad n = 0, 1, \dots.$$

Similarly, as the sequence $\{1 - \frac{1}{n+1} \mid n = 0, 1, ...\}$ is a set of uniqueness for H^2 , a function $h \in \mathrm{H}^2$ is the unique solution g in H^2 to the interpolation problem

$$g\left(\frac{n}{n+1}\right) = h\left(\frac{n}{n+1}\right), \quad n = 0, 1, \dots$$

The following proposition is immediate from Definition 2.1.

PROPOSITION 2.2. Fix a sequence of complex numbers w_0, w_1, w_2, \ldots . If f in $L^2([0,1])$ solves the moment problem

$$\int_{0}^{1} x^{n} f(x) \, dx = w_{n}, \qquad n = 0, 1, \dots,$$

then Uf is in H^2 and solves the interpolation problem

$$Uf\left(\frac{n}{n+1}\right) = (n+1)w_n, \quad n = 0, 1, \dots$$

If $h \in H^2$ solves the interpolation problem

$$h\left(\frac{n}{n+1}\right) = w_n, \quad n = 0, 1, \dots,$$

then U^*h is in $L^2([0,1])$ and solves the moment problem

$$\int_{0}^{1} x^{n} U^{*} h(x) \, dx = \frac{1}{n+1} w_{n}, \qquad n = 0, 1, \dots.$$

The correspondence between moments and interpolation described in the preceding proposition allows us to easily calculate the Sarason transform of many common functions. We illustrate this with the following two lemmas.

LEMMA 2.3. If $\alpha \in \mathbb{D}$, then

(2.1)
$$U^*(k_{\alpha})(x) = \frac{1}{1 - \bar{\alpha}} x^{\frac{\bar{\alpha}}{1 - \bar{\alpha}}}, \quad x \in [0, 1].$$

If $\operatorname{Re}\beta > -\frac{1}{2}$, then

$$U(x^{\beta}) = \frac{1}{\beta+1} k_{\frac{\bar{\beta}}{\bar{\beta}+1}}.$$

Proof. We note that the two assertions of the lemma are equivalent. Therefore it suffices to prove (2.1). Since the left and right hand sides of (2.1) are in $L^2([0, 1])$, it suffices to show that for each $n \ge 0$,

$$\langle x^n, U^* k_\alpha \rangle_{\mathrm{L}^2([0,1])} = \left\langle x^n, \frac{1}{1 - \bar{\alpha}} x^{\frac{\bar{\alpha}}{1 - \bar{\alpha}}} \right\rangle_{\mathrm{L}^2([0,1])}$$

But

$$\begin{split} \left\langle x^n, \frac{1}{1-\bar{\alpha}} x^{\frac{\bar{\alpha}}{1-\bar{\alpha}}} \right\rangle_{\mathcal{L}^2([0,1])} &= \frac{1}{1-\alpha} \left\langle x^n, x^{\frac{\bar{\alpha}}{1-\bar{\alpha}}} \right\rangle_{\mathcal{L}^2([0,1])} = \frac{1}{1-\alpha} \int_0^1 x^n \overline{x^{\frac{\bar{\alpha}}{1-\bar{\alpha}}}} \, dx \\ &= \frac{1}{1-\alpha} \frac{1}{n+\frac{\alpha}{1-\alpha}+1} = \frac{1}{n+1} \frac{1}{1-\frac{n}{n+1}\alpha} \\ &= \frac{1}{n+1} k_{\frac{n}{n+1}}(\alpha) = (Ux^n)(\alpha) = \langle Ux^n, k_\alpha \rangle_{\mathcal{H}^2} \\ &= \langle x^n, U^* k_\alpha \rangle_{\mathcal{L}^2([0,1])}. \quad \bullet \end{split}$$

For S a measurable set in [0, 1] let χ_S denote the characteristic function of S.

LEMMA 2.4. If $s \in [0, 1]$, then

(2.2)
$$U\chi_{[0,s]}(z) = \sqrt{s} \, e^{\frac{1}{2} \ln s} \, \frac{1+z}{1-z}$$

Proof. We first observe that

$$\begin{aligned} \frac{s^{n+1}}{n+1} &= \int_{0}^{s} x^{n} \, dx = \langle \chi_{[0,s]}, x^{n} \rangle_{\mathcal{L}^{2}([0,1])} = \langle U\chi_{[0,s]}, Ux^{n} \rangle_{\mathcal{H}^{2}} \\ &= \left\langle U\chi_{[0,s]}, \frac{1}{n+1} k_{\frac{n}{n+1}} \right\rangle_{\mathcal{H}^{2}} \\ &= \frac{1}{n+1} U\chi_{[0,s]} \left(\frac{n}{n+1}\right), \end{aligned}$$

so that

$$U\chi_{[0,s]}\left(\frac{n}{n+1}\right) = s^{n+1}$$

for all $n \ge 0$. On the other hand, if for w > 0 we let E_w denote the singular inner function defined by

$$E_w(z) = e^{-w\frac{1+z}{1-z}} = e^w e^{-\frac{2w}{1-z}}$$

then

$$E_w\left(\frac{n}{n+1}\right) = e^w e^{-2w(n+1)} = e^w (e^{-2w})^{n+1}$$

for all $n \ge 0$. Hence, if we choose $w = -\frac{1}{2} \ln s$, then

$$U\chi_{[0,s]}\left(\frac{n}{n+1}\right) = e^{-w}E_w\left(\frac{n}{n+1}\right)$$

for all $n \ge 0$. Since $\left\{1 - \frac{1}{n+1} \mid n \ge 0\right\}$ is a set of uniqueness for H^2 , it follows that

$$U\chi_{[0,s]}(z) = e^{-w}E_w(z)$$

for all $z \in \mathbb{D}$, which implies (2.2).

2.3. A formula for the Sarason transform

PROPOSITION 2.5. If $f \in L^2([0,1])$, then

$$Uf(z) = \frac{1}{1-z} \int_{0}^{1} f(x) x^{\frac{z}{1-z}} dx \quad \text{for all } z \in \mathbb{D}.$$

Proof.

$$\begin{split} Uf(z) &= \langle Uf, k_z \rangle_{\mathrm{H}^2} = \langle f, U^* k_z \rangle_{\mathrm{L}^2([0,1])} \\ &= \left\langle f, \frac{1}{1-\bar{z}} x^{\frac{\bar{z}}{1-\bar{z}}} \right\rangle_{\mathrm{L}^2([0,1])} \quad \text{(Lemma 2.3)} \\ &= \frac{1}{1-z} \int_0^1 f(x) x^{\frac{z}{1-z}} \, dx. \quad \bullet \end{split}$$

To obtain a nonrigorous but highly interesting proof of the proposition, let us define the Sarason transform $S\{\mu\}$, of a measure μ on [0, 1], to be the holomorphic function

$$S\{\mu\}(z) = \frac{1}{1-z} \int x^{\frac{z}{1-z}} d\mu(x), \quad |z| < 1.$$

Note that

$$\mathcal{S}(\chi_{[0,s]}) = \sqrt{s} \, e^{\ln \sqrt{s} \, \frac{1+z}{1-z}} = U(\chi_{[0,s]})$$

We have

$$S\{\delta_s\} = \frac{1}{1-z}s^{\frac{1}{1-z}-1} = \frac{1}{1-z}s^{\frac{z}{1-z}}.$$

So formally we get

$$Uf = \mathcal{S}\left(\int_{0}^{1} f(x)\delta_{x} \, dx\right) = \int_{0}^{1} f(x)\mathcal{S}(\delta_{x}) \, dx = \int_{0}^{1} f(x)\frac{1}{1-z}x^{\frac{z}{1-z}} \, dx,$$

the formula in Proposition 2.5.

2.4. Transforms of H, V and X. The Hardy operator H was defined by (1.2). Let X denote multiplication by x on $L^2([0,1])$, and define the Volterra operator $V : L^2([0,1]) \to L^2([0,1])$ by V = XH, so

$$Vf(x) = \int_{0}^{x} f(t) \, dt.$$

All three of these are monomial operators, and have simple descriptions in terms of monomials:

$$H: x^n \mapsto \frac{1}{n+1}x^n, \quad X: x^n \mapsto x^{n+1}, \quad V: x^n \mapsto \frac{1}{n+1}x^{n+1}$$

If T is a bounded operator on $L^2([0,1])$, let us write $\hat{T} = UTU^*$ for the unitarily equivalent operator on H^2 . Monomial operators then become operators that map kernel functions to multiples of other kernel functions, which are adjoints of weighted composition operators. For more about weighted composition operators, see e.g. [5, 8, 11, 13, 14, 17, 22].

We shall let M_g denote the operator of multiplication by g, and C_β denote composition with β . It was observed in [20] that it is possible for the product

10

 $M_g C_\beta$ to be bounded even when M_g is not. The following theorem is proved in [3].

THEOREM 2.6. The operator $T : L^2([0,1]) \to L^2([0,1])$ is a monomial operator if and only if $\widehat{T^*} : H^2 \to H^2$ is a bounded operator of the form $M_q C_\beta$ for some holomorphic $\beta : \mathbb{D} \to \mathbb{D}$ and some $g \in H^2$.

Let $\gamma(z) = \frac{1}{2-z}$. This maps \mathbb{D} to \mathbb{D} , and maps $\frac{n}{n+1}$ to $\frac{n+1}{n+2}$. Using Lemma 2.3 and the preceding formulas, it is easy to verify the following. We shall let S denote the unilateral shift, the operator of multiplication by z on H^2 .

PROPOSITION 2.7. We have

$$\widehat{H} = 1 - S^*, \quad \widehat{X} = S^* C^*_{\gamma}, \quad \widehat{V} = (1 - S^*) C^*_{\gamma}$$

The fact that 1 - H is unitarily equivalent to a backward shift operator was first proved in [10], and a proof similar to ours is in [23]. In [18] Fricain and Lefèvre study other properties that can be ported between $L^2([0, 1])$ and the Hardy space via the Sarason transform (they actually look at the Hardy space of the right half-plane).

2.5. The Sarason transform and Lat(V). The invariant subspaces of the Volterra operator were described by Brodskiĭ [9] and Donoghue [15].

THEOREM 2.8. The space $\mathcal{M} \subseteq L^2([0,1])$ is a closed invariant subspace for V if and only if $\mathcal{M} = L^2([\rho,1])$ for some $\rho \in [0,1]$.

How do these spaces transform under the Sarason transform? For $s \in (0, 1]$ let Φ_s be the singular inner function defined by

$$\Phi_s(z) = e^{\frac{1}{2}\ln s \frac{1+z}{1-z}}, \quad z \in \mathbb{D}.$$

For $s \in [0, 1]$, define orthogonal projections P_s^{\pm} on $L^2([0, 1])$ by the formulas

$$P_s^- f = \chi_{[0,s]} f$$
 and $P_s^+ f = \chi_{[s,1]} f$, $f \in L^2([0,1])$.

LEMMA 2.9.

$$U \operatorname{ran} P_s^- = \Phi_s \operatorname{H}^2$$
 and $U \operatorname{ran} P_s^+ = \Phi_s \operatorname{H}^{2^{\perp}}$

Proof. As ran P_s^- is invariant for H^* , it follows from Proposition 2.7 that $U \operatorname{ran} P_s^-$ is invariant for the shift S, and is therefore of the form $u\mathrm{H}^2$ for some inner function u by Beurling's theorem [6]. Moreover, u is a constant multiple of the projection of 1 onto the invariant subspace. By Lemma 2.4, the projection of 1 is $\sqrt{s} \Phi_s$, so $u = \Phi_s$.

2.6. The Sarason transform and the Laplace transform. Recall that the Laplace transform is defined by the formula

$$\mathcal{L}{f}(s) = \int_{0}^{\infty} e^{-st} f(t) \, dt.$$

Further, if for $f \in L^2([0,1])$ we define f^{\sim} by the formula

$$f^{\sim}(t) = e^{-\frac{t}{2}} f(e^{-t}), \quad t \in (0, \infty),$$

then the assignment $f \mapsto f^{\sim}$ is a Hilbert space isomorphism from $L^2([0, 1])$ onto $L^2((0, \infty))$. By making the substitution $x = e^{-t}$, we find that

$$\int_{0}^{1} f(x) x^{\frac{z}{1-z}} dx = \int_{\infty}^{0} f(e^{-t}) e^{-t \frac{z}{1-z}} (-e^{-t}) dt = \int_{0}^{\infty} e^{-\frac{t}{2}} f(e^{-t}) e^{-t \left(\frac{z}{1-z}+\frac{1}{2}\right)} dt$$
$$= \int_{0}^{\infty} f^{\sim}(t) e^{-t \left(\frac{1}{2}\frac{1+z}{1-z}\right)} dt = \mathcal{L}\{f^{\sim}\} \left(\frac{1}{2}\frac{1+z}{1-z}\right).$$

Hence,

$$Uf(z) = \frac{1}{1-z} \mathcal{L}\lbrace f^{\sim} \rbrace \left(\frac{1}{2} \frac{1+z}{1-z} \right).$$

So after changes of variables from $L^2([0,1])$ of the disc to $L^2((0,\infty))$ and from H^2 to H^2 of the right half-plane, the Sarason transform is simply the Laplace transform.

3. The inverse Sarason transform. We know from Proposition 2.7 that $1 - H^*$ is unitarily equivalent to the unilateral shift. Let us find what the orthonormal basis z^n in H² corresponds to. This was first done in [10], and studied further in [23]. In this section we shall use the notation $\phi \sim g$ to mean that the function $f \in L^2([0, 1])$ is mapped to $\phi \in H^2$ by the Sarason transform.

Lemma 3.1.

(3.1)
$$(H^*)^j 1 = (-1)^j \frac{(\ln x)^j}{j!}.$$

Proof. We proceed by induction. Clearly, (3.1) holds when j = 0. Assume $j \ge 0$ and (3.1) holds. Then

$$(H^*)^{j+1} 1 = H^*((H^*)^j 1) = \frac{(-1)^j}{j!} H^*(\ln x)^j = \frac{(-1)^j}{j!} \int_x^1 \frac{(\ln t)^j}{t} dt$$
$$= \frac{(-1)^j}{j!} \int_{\ln x}^0 u^j \, du = (-1)^{j+1} \frac{(\ln x)^{j+1}}{(j+1)!}.$$

The following result is proved in [10] and [23]; we include a proof for expository reasons.

Lemma 3.2.

$$z^n \sim \sum_{j=0}^n \binom{n}{j} \frac{(\ln x)^j}{j!}.$$

Proof. We have $S = 1 - \hat{H}^*$. Therefore, $z^n = S^n 1 \sim (1 - H^*)^n 1.$

But using Lemma 3.1,

$$(1 - H^*)^n 1 = \sum_{j=0}^n (-1)^j \binom{n}{j} (H^*)^j 1$$
$$= \sum_{j=0}^n (-1)^j \binom{n}{j} \left((-1)^j \frac{(\ln x)^j}{j!} \right) = \sum_{j=0}^n \binom{n}{j} \frac{(\ln x)^j}{j!}.$$

Define

$$e_n(x) = \sum_{j=0}^n \binom{n}{j} \frac{(\ln x)^j}{j!}.$$

We just proved that $e_n = (1 - H^*)^n 1$. The fact that the functions e_n are an orthonormal basis is already well-known. Indeed, the Laguerre polynomials

$$p_n(t) = \sum_{j=0}^n \binom{n}{j} \frac{(-t)^j}{j!}$$

are orthogonal polynomials of norm 1 in $L^2([0,\infty))$ with weight e^{-t} . By the change of variables $x = \ln \frac{1}{t}$, we deduce immediately that $e_n(x)$ is an orthonormal basis for $L^2([0,1])$.

LEMMA 3.3. If $f \in \mathbf{H}^2$ extends to be analytic on a neighborhood of 1, then

$$f(z) \sim \sum_{j=0}^{\infty} f^{(j)}(1) \frac{(\ln x)^j}{(j!)^2}.$$

Proof. If

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

is the power series representation of f, then by Lemma 3.2,

$$f(z) \sim \sum_{n=0}^{\infty} a_n \left(\sum_{j=0}^n \binom{n}{j} \frac{(\ln x)^j}{j!} \right)$$
$$= \sum_{j=0}^{\infty} \frac{(\ln x)^j}{(j!)^2} \left(\sum_{n=j}^{\infty} \frac{n!}{(n-j)!} a_n \right) = \sum_{j=0}^{\infty} f^{(j)}(1) \frac{(\ln x)^j}{(j!)^2}.$$

As a reality check we verify the formula in Lemma 2.3 by using Lemma 3.3. Note that

$$k_{\alpha}^{(j)}(1) = \frac{j!\bar{\alpha}^{j}}{(1-\bar{\alpha})^{j+1}}.$$

Therefore, Lemma 3.3 implies that

$$k_{\alpha}(z) \sim \sum_{j=0}^{\infty} \frac{j!\bar{\alpha}^{j}}{(1-\bar{\alpha})^{j+1}} \frac{(\ln x)^{j}}{(j!)^{2}} = \frac{1}{1-\bar{\alpha}} \sum_{j=0}^{\infty} \frac{1}{j!} \left(\frac{\bar{\alpha}}{1-\bar{\alpha}}\ln x\right)^{j}$$
$$= \frac{1}{1-\bar{\alpha}} e^{\frac{\bar{\alpha}}{1-\bar{\alpha}}\ln x} = \frac{1}{1-\bar{\alpha}} x^{\frac{\bar{\alpha}}{1-\bar{\alpha}}}.$$

The formula in Lemma 3.3 reminds one of Bessel functions. Indeed,

$$e^{z-1} \sim \sum_{j=0}^{\infty} \frac{(\ln x)^j}{(j!)^2} = \sum_{j=0}^{\infty} \frac{(-1)^j}{(j!)^2} (-\ln x)^j$$
$$= \sum_{j=0}^{\infty} \frac{(-1)^j}{(j!)^2} \left(\frac{\sqrt{\ln x^{-4}}}{2}\right)^{2j} = J_0(\sqrt{\ln x^{-4}}).$$

PROPOSITION 3.4. If $f \in H^2$ extends to be analytic on a neighborhood of 1, then $\Phi: (0,1] \to \mathbb{C}$ defined by

$$\Phi(x) = U^* f(x), \quad x \in (0,1],$$

extends holomorphically to $\mathbb{C}\setminus(-\infty, 0]$. Furthermore, if f_{-1} denotes the function defined by $f_{-1}(z) = f(-z)$, then

$$U^* f_{-1}(x) = \Phi(1/x), \quad x \in (0, 1].$$

Proof. Observe that as f is assumed to be analytic on a neighborhood of 1, the Cauchy–Hadamard radius of convergence formula implies that F, defined by the formula

$$F(w) = \sum_{k=0}^{\infty} f^{(k)}(1) \frac{w^k}{(k!)^2},$$

is an entire function. Consequently, as Lemma 3.3 implies that

$$\Phi(x) = U^* f(x) = F(\ln x)$$

and $\ln x$ extends holomorphically to $\mathbb{C} \setminus (-\infty, 0]$, also Φ extends holomorphically to $\mathbb{C} \setminus (-\infty, 0]$.

To see the second assertion of the lemma, note using Lemma 3.3 (with f replaced with f_{-1}) that

$$U^* f_{-1}(x) = \sum_{k=0}^{\infty} f_{-1}^{(k)}(1) \frac{(\ln x)^k}{(k!)^2} = \sum_{k=0}^{\infty} f^{(k)}(1) \frac{(\ln x)^k}{(k!)^2}.$$

4. Proofs and auxiliary results

4.1. Convergence of subspaces

PROPOSITION 4.1. Let \mathcal{M}_n be a sequence of closed subspaces of a Hilbert space \mathcal{H} . The following are equivalent.

- (i) $\mathcal{M}_n \to \mathcal{M}$.
- (ii) $P_{\mathcal{M}_n}P_{\mathcal{M}} \to P_{\mathcal{M}}$ in the strong operator topology on $\mathcal{B}(\mathcal{H})$, and there is no larger space $\mathcal{N} \supseteq \mathcal{M}$ such that $P_{\mathcal{M}_n}P_{\mathcal{N}} \to P_{\mathcal{N}}$.
- (iii) $P_{\mathcal{M}_n}P_{\mathcal{M}} \to P_{\mathcal{M}}$ in the weak operator topology on $\mathcal{B}(\mathcal{H})$, and there is no larger space $\mathcal{N} \supseteq \mathcal{M}$ such that $P_{\mathcal{M}_n}P_{\mathcal{N}} \to P_{\mathcal{N}}$.

Proof. (i) \Rightarrow (ii). Let $f \in \mathcal{H}$. Let $P_{\mathcal{M}}f = g$. By (i), there exist $g_n \in \mathcal{M}_n$ such that $||g_n - g|| \to 0$. Therefore

$$\|P_{\mathcal{M}_n}g - P_{\mathcal{M}}g\| \le \|g_n - g\| \to 0,$$

so $P_{\mathcal{M}_n}P_{\mathcal{M}} \to P_{\mathcal{M}}$ SOT.

If a space $\mathcal{N} \supseteq \mathcal{M}$ existed for which $P_{\mathcal{M}_n} P_{\mathcal{N}} \to P_{\mathcal{N}}$, let $h \in \mathcal{N} \ominus \mathcal{M}$. Let $h_n = P_{\mathcal{M}_n} h$. Then $h_n \in \mathcal{M}_n$ and $h_n \to h$. By (i), this means $h \in \mathcal{M}$, so h = 0 and $\mathcal{N} = \mathcal{M}$.

(ii) \Rightarrow (i). If $f \in \mathcal{M}$, then $P_{\mathcal{M}_n}f \to f$, so $f \in \lim \mathcal{M}_n$. If there were some $h = \lim g_n$ for a sequence $g_n \in \mathcal{M}_n$, then $\mathcal{N} = \mathcal{M} + \mathbb{C}h$ would satisfy $P_{\mathcal{M}_n}P_{\mathcal{N}} \to P_{\mathcal{N}}$ SOT. So by (ii), this means $h \in \mathcal{M}$, so $\mathcal{M} = \lim \mathcal{M}_n$.

(ii) \Leftrightarrow (iii). This is because a sequence of projections in a Hilbert space converges in the WOT if and only if it converges in the SOT. Indeed, suppose $Q_n \to Q$ WOT, and Q and each Q_n are projections. Then

$$||(Q-Q_n)f||^2 = \langle Qf, f \rangle - 2\operatorname{Re}\langle Q_nf, Qf \rangle + \langle Q_nf, f \rangle \to 0. \blacksquare$$

4.2. Proof of Theorem 1.17. We have already shown that $\{e_n\}$ is an orthonormal basis. Clearly, J is unitary, so must be given by Theorem 1.15 for some c(s) and $\tau(s)$. As J1 = 1, we have $\tau(0) = 0$ and c(0) = 1. As $J^2 = 1$, we have $\tau(\tau(s)) = s$ and $c(\tau(s))c(s) = 1$.

So τ is an automorphism of S that fixes 0 and has period 2. Once we know $\tau'(0)$, this will determine τ uniquely. To calculate $\tau'(0)$, note that

$$e_1(x) = 1 + \frac{\partial}{\partial s} x^s \bigg|_{s=0}$$

Therefore

$$Je_1(x) = -1 + \frac{\partial}{\partial s} [c(s)x^{\tau(s)}]_{s=0} = 1 + c'(0) + c(0)\tau'(0)\ln x.$$

This yields $\tau'(0) = -1$, so (1.4) hold.

4.3. Proof of Corollary 1.12

LEMMA 4.2. Suppose T is a bounded monomial operator given by

(4.1)
$$Tx^s = c(s)x^{\tau(s)}$$

Then τ is a holomorphic function from S to S, and c is a holomorphic function on S.

Proof. The map $s \mapsto x^s$ is a holomorphic map from S to $L^2([0, 1])$. Therefore, for each $t \in S$, the map

$$s \mapsto \langle x^s, T^*x^t \rangle = \frac{c(s)}{1 + \tau(s) + \overline{t}}$$

is holomorphic. Letting t = 0 and 1 and taking the quotient, we find that

$$\frac{2+\tau(s)}{1+\tau(s)} = 1 + \frac{1}{1+\tau(s)}$$

is a meromorphic function of s. Hence τ is meromorphic in \mathbb{S} . Moreover, τ cannot have a pole, since otherwise in a neighborhood of this pole it would take on all values in a neighborhood of ∞ , including ones not in \mathbb{S} . Therefore τ is holomorphic, and consequently so is c(s) since we have

$$c(s) = (1 + \tau(s)) \langle x^s, T^*1 \rangle. \blacksquare$$

Proof of Corollary 1.12. If T maps Lat(H) to Lat(H), it must be a monomial operator, since each monomial function spans a one-dimensional H-invariant subspace.

Conversely, suppose T is a monomial operator given by (4.1), and $\mathcal{M} \in \text{Lat}(H)$. By Lemma 4.2, the function τ is a holomorphic map from \mathbb{S} to \mathbb{S} . Define the function $\phi \in \text{H}^{\infty}(\mathbb{D}(1,1))$ by

$$\phi(z) = \frac{1}{1 + \tau\left(\frac{1-z}{z}\right)}$$

Then for every $s \in \mathbb{S}$ we have

$$\phi\left(\frac{1}{1+s}\right) = \frac{1}{1+\tau(s)}$$

Therefore $HT = T\phi(H)$, since they agree on all monomials, so $HT\mathcal{M} = T\phi(H)\mathcal{M} \subseteq T\mathcal{M}$, as required.

REMARK 4.3. We used the fact that if $\phi \in \mathrm{H}^{\infty}(\mathbb{D}(1,1))$, then $\phi(H)$ is a bounded operator. We define $\phi(H)$ to be the monomial operator

$$\phi(H): x^s \mapsto \phi\left(\frac{1}{1+s}\right)x^s.$$

This will be bounded by M if and only if $M^2 - \phi(H)^* \phi(H) \ge 0$, which is equivalent to

(4.2)
$$\frac{M^2 - \phi\left(\frac{1}{1+s}\right)\overline{\phi\left(\frac{1}{1+t}\right)}}{1+s+\overline{t}} \ge 0.$$

The fact that (4.2) is equivalent to the assertion that ϕ has norm at most M in $\mathrm{H}^{\infty}(\mathbb{D}(1,1))$ is, after a change of variables, the content of Pick's theorem [26].

4.4. Proof of Theorem 1.18. Let T be a unitary monomial operator, given by

(4.3)
$$Tx^s = c(s)x^{\tau(s)}.$$

By Theorem 1.15, we know that τ is a holomorphic automorphism of S. It is well-known that holomorphic automorphisms of the upper half-plane are given by linear fractional transformations with coefficients from $SL(2, \mathbb{R})$. So any holomorphic automorphism of S is of the form

(4.4)
$$\tau(s) = \frac{A(s+\frac{1}{2}) - iB}{iC(s+\frac{1}{2}) + D} - \frac{1}{2};$$

where $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ is in SL(2, \mathbb{R}). Let $\sigma(s) = \frac{-s}{1+2s}$.

CASE (i): $\tau(\infty) = \infty$. Then τ is of the form $\tau(s) = \alpha s + \beta + i\gamma$, where $\alpha > 0, \beta, \gamma \in \mathbb{R}$, and $\beta = \frac{\alpha - 1}{2}$. As T is given by (4.3) and

$$c(s) = c_0 \frac{1 + \bar{\beta} + \alpha s + \beta}{1 + s} = c_0 \alpha,$$

we have

$$T: x^s \mapsto c_0 \alpha x^\beta x^{\alpha s}.$$

Hence

$$T: f(x) \mapsto c_0 \alpha x^\beta f(x^\alpha).$$

Therefore

$$TL^{2}([\rho, 1]) = L^{2}[\rho^{\frac{1}{\alpha}}, 1] = \mathcal{A}_{1, \frac{1}{2\alpha} \log \frac{1}{\rho}}.$$

CASE (ii): $\tau(\infty) = -\frac{1}{2}$. Then $\sigma \circ \tau(\infty) = \infty$. So by Case (i), we have $JTL^2([\rho, 1]) = A_{1,w}$

for some w. Therefore

$$TL^2([\rho, 1]) = A_{-1,w}.$$

CASE (iii): $\tau(\infty) = -\frac{1}{2} + i\delta$. Then

$$M_{x^{-i\delta}}T: x^s \mapsto \tilde{c}(s)x^{\tilde{\tau}(s)},$$

where $\tilde{\tau}(\infty) = -\frac{1}{2}$. By Case (ii), we have

$$TL^2([\rho, 1]) = M_{x^{i\delta}} \mathcal{A}_{-1,\wp} = \mathcal{A}_{\tau,w},$$

for $\tau = \frac{2i\delta+1}{2i\delta-1}$ and $w = \frac{\wp}{1+4\delta^2}$.

Uniqueness of representation: We need to show that if $\mathcal{A}_{\tau,w} = \mathcal{A}_{\tau',w'}$, then $\tau' = \tau$ and w' = w. Observe that if U is a unitary, and $P_{\mathcal{M}}f = g$, then $P_{\mathcal{U}\mathcal{M}}Uf = Ug$.

Let us calculate $P_{\mathcal{A}_{\tau,w}} x^s$. If $\tau = 1$, then

$$P_{\mathcal{A}_{1,w}}x^s = \chi_{[e^{-2w},1]}x^s,$$

and

(4.5)
$$\|P_{\mathcal{A}_{1,w}}x^s\|^2 = \frac{1}{1+2\operatorname{Re} s} [1-e^{-2w(1+2\operatorname{Re} s)}].$$

Otherwise, $\tau = \frac{2i\delta + 1}{2i\delta - 1}$ for some $\delta \in \mathbb{R}$. Then

$$P_{\mathcal{A}_{\tau,w}}x^s = M_{x^{i\delta}}JP_{\mathcal{A}_{1,\wp}}Jx^{s-i\delta}.$$

From (1.4),

$$Jx^{s-i\delta} = \frac{1}{1+2s-2i\delta}x^{\frac{-s+i\delta}{1+2s-2i\delta}}.$$

Therefore

$$\begin{aligned} \|P_{\mathcal{A}_{\tau,w}}x^{s}\|^{2} &= \|P_{\mathcal{A}_{1,\wp}}Jx^{s-i\delta}\|^{2} \\ &= \frac{1}{|1+2s-2i\delta|^{2}} \int_{e^{-2\wp}}^{1} x^{2\operatorname{Re}\frac{-s+i\delta}{1+2s-2i\delta}} dx \end{aligned}$$

When s = u + iv, this gives

(4.6)
$$\|P_{\mathcal{A}_{\tau,w}}x^s\|^2 = \frac{1}{1+2u} \Big[1 - e^{-2\wp \frac{1+2u}{(1+2u)^2 + 4(\delta-v)^2}}\Big].$$

Comparing (4.5) and (4.6), we see that τ and w are completely determined by $\|P_{\mathcal{A}_{\tau,w}}x^s\|^2$.

5. Proofs using Hardy space theory. Although Theorems 1.14 and 1.19 are stated without using the language of Hardy spaces, the authors do not know how to prove them directly.

5.1. Proof of 1.14. Let \mathcal{M} be in Lat(H). Define a sequence $S \subset \mathbb{S}$ by $S = \{s \mid \langle f, x^s \rangle = 0 \ \forall f \in \mathcal{M}\}$. The number s will occur in S with multiplicity m where m is the largest number such that $\mathcal{M} \perp \{x^s, (\ln x)x^s, \ldots, (\ln x)^{m-1}x^s\}$. Let $\mathcal{M}_0 = \mathcal{M}(S)$.

To see that $\mathcal{M} = \overline{\mathcal{M}_0 + \mathcal{M}_1}$ for some singular space \mathcal{M}_1 , we use the Sarason transform to move to H^2 . Then \mathcal{M} becomes $(BSH^2)^{\perp}$, where B is a Blaschke product and S is a singular inner function. As $BSH^2 = BH^2 \cap SH^2$, we have

$$(BSH^2)^{\perp} = \overline{(BH^2)^{\perp} + (SH^2)^{\perp}},$$

and \mathcal{M}_1 is the inverse Sarason transform of $(SH^2)^{\perp}$.

5.2. Proof of 1.19. Let $S_{\tau,w}$ denote the singular inner function

$$S_{\tau,w}(z) = \exp\left(-w\frac{\tau+z}{\tau-z}\right).$$

Let $U: L^2 \to H^2$ be the Sarason transform. By Lemma 2.9 we have

$$U\mathcal{A}_{1,w}U^* = (S_{1,w}\mathrm{H}^2)^{\perp}.$$

18

We wish to extend this to other values of τ .

Lemma 5.1.

(5.1)
$$UM_{x^{ic}}U^*(S_{-1,\wp}) = FS_{\tau,w},$$

where $\tau = \frac{2ic+1}{2ic-1}, w = \frac{1}{1+4c^2}\wp$ and $F(z) = \exp(-2icw)\frac{1}{1+ic-icz}$

Proof. Observe first that $\widehat{M_{x^{-ic}}}$ is a unitary operator that takes k_{α} to $\overline{\phi(\alpha)}k_{\psi(\alpha)}$, where

$$\phi(z) = \frac{1}{1 + ic - icz}, \quad \psi(z) = \frac{(1 - ic)z + ic}{1 + ic - icz}.$$

Therefore

$$\widehat{M_{x^{-ic}}} = C_{\psi}^* M_{\phi}^*,$$

and so

(5.2)
$$\widehat{M_{x^{ic}}} = M_{\phi} C_{\psi}.$$

We have

$$C_{\psi}S_{-1,\wp}(z) = \exp\left(\wp \; \frac{-1+\psi(z)}{-1-\psi(z)}\right).$$

A calculation shows that

$$\frac{-1+\psi(z)}{-1-\psi(z)} = \frac{1}{1+4c^2} \frac{\tau+z}{\tau-z} - \frac{2ic}{1+4c^2}$$

Therefore $C_{\psi}S_{-1,\wp}(z)$ is a unimodular constant times $S_{\tau,w}$, and (5.1) holds.

LEMMA 5.2.

(5.3)
$$U\mathcal{A}_{\tau,w}U^* = (S_{\tau,w}\mathrm{H}^2)^{\perp}$$

Proof. We have already proved the case $\tau = 1$, so assume $\tau \neq 1$. Consider next $\tau = -1$. Then

$$U\mathcal{A}_{-1,w}U^* = UJ\mathcal{A}_{1,w}U^* = UJU^*U\mathcal{A}_{1,w}U^* = UJU^*(S_{1,w}\mathrm{H}^2)^{\perp}.$$

As $UJU^*f(z) = f(-z)$, we have

$$UJU^*(S_{1,w}H^2) = (S_{-1,w}H^2),$$

 \mathbf{SO}

$$UJU^*(S_{1,w}\mathrm{H}^2)^{\perp} = (S_{-1,w}\mathrm{H}^2)^{\perp}.$$

For $\tau \neq \pm 1$, we have, with ϕ and ψ as in (5.2) and F as in Lemma 5.1,

$$U\mathcal{A}_{\tau,w}^{\perp}U^* = UM_{x^{ic}}J\mathcal{A}_{1,\wp}^{\perp}U^* = UM_{x^{ic}}U^*(S_{-1,\wp}\mathrm{H}^2)$$
$$= \{\phi(z)F(z)S_{\tau,w}(z)h(\psi(z)) \mid h \in \mathrm{H}^2\}.$$

As F and ϕ are outer and ψ is an automorphism of \mathbb{D} , this proves

$$U\mathcal{A}_{\tau,w}^{\perp}U^* = S_{\tau,w}\mathrm{H}^2,$$

and hence (5.3).

Proof of Theorem 1.19. From Lemma 5.2, we have, for distinct points τ_k ,

$$U\mathcal{M}\left(\sum_{k=1}^{n} w_k \delta_{\tau_k}\right) U^* = \sum_{k=1}^{n} (S_{\tau_k, w_k} \mathrm{H}^2)^{\perp} = \left(\left(\prod_{k=1}^{n} S_{\tau_k, w_k}\right) \mathrm{H}^2\right)^{\perp}.$$

Suppose that $\mu_n \to \mu$ weak-*, where μ and each μ_n are singular. Define singular inner functions by

$$\varphi_n(z) = \exp\left[-\int \frac{e^{i\theta} + z}{e^{i\theta} - z} d\mu_n(\theta)\right], \quad \varphi(z) = \exp\left[-\int \frac{e^{i\theta} + z}{e^{i\theta} - z} d\mu(\theta)\right].$$

Then $\|\varphi_n - \varphi\|_{H^2} \to 0$. Indeed, φ_n tends to φ weakly in H^2 , since the functions all have norm 1 and converge pointwise on \mathbb{D} . Therefore

$$\|\varphi_n - \varphi\|^2 = 2 - 2 \operatorname{Re}\langle\varphi_n, \varphi\rangle \to 0$$

This means that not only do the Toeplitz operators $T_{\overline{\varphi}_n}$ converge to $T_{\overline{\varphi}}$ in the strong operator topology, but $T_{\varphi_n}T_{\overline{\varphi}_n}$ converges to $T_{\varphi}T_{\overline{\varphi}}$ in the SOT. This is proved in [25, p. 34]; for the convenience of the reader, we include the proof. Let $f \in \mathrm{H}^2$. Then

$$\begin{aligned} \|T_{\varphi_n}T_{\bar{\varphi}_n}f - T_{\varphi}T_{\overline{\varphi}}f\| &\leq \|T_{\varphi_n}(T_{\overline{\varphi}_n} - T_{\overline{\varphi}})f\| + \|(T_{\varphi_n} - T_{\varphi})T_{\overline{\varphi}}f\| \\ &\leq \sup_n \|\varphi_n\|_{H^{\infty}} \|(T_{\overline{\varphi}_n} - T_{\overline{\varphi}})f\| + \left(\int |\varphi_n - \varphi|^2 |T_{\bar{\varphi}}f|^2\right)^{1/2}. \end{aligned}$$

The first term tends to zero because $T_{\overline{\varphi}_n}$ tends to $T_{\overline{\varphi}}$ in the SOT, and the second term tends to zero because φ_n tends to φ in measure and $|\varphi_n - \varphi| \leq 2$. As $T_{\varphi_n} T_{\overline{\varphi}_n}$ is the projection onto $(\varphi_n H^2)^{\perp}$, this means by Proposition 4.1 that the spaces $(\varphi_n H^2)^{\perp}$ converge to $(\varphi H^2)^{\perp}$. Applying the inverse Sarason transform, we conclude that $\mathcal{M}(\mu_n)$ converges to $\mathcal{M}(\mu)$.

Acknowledgements. J.M. was partially supported by National Science Foundation Grant DMS 2054199.

References

- J. Agler and J. E. McCarthy, Beurling's theorem for the Hardy operator on L²[0, 1], preprint, 2022.
- J. Agler and J. E. McCarthy, A generalization of Hardy's operator and an asymptotic Müntz-Szász theorem, Expo. Math. 40 (2022), 920–930.
- [3] J. Agler and J. E. M^cCarthy, *Monomial operators*, Acta Sci. Math. (Szeged) 88 (2022), 371–381.
- [4] J. M. Almira, Müntz type theorems. I, Surv. Approx. Theory 3 (2007), 152–194.
- [5] W. Arendt, I. Chalendar, M. Kumar, and S. Srivastava, Asymptotic behaviour of the powers of composition operators on Banach spaces of holomorphic functions, Indiana Univ. Math. J. 67 (2018), 1571–1595.
- [6] A. Beurling, On two problems concerning linear transformations in Hilbert space, Acta Math. 81 (1948), 239–255.

- [7] P. Borwein and T. Erdélyi, *Polynomials and Polynomial Inequalities*, Grad. Texts in Math. 161, Springer, New York, 1995.
- [8] P. S. Bourdon and S. K. Narayan, Normal weighted composition operators on the Hardy space H²(U), J. Math. Anal. Appl. 367 (2010), 278–286.
- [9] M. S. Brodskii, On a problem of I. M. Gel'fand, Uspekhi Mat. Nauk (N.S.) 12 (1957), no. 2 (74), 129–132 (in Russian).
- [10] A. Brown, P. R. Halmos, and A. L. Shields, *Cesàro operators*, Acta Sci. Math. (Szeged) 26 (1965), 125–137.
- [11] I. Chalendar and J. R. Partington, Weighted composition operators: isometries and asymptotic behaviour, J. Operator Theory 86 (2021), 189–201.
- [12] J. A. Clarkson and P. Erdős, Approximation by polynomials, Duke Math. J. 10 (1943), 5–11.
- [13] C. C. Cowen and E. A. Gallardo-Gutiérrez, A new class of operators and a description of adjoints of composition operators, J. Funct. Anal. 238 (2006), 447–462.
- [14] C. C. Cowen and E. Ko, Hermitian weighted composition operators on H², Trans. Amer. Math. Soc. 362 (2010), 5771–5801.
- [15] W. F. Donoghue, Jr., The lattice of invariant subspaces of a completely continuous quasi-nilpotent transformation, Pacific J. Math. 7 (1957), 1031–1035.
- [16] T. Erdélyi and W. B. Johnson, The "full Müntz theorem" in $L_p[0,1]$ for 0 , J. Anal. Math. 84 (2001), 145–172.
- [17] F. Forelli, The isometries of H^p, Canad. J. Math. 16 (1964), 721–728.
- [18] E. Fricain and P. Lefèvre, L²-Müntz spaces as model spaces, Complex Anal. Oper. Theory 13 (2019), 127–139.
- [19] S. R. Garcia, J. Mashreghi, and W. T. Ross. Introduction to Model Spaces and Their Operators, Cambridge Stud. Adv. Math. 148, Cambridge Univ. Press, Cambridge, 2016.
- [20] G. Gunatillake, Compact weighted composition operators on the Hardy space, Proc. Amer. Math. Soc. 136 (2008), 2895–2899.
- [21] G. H. Hardy, Note on a theorem of Hilbert, Math. Z. 6 (1920), 314–317.
- [22] T. Kriete and J. Moorhouse, Linear relations in the Calkin algebra for composition operators, Trans. Amer. Math. Soc. 359 (2007), 2915–2944.
- [23] N. Kruglyak, L. Maligranda, and L.-E. Persson, Structure of the Hardy operator related to Laguerre polynomials and the Euler differential equation, Rev. Mat. Complut. 19 (2006), 467–476.
- [24] Ch. H. Müntz, Uber den Approximationssatz von Weierstraß, in: H. A. Schwartz-Festschrift, Springer, Berlin, 1914, 303–312.
- [25] N. K. Nikol'skiĭ, Treatise on the Shift Operator. Spectral Function Theory, Grundlehren Math. Wiss. 273, Springer, Berlin, 1986.
- [26] G. Pick, Über die Beschränkungen analytischer Funktionen, welche durch vorgegebene Funktionswerte bewirkt werden, Math. Ann. 77 (1915), 7–23.
- [27] D. Sarason, A remark on the Volterra operator, J. Math. Anal. Appl. 12 (1965), 244–246.
- [28] O. Szász, Über die Approximation stetiger Funktionen durch lineare Aggregate von Potenzen, Math. Ann. 77 (1916), 482–496.
- [29] O. Szász, On closed sets of rational functions, Ann. Mat. Pura Appl. (4) 34 (1953), 195–218.

Jim Agler Department of Mathematics University of California at San Diego La Jolla, CA 92093-0112, U.S.A. E-mail: jagler@ucsd.edu John E. McCarthy Department of Mathematics Washington University in St. Louis St. Louis, MO 63130-4899, U.S.A. E-mail: mccarthy@wustl.edu