

Exam 2

Math 309

November 9, 2016

You may use a scientific calculator, but no notes, graphing calculators, or other electronic devices allowed. Fill out your scantron cards with your name, *including your official "preferred" name*, and your ID number.

Part I: Multiple Choice (30) Each problem in this section is worth five points.

1. If $A = \begin{bmatrix} -12 & 0 & 13 & 42 & 3 & -62 & 0 & 98 & 0 & 12 \\ 0 & -2 & 65 & 55 & 55 & 48 & 7 & 48 & 19 & 0 \\ 0 & 0 & 14 & 14 & 15 & 16 & -3 & 223 & 23 & 42 \\ 0 & 0 & 0 & 8 & 42 & 62 & 27 & -28 & 12 & -10 \\ 0 & 0 & 0 & 0 & -51 & -13 & 17 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -6 & 23 & 86 & -9 & \frac{2}{3} \\ 0 & 0 & 0 & 0 & 0 & 0 & 12 & 0 & 0 & \pi \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 11 & 2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -4 & 101 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 100 \end{bmatrix}$, then

- A) $\det A < 0$ B) $\det A = 0$ C) $\det A > 0$ D) $\det A$ is not a real number
E) none of the above
2. If A and B are 3×3 matrices, $\det(A) = 3$ and $\det(B) = 2$, what is $\det((2A)B^{-1})$?
A) 1 B) -1 C) 4 D) 12 E) -12 F) 24
G) -24 H) none of the above
3. Let A be an $m \times n$ matrix such that there is one free variable in the system $A\mathbf{x} = \mathbf{0}$. What is the rank of A ?
A) m B) n C) $n - m$ D) $m-1$ E) $n-1$ F) $n-m+1$
G) $m + n + 1$ H) none of the above

4. Find all a such that $\det \begin{bmatrix} a & 0 & 2 & 4 \\ 0 & a & 0 & 0 \\ 0 & 1 & 4 & 5 \\ 0 & 3 & 3 & 4 \end{bmatrix} = \det \begin{bmatrix} 4 & 0 & 0 \\ -2 & a & 1 \\ 3 & 1 & 1 \end{bmatrix}$

- A) all real numbers a B) no real numbers a C) 0, -2 D) -2, 2
 E) 2 F) 4 G) 2, 4 H) none of the above

5. A linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is given by $T(\mathbf{x}) = A\mathbf{x}$ for a certain matrix A . If S a circle of radius 2, the image under the transformation $T(S)$ is an ellipse with area 8π . Which of the following matrices might be A ?

A) $\begin{bmatrix} 517 & 330 \\ -123 & 223 \end{bmatrix}$ B) $\begin{bmatrix} 1 & 0 \\ -1 & 2 \end{bmatrix}$ C) $\begin{bmatrix} 1 & 1 \\ -2 & 2 \end{bmatrix}$ D) $\begin{bmatrix} 2 & 0 \\ -12 & -3 \end{bmatrix}$

E) $\begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$ F) $\begin{bmatrix} 6 & 0 \\ -1 & 6 \end{bmatrix}$ G) $\begin{bmatrix} 2 & 3 \\ -1 & 2 \end{bmatrix}$ H) $\begin{bmatrix} 6 & 0 \\ 0 & 6 \end{bmatrix}$

I) None of the above

6. For an $m \times n$ matrix A , which of the following is not equal to the rest?

- A) the rank of A B) the dimension of the column space of A C) the dimension of the row space of A
 D) the dimension of the null space of A
 E) the number of pivot columns of A

14. The number of *free variables* in the equation $A\mathbf{x} = \mathbf{b}$ equals the dimension $\text{Nul}(A)$.

A) True

B) False

15. The vectors $\begin{bmatrix} 1 \\ 10 \\ -1 \\ 3 \\ 7 \end{bmatrix}$, $\begin{bmatrix} 5 \\ 0 \\ 3 \\ 1 \\ 2 \end{bmatrix}$, $\begin{bmatrix} 1 \\ 1 \\ 1 \\ 2 \\ 1 \end{bmatrix}$, and $\begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{bmatrix}$ span \mathbb{R}^5 .

A) True

B) False

16. If E_1, E_2, \dots, E_n are elementary matrices and $E_1 \cdot E_2 \cdot \dots \cdot E_n A = B$, then $\det A = \det B$.

A) True

B) False

Part III: Short Answer(20)

You do not need to show any work for this part, just answer each question or fill in the blank.

17. Let $\mathbf{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ (in coordinates of the standard basis for \mathbb{R}^2) and let $\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} -1 \\ 2 \end{bmatrix} \right\}$ be an alternate basis. Find $[\mathbf{x}]_{\mathcal{B}}$.

18. Let $u_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$, $u_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$, $v_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$, and $v_2 = \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}$, and let $U = \text{span}\{u_1, u_2\}$, $V = \text{span}\{v_1, v_2\}$. Find a nonzero vector which is in the subspace $U \cap V$.

19. The matrix $A = \begin{bmatrix} -2 & -5 & 8 & 0 & -17 \\ 1 & 3 & -5 & 1 & 5 \\ 3 & 11 & -19 & 7 & 1 \\ 1 & 7 & -13 & 5 & -3 \end{bmatrix}$ reduces to $\begin{bmatrix} 1 & 3 & -5 & 1 & 5 \\ 0 & 1 & -2 & 2 & -7 \\ 0 & 0 & 0 & -4 & 20 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$.

(a) $\text{rank}(A) =$

Give a basis for:

(b) $\text{row}(A)$

(c) $\text{Col}(A)$

(d) $\text{Nul}(A)$

Part IV: Longer Answer(20)

20. Clearly state whether each statement is true or false. If the statement is true, give a proof. If the statement is false, give a counterexample or a reason why it cannot be true.

- (a) If A is a 2×2 matrix, then the determinant of A is equal to the determinant of its transpose. That is,

$$\det(A^T) = \det(A).$$

- (b) If A is an $n \times n$ matrix, then the determinant of A is equal to the determinant of its inverse. That is,

$$\det(A^{-1}) = \det(A).$$

- (c) If H and K are subspaces of a vector space V , then the intersection $H \cap K$ is also a subspace.

21. For each matrix below, find the determinant of A.

$$(a) A = \begin{bmatrix} 12 & 4 \\ -1 & 2 \end{bmatrix}$$

$$(b) A = \begin{bmatrix} 10 & 0 & 0 & 5 \\ 1 & 2 & 0 & 0 \\ 4 & -9 & 5 & 5 \\ 4 & -9 & 5 & -2 \end{bmatrix}$$

$$(c) A = \begin{bmatrix} \frac{1}{2} & \frac{5}{8} & \frac{1}{3} \\ 12 & 15 & 8 \\ \frac{1}{30} & 32 & 13 \end{bmatrix}$$