

# Chapter III

## Topological Spaces

### 1. Introduction

In Chapter I we looked at properties of sets, and in Chapter II we added some additional structure to a set – a distance function  $d$  – to create a pseudometric space. We then saw some of the most basic fundamental definitions and properties of pseudometric spaces. There is much more, and some of the most useful and interesting properties of pseudometric spaces will be discussed in Chapter IV. But in Chapter III we look at an important generalization.

We observed, early in Chapter II, that the idea of continuity (in calculus) depends on talking about “nearness,” so we used a distance function  $d$  to make the idea of “nearness” precise. In that way, we were able to extend the definition of continuity from  $\mathbb{R}^n$  to pseudometric spaces. The distance function  $d$  also led us to the idea of open sets in a pseudometric space. From there we developed properties of closed sets, closures, interiors, frontiers, dense sets, continuity, and sequential convergence.

An important observation in Chapter II was that open (or closed) sets are really all that we require to talk about many of these ideas. In other words, we can often do what's necessary using the open sets without knowing which specific  $d$  generated the open sets: the topology  $\mathcal{T}_d$  is what really matters. For example,  $\text{int } A$  is defined in  $(X, d)$  in terms of the open sets, so  $\text{int } A$  doesn't change if  $d$  is replaced with a different but equivalent metric  $d'$  – one that generates the same open sets. Similarly, changing  $d$  to an equivalent metric  $d'$  doesn't affect closures, continuity, or convergent sequences. In summary: for many purposes  $d$  is logically unnecessary once  $d$  has done the job of creating the topology  $\mathcal{T}_d$  (although having  $d$  might still be convenient).

This suggests a way to generalize our work. For a particular set  $X$  we can simply assign a topology – that is, a collection of “open” sets given without any mention of a pseudometric that might have generated them. Of course when we do this, we want the “open sets” to “behave the way open sets should behave” as described in Theorem II.2.11. This leads us to the definition of a topological space.

### 2. Topological Spaces

**Definition 2.1** A topology  $\mathcal{T}$  on a set  $X$  is a collection of subsets of  $X$  such that

- i)  $\emptyset, X \in \mathcal{T}$
- ii) if  $O_\alpha \in \mathcal{T}$  for each  $\alpha \in A$ , then  $\bigcup_{a \in A} O_a \in \mathcal{T}$
- iii) if  $O_1, \dots, O_n \in \mathcal{T}$ , then  $O_1 \cap \dots \cap O_n \in \mathcal{T}$ .

A set  $O \subseteq X$  is called open if  $O \in \mathcal{T}$ . The pair  $(X, \mathcal{T})$  is called a topological space.

*Sometimes we will just refer to “a topological space  $X$ .” In that case, it is assumed that there is some topology  $\mathcal{T}$  on  $X$  but, for short, we are just not bothering to write down the “ $\mathcal{T}$ ”)*

We emphasize that in a topological space there is no distance function  $d$ : therefore phrases like “distance between two points” and “ $\epsilon$ -ball” make no sense in  $(X, \mathcal{T})$ . There is no preconceived idea about what “open” means: to say “ $O$  is open” means nothing more or less than “ $O \in \mathcal{T}$ .”

In a topological space  $(X, \mathcal{T})$ , we can go on to define closed sets and isolated points just as we did in pseudometric spaces.

**Definition 2.2** A subset  $F$  in  $(X, \mathcal{T})$  is called closed if  $X - F$  is open, that is, if  $X - F \in \mathcal{T}$ .

**Definition 2.3** A point  $a \in (X, \mathcal{T})$  is called isolated if  $\{a\}$  is open, that is, if  $\{a\} \in \mathcal{T}$ .

The proof of the following theorem is the same as it was for pseudometric spaces; we just take complements and apply properties of open sets.

**Theorem 2.4** In any topological space  $(X, \mathcal{T})$

- i)  $\emptyset$  and  $X$  are closed
- ii) if  $F_\alpha$  is closed for each  $\alpha \in A$ , then  $\bigcap_{\alpha \in A} F_\alpha$  is closed
- iii) if  $F_1, \dots, F_n$  are closed, then  $\bigcup_{i=1}^n F_i$  is closed.

More informally, ii) and iii) state that intersections and finite unions of closed sets are closed.

**Proof** Read the proof for Theorem II.4.2. •

For a particular topological space  $(X, \mathcal{T})$ , it might or might not be possible to find a pseudometric  $d$  on  $X$  that “creates” this topology – that is, one for which  $\mathcal{T}_d = \mathcal{T}$ .

**Definition 2.5** A topological space  $(X, \mathcal{T})$  is called pseudometrizable if there exists a pseudometric  $d$  on  $X$  for which  $\mathcal{T}_d = \mathcal{T}$ . If  $d$  is a metric, then  $(X, \mathcal{T})$  is called metrizable.

### Examples 2.6

1) Suppose  $X$  is a set and  $\mathcal{T} = \{\emptyset, X\}$ .  $\mathcal{T}$  is called the trivial topology on  $X$  and it is the smallest possible topology on  $X$ .  $(X, \mathcal{T})$  is called a trivial topological space. The only open (or closed) sets are  $\emptyset$  and  $X$ . If we put the trivial pseudometric  $d$  on  $X$ , then  $\mathcal{T}_d = \mathcal{T}$ . So a trivial topological space turns out to be pseudometrizable.

At the opposite extreme, suppose  $\mathcal{T} = \mathcal{P}(X)$ . Then  $\mathcal{T}$  is called the discrete topology on  $X$  and it is the largest possible topology on  $X$ .  $(X, \mathcal{T})$  is called a discrete topological space. Every subset is open (and also closed). Every point of  $X$  is isolated. If we put the discrete unit metric  $d$  (or any equivalent metric) on  $X$ , then  $\mathcal{T}_d = \mathcal{T}$ . So a discrete topological space is metrizable.

2) Suppose  $X = \{0, 1\}$  and let  $\mathcal{T} = \{\emptyset, \{1\}, X\}$ .  $(X, \mathcal{T})$  is a topological space called Sierpinski space. In this case it is not possible to find a pseudometric  $d$  on  $X$  for which  $\mathcal{T}_d = \mathcal{T}$ , so Sierpinski space is not pseudometrizable. To see this, consider any pseudometric  $d$  on  $X$ .

If  $d(0, 1) = 0$ , then  $d$  is the trivial pseudometric on  $X$  and  $\{\emptyset, X\} = \mathcal{T}_d \neq \mathcal{T}$ .

If  $d(0, 1) = \delta > 0$ , then the open ball  $B_\delta(0) = \{0\} \in \mathcal{T}_d$ , so  $\mathcal{T}_d \neq \mathcal{T}$ .

(In this case  $\mathcal{T}_d$  is actually the discrete topology:  $d$  is just a rescaling of the discrete unit metric.)

Another possible topology on  $X = \{0, 1\}$  is  $\mathcal{T}' = \{\emptyset, \{0\}, X\}$ , although  $(X, \mathcal{T})$  and  $(X, \mathcal{T}')$  seem very much alike: both are two-point spaces, each with exactly one isolated point. One space can be obtained from the other simply renaming “0” and “1” as “1” and “0” respectively. Such “topologically identical” spaces are called “homeomorphic.” (We will give a precise definition later in this chapter.)

3) For a set  $X$ , let  $\mathcal{T} = \{O \subseteq X : O = \emptyset \text{ or } X - O \text{ is finite}\}$ .  $\mathcal{T}$  is a topology on  $X$ :

i) Clearly,  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ .

ii) Suppose  $O_\alpha \in \mathcal{T}$  for each  $\alpha \in A$ . If  $\bigcup_{\alpha \in A} O_\alpha = \emptyset$ , then  $\bigcup_{\alpha \in A} O_\alpha \in \mathcal{T}$ . Otherwise there is at least one  $O_{\alpha_0} \neq \emptyset$ . Then  $X - O_{\alpha_0}$  is finite, so  $X - \bigcup_{\alpha \in A} O_\alpha = \bigcap_{\alpha \in A} (X - O_\alpha) \subseteq X - O_{\alpha_0}$ . Therefore  $X - \bigcup_{\alpha \in A} O_\alpha$  is also finite, so  $\bigcup_{\alpha \in A} O_\alpha \in \mathcal{T}$ .

iii) If  $O_1, \dots, O_n \in \mathcal{T}$  and some  $O_i = \emptyset$ , then  $\bigcap_{i=1}^n O_i = \emptyset$  so  $\bigcap_{i=1}^n O_i \in \mathcal{T}$ . Otherwise each  $O_i$  is nonempty, so each  $X - O_i$  is finite. Then  $(X - O_1) \cup \dots \cup (X - O_n) = X - \bigcap_{i=1}^n O_i$  is finite, so  $\bigcap_{i=1}^n O_i \in \mathcal{T}$ .

In  $(X, \mathcal{T})$ , a set  $F$  is closed iff  $F = \emptyset$  or  $F$  is finite. Because the open sets are  $\emptyset$  and the complements of finite sets,  $\mathcal{T}$  is called the cofinite topology on  $X$ .

If  $X$  is a finite set, then the cofinite topology is the same as the discrete topology on  $X$ . (Why?) If  $X$  is infinite, then no point in  $(X, \mathcal{T})$  is isolated.

Suppose  $X$  is an infinite set with the cofinite topology  $\mathcal{T}$ . If  $U$  and  $V$  are nonempty open sets, then  $X - U$  and  $X - V$  must be finite so  $(X - U) \cup (X - V) = X - (U \cap V)$  is finite. Since  $X$  is infinite, this means that  $U \cap V \neq \emptyset$  (*in fact,  $U \cap V$  must be infinite*). Therefore every pair of nonempty open sets in  $(X, \mathcal{T})$  has nonempty intersection! This shows us that an infinite cofinite space  $(X, \mathcal{T})$  is not pseudometrizable:

i) if  $d$  is the trivial pseudometric on  $X$ , then certainly  $\mathcal{T}_d \neq \mathcal{T}$ , and

ii) if  $d$  is not the trivial pseudometric on  $X$ , then there exist points  $a, b \in X$  for which  $d(a, b) = \delta > 0$ . In that case,  $B_{\delta/2}(a)$  and  $B_{\delta/2}(b)$  would be disjoint nonempty open sets in  $\mathcal{T}_d$ , so  $\mathcal{T}_d \neq \mathcal{T}$ .

4) On  $\mathbb{R}$ , let  $\mathcal{T} = \{(a, \infty) : a \in \mathbb{R}\} \cup \{\emptyset, \mathbb{R}\}$ . It is easy to verify that  $\mathcal{T}$  is a topology on  $\mathbb{R}$ , called the right-ray topology. Is  $(\mathbb{R}, \mathcal{T})$  metrizable or pseudometrizable?

If  $X = \emptyset$ , then  $\mathcal{T} = \{\emptyset\}$  is the only possible topology on  $X$ , and  $\mathcal{T} = \{\emptyset, \{a\}\}$  is the only possible topology on a singleton set  $X = \{a\}$ . But for  $|X| > 1$ , there are many possible topologies on  $X$ . For example, there are four possible topologies on the set  $X = \{a, b\}$ . These are the trivial topology, the discrete topology,  $\{\emptyset, \{a\}, X\}$ , and  $\{\emptyset, \{b\}, X\}$  although, as mentioned earlier, the last two can be considered as “topologically identical.”

If  $\mathcal{T}$  is a topology on  $X$ , then  $\mathcal{T}$  is a collection of subsets of  $X$ , so  $\mathcal{T} \subseteq \mathcal{P}(X)$ . This means that  $\mathcal{T} \in \mathcal{P}(\mathcal{P}(X))$ , so  $|\mathcal{P}(\mathcal{P}(X))| = 2^{|\mathcal{P}(X)|} = 2^{2^{|X|}}$  is an upper bound for the number of possible topologies on  $X$ . For example, there are no more than  $2^{2^7} \approx 3.4 \times 10^{38}$  topologies on a set  $X$  with 7 elements. But this upper bound is actually very crude, as the following table (given without proof) indicates:

$n =  X $	Actual number of distinct topologies on $X$
0	1
1	1
2	4
3	29
4	355
5	6942
6	209527
7	9535241 (many less than $10^{38}$ )

Counting topologies on finite sets is really a question about combinatorics and we will not pursue this topic.

Each concept we defined for pseudometric spaces can be carried over directly to topological spaces if the concept was defined in topological terms – that is, in terms of open (or closed) sets. This applies, for example, to the definitions of interior, closure, and frontier in pseudometric spaces, so these definitions can also be carried over verbatim to a topological space  $(X, \mathcal{T})$ .

**Definition 2.7** Suppose  $A \subseteq (X, \mathcal{T})$ . We define

$$\begin{aligned}\text{int}_X A &= \text{the interior of } A \text{ in } X = \bigcup\{O : O \text{ is open and } O \subseteq A\} \\ \text{cl}_X A &= \text{the closure of } A \text{ in } X = \bigcap\{F : F \text{ is closed and } F \supseteq A\} \\ \text{Fr}_X A &= \text{the frontier (or boundary) of } A \text{ in } X = \text{cl}_X A \cap \text{cl}_X(X - A)\end{aligned}$$

As before, we will drop the subscript “ $X$ ” when the context makes it clear.

The properties for the operators  $\text{cl}$ ,  $\text{int}$ , and  $\text{Fr}$  (except those that mention a pseudometric  $d$  or an  $\epsilon$ -ball) remain true. The proofs in the preceding chapter were deliberately phrased in topological terms so they would carry over to the more general setting of topological spaces.

**Theorem 2.8** Suppose  $A \subseteq (X, d)$ . Then

- 1) a)  $\text{int } A$  is the largest open subset of  $A$  (that is, if  $O$  is open and  $O \subseteq A$ , then  $O \subseteq \text{int } A$ ).  
b)  $A$  is open iff  $A = \text{int } A$  (*Note:  $\text{int } A \subseteq A$  is true for every set  $A$  so we could say:  $A$  is open iff  $A \subseteq \text{int } A$ .*)  
c)  $x \in \text{int } A$  iff there is an open set  $O$  such that  $x \in O \subseteq A$
- 2) a)  $\text{cl } A$  is the smallest closed set containing  $A$  (that is, if  $F$  is closed and  $F \supseteq A$ , then  $F \supseteq \text{cl } A$ ).  
b)  $A$  is closed iff  $A = \text{cl } A$  (*Note:  $A \subseteq \text{cl } A$  is true for every set  $A$ , so we could say:  $A$  is closed iff  $\text{cl } A \subseteq A$ .*)  
c)  $x \in \text{cl } A$  iff for every open set  $O$  containing  $x$ ,  $O \cap A \neq \emptyset$
- 3) a)  $\text{Fr } A$  is closed and  $\text{Fr } A = \text{Fr}(X - A)$ .  
b)  $x \in \text{Fr } A$  iff for every open set  $O$  containing  $x$ ,  $O \cap A \neq \emptyset$  and  $O \cap (X - A) \neq \emptyset$   
c)  $A$  is clopen iff  $\text{Fr } A = \emptyset$ .

*See the proof of Theorem II.4.5*

At this point, we add a few additional facts about these operators. Some of the proofs are left as exercises.

**Theorem 2.9** Suppose  $A, B$  are subsets of a topological space  $(X, \mathcal{T})$ . Then

- 1)  $\text{cl}(A \cup B) = \text{cl}(A) \cup \text{cl}(B)$
- 2)  $\text{cl } A = A \cup \text{Fr } A$
- 3)  $\text{int } A = A - \text{Fr } A = X - \text{cl}(X - A)$
- 4)  $\text{Fr } A = \text{cl } A - \text{int } A$
- 5)  $X = \text{int } A \cup \text{Fr } A \cup \text{int } (X - A)$ , and these 3 sets are disjoint.

**Proof** 1)  $A \subseteq A \cup B$  so, from the definition of closure, we have  $\text{cl } A \subseteq \text{cl}(A \cup B)$ .

Similarly,  $\text{cl } B \subseteq \text{cl}(A \cup B)$ . Therefore  $\text{cl } A \cup \text{cl } B \subseteq \text{cl}(A \cup B)$ .

On the other hand,  $\text{cl } A \cup \text{cl } B$  is the union of two closed sets, so  $\text{cl } A \cup \text{cl } B$  is closed and  $\text{cl } A \cup \text{cl } B \supseteq A \cup B$ , so  $\text{cl } A \cup \text{cl } B \supseteq \text{cl}(A \cup B)$ . (*As an exercise, try proving 1) instead using the characterization of closures given above in Theorem 2.8.2c.*)

Is 1) true if “ $\cup$ ” is replaced by “ $\cap$ ”?

2) Suppose  $x \in \text{cl } A$  but  $x \notin A$ . If  $O$  is any open set containing  $x$ , then  $O \cap A \neq \emptyset$  (because  $x \in \text{cl } A$ ) and  $O \cap (X - A) \neq \emptyset$  (because the intersection contains  $x$ ). Therefore  $x \in \text{Fr } A$ , so  $x \in A \cup \text{Fr } A$ .

Conversely, suppose  $x \in A \cup \text{Fr } A$ . If  $x \in A$ , then  $x \in \text{cl } A$ . And if  $x \notin A$ , then  $x \in \text{Fr } A = \text{cl } A \cap \text{cl } (X - A)$ , so  $x \in \text{cl } A$ . Therefore  $\text{cl } A = A \cup \text{Fr } A$ .

The proofs of 3) – 5) are left as exercises. •

Theorem 2.9 shows us that complements, closures, interiors and frontiers are interrelated and therefore some of these operators are redundant. That is, if we wanted to very “economical,” we could discard some of them. For example, we could avoid using “Fr” and “int” and just use “cl” and complement because  $\text{Fr } A = \text{cl } A \cap \text{cl } (X - A)$  and  $\text{int } A = A - \text{Fr } A = A - (\text{cl } A \cap \text{cl } (X - A))$ . Of course, the most economical way of doing things is not necessarily the most convenient. (*Could we get by only using complements and “Fr” – that is, can we define “int” and “cl” in terms of “Fr” and complements? Or could we use just “int” and complements?*)

Here is a famous related problem from the early days of topology: for  $A \subseteq (X, \mathcal{T})$ , is there an upper bound for the number of different subsets of  $X$  which might be created from  $A$  using only complements and closures, repeated in any order? (*As we just observed, using the interior and frontier operators would not help to create any additional sets.*) For example, one might start with  $A$  and then consider such sets as  $\text{cl } A$ ,  $X - \text{cl } A$ ,  $\text{cl } (X - \text{cl } (X - A))$ , and so on. An old theorem of Kuratowski (1922) says that for any set  $A$  in any space  $(X, \mathcal{T})$ , the upper bound is 14. Moreover, this upper bound is “sharp” – there is a set  $A \subseteq \mathbb{R}$  from which 14 different sets can actually be obtained! Can you find such a set?

**Definition 2.10** Suppose  $(X, \mathcal{T})$  is a topological space and  $D \subseteq X$ .  $D$  is called dense in  $X$  if  $\text{cl } D = X$ . The space  $(X, \mathcal{T})$  is called separable if there exists a countable dense set  $D$  in  $X$ .

**Example 2.11** Let  $\mathcal{T}$  be the cofinite topology on  $\mathbb{R}$ . Let  $\mathbb{E} = \{2, 4, 6, \dots\}$ .

$\text{int } \mathbb{E} = \emptyset$ , because each nonempty open set  $O$  has finite complement and therefore  $O \not\subseteq \mathbb{E}$ . In fact, for any  $B \subseteq \mathbb{R}$  : if  $\mathbb{R} - B$  is infinite, then  $\text{int } B = \emptyset$

/  
 $\text{cl } \mathbb{E} = \mathbb{R}$  because the only closed set containing  $\mathbb{E}$  is  $\mathbb{R}$ . Therefore  $(\mathbb{R}, \mathcal{T})$  is separable.  
In fact, any infinite set  $B$  is dense.

$$\text{Fr } \mathbb{E} = \text{cl } (\mathbb{E}) \cap \text{cl } (\mathbb{R} - \mathbb{E}) = \mathbb{R} \cap \mathbb{R} = \mathbb{R}$$

$(\mathbb{R}, \mathcal{T})$  is not pseudometrizable (*why?*)

**Example 2.12** Let  $\mathcal{T}$  be the right-ray topology on  $\mathbb{R}$ . In  $(\mathbb{R}, \mathcal{T})$ ,

$$\text{int } \mathbb{Z} = \emptyset$$

$$\text{cl } \mathbb{Z} = \mathbb{R}, \text{ so } (\mathbb{R}, \mathcal{T}) \text{ is separable}$$

$$\text{Fr } \mathbb{Z} = \mathbb{R}$$

Any two nonempty open sets intersect, so  $(\mathbb{R}, \mathcal{T})$  is not metrizable. Is it pseudometrizable?

### 3. Subspaces

There is a natural way to create a new topological space from a subset of a given topological space. The new space is called a subspace (not merely a subset) of the original space.

**Definition 3.1** Suppose  $(X, \mathcal{T})$  is a topological space and that  $A \subseteq X$ . The subspace topology on  $A$  is defined as  $\mathcal{T}_A = \{A \cap O : O \in \mathcal{T}\}$  and  $(A, \mathcal{T}_A)$  is called a subspace of  $(X, \mathcal{T})$ .

We sometimes call  $O \cap A$  the “restriction of  $O$  to  $A$ ” or the “trace of  $O$  on  $A$ ”

*You should check the conditions in Definition 2.1 are satisfied: in other words, that  $\mathcal{T}_A$  really is a topology on  $A$ . When we say that  $A$  is a subspace of  $(X, \mathcal{T})$ , we mean that  $A$  is a subset of  $X$  with the subspace topology  $\mathcal{T}_A$ . To indicate that  $A$  is a subspace, we sometimes write  $A \subseteq (X, \mathcal{T})$  rather than  $A \subseteq X$ .*

**Example 3.2** Consider  $\mathbb{N} \subseteq \mathbb{R}$ , where  $\mathbb{R}$  has its usual topology. For each  $n \in \mathbb{N}$ , the interval  $(n-1, n+1)$  is open in  $\mathbb{R}$ . Therefore  $(n-1, n+1) \cap \mathbb{N} = \{n\}$  is open in the subspace  $\mathbb{N}$ , so every point  $n$  is isolated in the subspace. The subspace topology is the discrete topology. Notice: the subspace topology on  $\mathbb{N}$  is the same as what we get if we use the usual metric on  $\mathbb{N}$  to generate open sets in  $\mathbb{N}$ . Similarly, it is easy to check that in  $\mathbb{R}^2$  the subspace topology on the  $x$ -axis is the same as the usual metric topology on  $\mathbb{R}$ .

More generally: suppose  $A \subseteq X$ , where  $(X, d)$  is a pseudometric space. Then we can think of two ways to make  $A$  into a topological space.

- i)  $d$  gives a topology  $\mathcal{T}_d$  on  $X$ . Take the open sets in  $\mathcal{T}_d$  and intersect them with  $A$ . This gives us the subspace topology on  $A$ , which we call  $(\mathcal{T}_d)_A$  and  $(A, (\mathcal{T}_d)_A)$  is a subspace of  $(X, \mathcal{T})$ .
- 2) Or, we could view  $A$  as a pseudometric space by using  $d$  to measure distances in  $A$ . To be very precise,  $d : X \times X \rightarrow \mathbb{R}$ , so we make a “new” pseudometric  $d'$  defined by  $d' = d|A \times A$ . Then  $(A, d')$  is a pseudometric space and we can use  $d'$  to generate open sets in  $A$ : the topology  $\mathcal{T}_{d'}$ .

*Usually, we would be less compulsive about notation and continue to use the name “ $d$ ” also for the pseudometric on  $A$ . But for a moment it will be helpful to distinguish carefully between  $d$  and  $d' = d|A \times A$ .*

Fortunately, it turns out that 1) and 2) produce the same open sets in  $A$  : the open sets in  $(A, d')$  are just the open sets from  $(X, d)$  restricted to  $A$ . That's just what Theorem 3.3 says, in “fancier” notation.

**Theorem 3.3** Suppose  $A \subseteq X$ , where  $(X, d)$  is a pseudometric space. Then  $(\mathcal{T}_d)_A = \mathcal{T}_{d'}$ .

**Proof** If  $U \in (\mathcal{T}_d)_A$ , then  $U = O \cap A$  where  $O \in \mathcal{T}_d$ . Let  $a \in U$ . There is an  $\epsilon > 0$  such that  $B_\epsilon^d(a) \subseteq O$ . Since  $d' = d|A \times A$ , we get that  $B_\epsilon^{d'}(a) = B_\epsilon^d(a) \cap A \subseteq O \cap A = U$ , so  $U \in \mathcal{T}_{d'}$ .

Conversely, suppose  $U \in \mathcal{T}_{d'}$ . For each  $a \in U$  there is an  $\epsilon_a > 0$  such that  $B_{\epsilon_a}^{d'}(a) \subseteq U$ , and  $U = \bigcup_{a \in U} B_{\epsilon_a}^{d'}(a)$ . Let  $O = \bigcup_{a \in U} B_{\epsilon_a}^d(a)$ , an open set in  $\mathcal{T}_d$ . Since  $B_{\epsilon_a}^{d'}(a) = B_{\epsilon_a}^d(a) \cap A$ , we get  $O \cap A = (\bigcup_{a \in U} B_{\epsilon_a}^d(a)) \cap A = (\bigcup_{a \in O} B_{\epsilon_a}^d(a) \cap A) = \bigcup_{a \in U} B_{\epsilon_a}^d(a) = U$ . Therefore  $U \in (\mathcal{T}_d)_A$ . •

**Exercise** Verify that in any topological space  $(X, \mathcal{T})$

- i) If  $U$  is open in  $(X, \mathcal{T})$  and  $A$  is an open set in the subspace  $U$ , then  $A$  is open in  $X$ . (“*An open subset of an open set is open.*”)
- ii) If  $F$  is closed in  $(X, \mathcal{T})$  and  $A$  is a closed set in the subspace  $F$ , then  $F$  is closed in  $X$ . (“*A closed subset of a closed set is closed.*”)

## 4. Neighborhoods

**Definition 4.1** Let  $(X, \mathcal{T})$  be a topological space and suppose  $N \subseteq X$ . If  $x \in \text{int } N$ , then we say that  $N$  is a neighborhood of  $x$ .

The collection  $\mathcal{N}_x = \{N \subseteq X : N \text{ is a neighborhood of } x\}$  is called the neighborhood system at  $x$ .

Note that

- 1)  $\mathcal{N}_x \neq \emptyset$ , because every point  $x$  has at least one neighborhood – for example,  $X \in \mathcal{N}_x$ .
- 2) If  $N_1$  and  $N_2 \in \mathcal{N}_x$ , then  $x \in \text{int } N_1 \cap \text{int } N_2 = (\text{why?}) \text{int } (N_1 \cap N_2)$ .  
Therefore  $N_1 \cap N_2 \in \mathcal{N}_x$ .
- 3) If  $N \in \mathcal{N}_x$  and  $N \subseteq N'$ , then  $x \in \text{int } N \subseteq \text{int } N'$ , so  $N' \in \mathcal{N}_x$  (that is, if  $N'$  contains a neighborhood of  $x$ , then  $N'$  is also a neighborhood of  $x$ .)

Just as in pseudometric spaces, it is clear that a set  $O$  in  $(X, \mathcal{T})$  is open iff  $O$  is a neighborhood of each of its points. (See Theorem II.5.4)

In a pseudometric space, we use the  $\epsilon$ -balls centered at  $x$  to measure “nearness” to  $x$ . For example, if “every  $\epsilon$ -ball in  $\mathbb{R}$  centered at  $x$  contains an irrational number,” this tells us that “there are irrational numbers arbitrarily near to  $x$ .” Of course, we could convey the same information in terms of neighborhoods by saying “every neighborhood of  $x$  in  $\mathbb{R}$  contains an irrational number.” Or, instead, we could say it in terms of “open sets”: “every open set in  $\mathbb{R}$  containing  $x$  contains an irrational number.” These are all equivalent ways to say “there are irrational numbers arbitrarily near to  $x$ .” *This isn't that surprising since open sets and neighborhoods in  $(X, d)$  were defined in terms of  $\epsilon$ -balls.*

In a topological space  $(X, \mathcal{T})$  we don't have  $\epsilon$ -balls, but we still have open sets and neighborhoods. We now think of the neighborhoods in  $\mathcal{N}_x$  (or, if we prefer, the collection of open sets containing  $x$ ) as the tool we use to talk about “nearness to  $x$ .”

For example, suppose  $X$  has the trivial topology  $\mathcal{T}$ . For any  $x \in X$ , the only neighborhood of  $x$  is  $X$ : therefore every  $y$  in  $X$  is in every neighborhood of  $x$ : the neighborhoods of  $x$  are unable to “separate”  $x$  and  $y$ , and that's analogous to having  $d(x, y) = 0$  (if we had a pseudometric). In that sense, all points in  $(X, \mathcal{T})$  are “very close together”: so close together, in fact, that they are “indistinguishable.” The neighborhoods of  $x$  tell us this.

At the opposite extreme, suppose  $X$  has the discrete topology  $\mathcal{T}$  and that  $x \in X$ . If  $x \in W$  then (*since  $W$  is open*),  $W$  is a neighborhood of  $x$ . The smallest neighborhood of  $x$  is  $N = \{x\}$ . So every point  $x$  has a neighborhood  $N$  that excludes all other points  $y$ : for every  $y \neq x$ , we could

say “ $y$  is not within the neighborhood  $N$  of  $x$ .” This is analogous to saying “ $y$  is not within  $\epsilon$  of  $x$ ” (if we had a pseudometric). Because no point  $y$  is “within  $N$  of  $x$ ,” we call  $x$  isolated. The neighborhoods of  $x$  tell us this.

*Of course, if we prefer, we could use “ $U$  is an open set containing  $x$ ” instead of “ $N$  is a neighborhood of  $x$ ” to talk about nearness to  $x$ .*

The complete neighborhood system  $\mathcal{N}_x$  of a point  $x$  often contains more neighborhoods than we actually need to talk about nearness to  $x$ . For example, the open balls  $B_{\frac{1}{n}}(x)$  in a pseudometric space  $(X, d)$  are enough to let us talk about continuity at  $x$ . Therefore, we introduce the idea of a neighborhood base at  $x$  to choose a smaller collection of neighborhoods of  $x$  that is

- i) good enough for all our purposes, and
- ii) from which all the other neighborhoods of  $x$  can be obtained if we want them.

**Definition 4.2** A collection  $\mathcal{B}_x \subseteq \mathcal{N}_x$  is called a neighborhood base at  $x$  if for every neighborhood  $N$  of  $x$ , there is a neighborhood  $B \in \mathcal{B}_x$  such that  $x \in B \subseteq N$ . We refer to the sets in  $\mathcal{B}_x$  as basic neighborhoods of  $x$ .

According to the definition, each set in  $\mathcal{B}_x$  must be a neighborhood of  $x$ , but the collection  $\mathcal{B}_x$  may be much simpler than the whole neighborhood system  $\mathcal{N}_x$ . The crucial thing is that every neighborhood  $N$  of  $x$  must contain a basic neighborhood  $B$  of  $x$ .

**Example 4.3** In  $(X, d)$ , possible ways to choose a neighborhood base at  $x$  include:

- i)  $\mathcal{B}_x$  = the collection of all balls  $B_\epsilon(x)$ , or
- ii)  $\mathcal{B}_x$  = the collection of all balls  $B_\epsilon(x)$ , where  $\epsilon$  is a positive rational, or
- iii)  $\mathcal{B}_x$  = the collection of all balls  $B_{\frac{1}{n}}(x)$  for  $n \in \mathbb{N}$ , or
- iv)  $\mathcal{B}_x = \mathcal{N}_x$  (*The neighborhood system is always a base for itself, but it is not an “efficient” choice; the goal is to get a base  $\mathcal{B}_x$  that's much simpler than  $\mathcal{N}_x$ .*)

Which  $\mathcal{B}_x$  to use is our choice: each of i)-iv) gives a neighborhood base at  $x$ . But ii) or iii) might be more convenient – because ii) and iii) are countable neighborhood bases  $\mathcal{B}_x$ . (*If  $X = \mathbb{R}$ , for example, with the usual metric  $d$ , then the collections  $\mathcal{B}_x$  in i) and iv) are uncountable.*) Of the four, iii) is probably the simplest choice for  $\mathcal{B}_x$ .

Suppose we want to check whether some property that involves neighborhoods of  $x$  is true. Often all we need to do is to check whether the property holds for neighborhoods in the simpler collection  $\mathcal{B}_x$ . For example, in  $(X, d)$  :  $O$  is open iff  $O$  contains a neighborhood  $N$  of each  $x \in O$ . But that is true iff  $O$  contains a set  $B_{\frac{1}{n}}(x)$  around each of its points  $x$ .

Similarly,  $x \in \text{cl } A$  iff  $N \cap A \neq \emptyset$  for every  $N \in \mathcal{N}_x$  iff  $B \cap A \neq \emptyset$  for every  $B \in \mathcal{B}_x$ . For example, suppose we want to check, in  $\mathbb{R}$ , that  $1 \in \text{cl } \mathbb{P}$ . It is sufficient just to check that  $B_{\frac{1}{n}}(1) \cap \mathbb{P} \neq \emptyset$  for each  $n \in \mathbb{N}$ , because this implies that  $N \cap \mathbb{P} \neq \emptyset$  for every  $N \in \mathcal{N}_1$ .

Therefore it's often desirable to make an “efficient” choice of neighborhood base  $\mathcal{B}_x$  at each point  $x \in X$ .

**Definition 4.4** We say that a space  $(X, \mathcal{T})$  satisfies the first axiom of countability (or, more simply, that  $(X, \mathcal{T})$  is first countable) if at each point  $x \in X$ , it is possible to choose a countable neighborhood base  $\mathcal{B}_x$ .

**Example 4.5**

- 1) The preceding Example 4.3 shows that every pseudometric space is first countable.
- 2) If  $\mathcal{T}$  is the discrete topology on  $X$ , then  $(X, \mathcal{T})$  is first countable. In fact, at each point  $x$ , we can choose a neighborhood base that consists of a single set:  $\mathcal{B}_x = \{ \{x\} \}$ .
- 3) Let  $\mathcal{T}$  be the cofinite topology on an uncountable set  $X$ . For any  $x \in X$ , there cannot be a countable neighborhood base  $\mathcal{B}_x$  at  $x$ .

We prove this by contradiction: suppose we had a countable neighborhood base at some point  $x$ : call it  $\mathcal{B}_x = \{B_1, \dots, B_n, \dots\}$ .

For any  $y \neq x$ ,  $\{y\}$  is closed, so  $X - \{y\}$  is a neighborhood of  $x$ . Then, by the definition of neighborhood base, there is some  $k$  for which  $x \in \text{int } B_k \subseteq B_k \subseteq X - \{y\}$ . Therefore  $y \notin \bigcap_{n=1}^{\infty} \text{int } B_n$ .

But  $x \in \bigcap_{n=1}^{\infty} \text{int } B_n$  so  $\bigcap_{n=1}^{\infty} \text{int } B_n = \{x\}$ .

Then  $X - \{x\} = X - \bigcap_{n=1}^{\infty} \text{int } B_n = \bigcup_{n=1}^{\infty} (X - \text{int } B_n)$ . Since  $X - \text{int } B_n$  is finite (*why?*), this would mean that  $X - \{x\}$  is countable – which is impossible.

Since any pseudometric space is first countable, the example gives us another way to see that this space  $(X, \mathcal{T})$  is not pseudometrizable.

In  $(X, \mathcal{T})$ , the neighborhood system  $\mathcal{N}_x$  at each point  $x$  is completely determined by the topology  $\mathcal{T}$ , but  $\mathcal{B}_x$  is not. As the preceding examples illustrate, there are usually many possible choices for  $\mathcal{B}_x$ . (*Can you describe all the spaces  $(X, \mathcal{T})$  for which  $\mathcal{B}_x$  is uniquely determined at each point  $x$ ?*)

On the other hand, if we were given  $\mathcal{B}_x$  at each point  $x \in X$  we could

- 1) “reconstruct” the whole neighborhood system  $\mathcal{N}_x$ :

$$\mathcal{N}_x = \{N \subseteq X : \exists B \in \mathcal{B}_x \text{ such that } x \in B \subseteq N\}, \text{ and then we could}$$

- 2) “reconstruct” the whole topology  $\mathcal{T}$ :

$$\begin{aligned} \mathcal{T} &= \{O : O \text{ is a neighborhood of each of its points}\} \\ &= \{O : \forall x \in O \exists B \in \mathcal{B}_x x \in B \subseteq O\}, \text{ that is,} \\ &O \text{ is open iff } O \text{ contains a basic neighborhood of each of its points.} \end{aligned}$$

This illustrates one method of describing a topology: by telling the neighborhood basis  $\mathcal{B}_x$  at each point. Various effective methods to describe topologies are discussed in the next section.

## 5. Describing Topologies

How can a topological space be described? If  $X = \{0, 1\}$ , it is simple to give a topology by just writing  $\mathcal{T} = \{\emptyset, \{0\}, X\}$ . However, describing all the sets in  $\mathcal{T}$  explicitly is often not the easiest way to go.

In this section we look at three important – different but closely related – ways to define a topology on a set. All of them will be used throughout the course. A fourth method – by using a “closure operator” – is not used much nowadays. It is included just as an historical curiosity.

### A. Basic neighborhoods at each point

Suppose that at each point  $x \in (X, \mathcal{T})$  we have picked a neighborhood base  $\mathcal{B}_x$ . As mentioned above, the collections  $\mathcal{B}_x$  implicitly contain all the information about the topology: a set  $O$  is in  $\mathcal{T}$  iff  $O$  contains a basic neighborhood of each of its points. This suggests that if we start with just a set  $X$ , then we could define a topology on  $X$  if we begin by saying what the  $\mathcal{B}_x$ 's should be. Of course, we can't just put “random” sets in  $\mathcal{B}_x$ : the sets in each  $\mathcal{B}_x$  must “act the way basic neighborhoods are supposed to act.” And how is that? The next theorem describes the crucial behavior of a collection of basic neighborhoods at  $x$  in any topological space.

**Theorem 5.1** Suppose  $(X, \mathcal{T})$  is a topological space and that for each point  $x \in X$ ,  $\mathcal{B}_x$  is a neighborhood base.

- 1)  $\mathcal{B}_x \neq \emptyset$  and  $B \in \mathcal{B}_x \Rightarrow x \in B \subseteq X$
- 2) if  $B_1$  and  $B_2 \in \mathcal{B}_x$ , then  $\exists B_3 \in \mathcal{B}_x$  such that  $x \in B_3 \subseteq B_1 \cap B_2$
- 3) if  $B \in \mathcal{B}_x$ , then  $\exists I$  such that  $x \in I \subseteq B$  and,  
 $\forall y \in I, \exists B_y \in \mathcal{B}_y$  such that  $y \in B_y \subseteq I$
- 4)  $O \in \mathcal{T} \Leftrightarrow \forall x \in O \ \exists B \in \mathcal{B}_x$  such that  $x \in B \subseteq O$ .

**Proof** 1) Since  $X$  is a neighborhood of  $x$ , there is a  $B \in \mathcal{B}_x$  such that  $x \in B \subseteq X$ . Therefore  $\mathcal{B}_x \neq \emptyset$ . If  $B \in \mathcal{B}_x \subseteq \mathcal{N}_x$ , then  $B$  is a neighborhood of  $x$ , so  $x \in \text{int } B \subseteq B \subseteq X$ .

2) The intersection of the two neighborhoods  $B_1$  and  $B_2$  of  $x$  is a neighborhood of  $x$ . Therefore, by the definition of a neighborhood base, there is a set  $B_3 \in \mathcal{B}_x$  such that  $x \in B_3 \subseteq B_1 \cap B_2$ .

3) Let  $I = \text{int } B$ . Then  $x \in I \subseteq B$  and because  $I$  is open,  $I$  is a neighborhood of each of its points  $y$ . Since  $\mathcal{B}_y$  is a neighborhood base at  $y$ , there is a set  $B_y \in \mathcal{B}_y$  such that  $y \in B_y \subseteq I$ .

4)  $\Rightarrow$  : If  $O$  is open, then  $O$  is a neighborhood of each of its points  $x$ . Therefore for each  $x \in O$  there must be a set  $B \in \mathcal{B}_x$  such that  $x \in B \subseteq O$ .

$\Leftarrow$  : The condition implies that  $O$  contains a neighborhood of each of its points. Therefore  $O$  is a neighborhood of each of its points, so  $O$  is open. •

Theorem 5.1 lists the crucial features of the behavior of a neighborhood base at  $x$ . The next theorem tells us that we can put a topology on a set  $X$  by assigning a “properly behaved” collection of sets to become the basic neighborhoods at each point  $x$ .

**Theorem 5.2 (The Neighborhood Base Theorem)** Let  $X$  be a set. Suppose that for each  $x \in X$  we give a collection  $\mathcal{B}_x$  of subsets of  $X$  in such a way that conditions 1) - 3) of Theorem 5.1 are true. Define  $\mathcal{T} = \{O \subseteq X : \forall x \in O \ \exists B \in \mathcal{B}_x \text{ such that } x \in B \subseteq O\}$ . Then  $\mathcal{T}$  is a topology on  $X$  and  $\mathcal{B}_x$  is now a neighborhood base at  $x$  in  $(X, \mathcal{T})$ .

*Note: In Theorem 5.2, we do not ask that the  $\mathcal{B}_x$ 's satisfy condition 4) of Theorem 5.1 – since  $X$  is a set with no topology (yet), condition 4) would be meaningless. Rather, Condition 4) becomes the motivation for how to define a topology  $\mathcal{T}$  using the  $\mathcal{B}_x$ 's.*

**Proof** We need to prove three things: a)  $\mathcal{T}$  is a topology

- b) each  $B \in \mathcal{B}_x$  is now a neighborhood of  $x$  in  $(X, \mathcal{T})$ , and
- c) the collection  $\mathcal{B}_x$  is now a neighborhood base at  $x$ .

a) Clearly,  $\emptyset \in \mathcal{T}$ . If  $x \in X$  then, by condition 1), we can choose any  $B \in \mathcal{B}_x$  and  $x \in B \subseteq X$ . Therefore  $X \in \mathcal{T}$ .

Suppose  $O_\alpha \in \mathcal{T}$  for all  $\alpha \in A$ . If  $x \in \bigcup\{O_\alpha : \alpha \in A\}$ , then  $x \in O_{\alpha_0}$  for some  $\alpha_0 \in A$ . By definition of  $\mathcal{T}$ , there is a set  $B \in \mathcal{B}_x$  such that  $x \in B \subseteq O_{\alpha_0} \subseteq \bigcup\{O_\alpha : \alpha \in A\}$ , so  $\bigcup\{O_\alpha : \alpha \in A\} \in \mathcal{T}$ .

To finish a), it is sufficient to show that if  $O_1$  and  $O_2 \in \mathcal{T}$ , then  $O_1 \cap O_2 \in \mathcal{T}$ . Suppose  $x \in O_1 \cap O_2$ . By the definition of  $\mathcal{T}$ , there are sets  $B_1$  and  $B_2 \in \mathcal{B}_x$  such that  $x \in B_1 \subseteq O_1$  and  $x \in B_2 \subseteq O_2$ , so  $x \in B_1 \cap B_2 \subseteq O_1 \cap O_2$ . By condition 2), there is a set  $B_3 \in \mathcal{B}_x$  such that  $x \in B_3 \subseteq B_1 \cap B_2 \subseteq O_1 \cap O_2$ . Therefore  $O_1 \cap O_2 \in \mathcal{T}$ .

Therefore  $\mathcal{T}$  is a topology on  $X$  – so now we have a topological space  $(X, \mathcal{T})$  – and we must show that  $\mathcal{B}_x$  is a neighborhood base at  $x$  in  $(X, \mathcal{T})$ . Doing so involves the awkward-looking condition 3) – which we have not yet used.

b) If  $B \in \mathcal{B}_x$ , then  $x \in B$  (by condition 1) and (by condition 3), there is a set  $I \subseteq X$  such that  $x \in I \subseteq B$  and  $\forall y \in I, \exists B_y \in \mathcal{B}_y \text{ such that } y \in B_y \subseteq I$ . The underlined phrase states that  $I$  satisfies the condition for  $I \in \mathcal{T}$ , so  $I$  is open. Since  $I$  is open and  $x \in I \subseteq B$ ,  $B$  is a neighborhood of  $x$ , that is,  $\mathcal{B}_x \subseteq \mathcal{N}_x$ .

c) To complete the proof, we have to check that  $\mathcal{B}_x$  is a neighborhood base at  $x$ . If  $N$  is a neighborhood of  $x$ , then  $x \in \text{int } N$ . Since  $\text{int } N$  is open,  $\text{int } N$  must satisfy the criterion for membership in  $\mathcal{T}$ , so there is a set  $B \in \mathcal{B}_x$  such that  $x \in B \subseteq \text{int } N \subseteq N$ . Therefore  $\mathcal{B}_x$  forms a neighborhood base at  $x$ . •

**Example 5.3** For each  $x \in \mathbb{R}$ , let  $\mathcal{B}_x = \{[x, b) : b > x\}$ . We can easily check the conditions 1) - 3) from Theorem 5.2:

- 1) For each  $x \in \mathbb{R}$ , certainly  $\mathcal{B}_x \neq \emptyset$  and  $x \in [x, b)$  for each set  $[x, b) \in \mathcal{B}_x$
- 2) If  $B_1 = [x, b_1)$  and  $B_2 = [x, b_2)$  are in  $\mathcal{B}_x$ , then (in this example) we can choose  $B_3 = B_1 \cap B_2 = [x, b_3) \in \mathcal{B}_x$ , where  $b_3 = \min\{b_1, b_2\}$ .
- 3) If  $B = [x, b) \in \mathcal{B}_x$ , then (in this example) we can let  $I = B$ .  
If  $y \in I = [x, b)$ , pick  $c$  so  $y < c < b$ . Then  $B_y = [y, c) \in \mathcal{B}_y$  and  $y \in [y, c) \subseteq I$ .

According to Theorem 5.2,  $\mathcal{T} = \{O \subseteq \mathbb{R} : \forall x \in O \exists [x, b) \in \mathcal{B}_x \text{ such that } x \in [x, b) \subseteq O\}$  is a topology on  $\mathbb{R}$  and  $\mathcal{B}_x$  is a neighborhood base at  $x$  in  $(\mathbb{R}, \mathcal{T})$ . The space  $(\mathbb{R}, \mathcal{T})$  is called the Sorgenfrey line.

Notice that in this example each set  $[x, b) \in \mathcal{T}$ : that is, the sets in  $\mathcal{B}_x$  turn out to be open, not merely neighborhoods of  $x$ . (*This does not always happen.*)

It is easy to check that sets of form  $(-\infty, x)$  and  $[b, \infty)$  are open, so  $(-\infty, x) \cup [b, \infty) = \mathbb{R} - [x, b)$  is open. Therefore  $[x, b)$  is also closed. So at each point  $x$  in the Sorgenfrey line, there is a neighborhood base  $\mathcal{B}_x$  consisting of clopen sets.

We can write  $(a, c) = \bigcup_{n=1}^{\infty} [a + \frac{1}{n}, c)$ , so  $(a, c)$  is open in the Sorgenfrey line. Because every usual open set in  $\mathbb{R}$  is a union of sets of the form  $(a, c)$ , we conclude that every usual open set in  $\mathbb{R}$  is also open in the Sorgenfrey line. The usual topology on  $\mathbb{R}$  is strictly smaller than the Sorgenfrey topology:  $\mathcal{T}_d \neq \mathcal{T}$ .

$\mathbb{Q}$  is dense in the Sorgenfrey line: if  $x \in \mathbb{R}$ , then every basic neighborhood  $[x, b)$  of  $x$  intersects  $\mathbb{Q}$ , so  $x \in \text{cl } \mathbb{Q}$ . Therefore the Sorgenfrey line is separable. It is also clear that the Sorgenfrey line is first countable: at each point  $x$  the collection  $\{[x, x + \frac{1}{n}) : n \in \mathbb{N}\}$  is a countable neighborhood base.

**Example 5.4** Similarly, we can define the Sorgenfrey plane by putting a new topology on  $\mathbb{R}^2$ . At each point  $(x, y) \in \mathbb{R}^2$ , let  $\mathcal{B}_{(x,y)} = \{[x, b) \times [y, c) : b > x, c > y\}$ . The families  $\mathcal{B}_{(x,y)}$  satisfy the conditions in the Neighborhood Base Theorem, so they give a topology  $\mathcal{T}$  for which  $\mathcal{B}_{(x,y)}$  is a neighborhood base at  $(x, y)$ . A set  $O \subseteq \mathbb{R}^2$  is open iff: for each  $(x, y) \in O$ , there are  $b > x$  and  $c > y$  such that  $(x, y) \in [x, b) \times [y, c) \subseteq O$ . (*Make a sketch!*) You should check that the sets  $[x, b) \times [y, c) \in \mathcal{B}_{(x,y)}$  are actually clopen in the Sorgenfrey plane. It is also easy to check the usual topology  $\mathcal{T}_d$  on the plane is strictly smaller than the Sorgenfrey topology. It is clear that  $\mathbb{Q}^2$  is dense, so  $(\mathbb{R}^2, \mathcal{T})$  is separable. Is the Sorgenfrey plane first countable?

**Example 5.5** At each point  $p \in \mathbb{R}^2$ , let  $C_\epsilon(p) = \{x \in \mathbb{R}^2 : d(x, p) \leq \epsilon\}$  and define  $\mathcal{B}_p = \{C_\epsilon(p) : p \in \mathbb{R}^2\}$ . It is easy to check that the conditions 1)-3) of Theorem 5.2 are satisfied. (*For  $C_\epsilon(p)$  in condition 3), let  $I = B_\epsilon(p)$ .*) The topology generated by the  $\mathcal{B}_p$ 's is just the usual topology; the sets in  $\mathcal{B}_p$  are basic neighborhoods of  $p$  in the usual topology – as they should be – but the sets in  $\mathcal{B}_p$  did not turn out to be open sets.

**Example 5.6** Let  $\Gamma = \{(x, y) \in \mathbb{R}^2 : y \geq 0\}$  = the “closed upper half-plane.”

For a point  $p = (x, y) \in \Gamma$  with  $y > 0$ , let  $\mathcal{B}_p = \{B_\epsilon(p) : \epsilon < |y|\}$

For a point  $p = (x, 0) \in \Gamma$ , let

$\mathcal{B}_p = \{p\} \cup A : A \text{ is a usual open disc in the upper half-plane, tangent to the } x\text{-axis at } p\}.$

It is easy to check that the collections  $\mathcal{B}_p$  satisfy the conditions 1)-3) of Theorem 5.2 and therefore give a topology on  $\Gamma$ . In this topology, the sets in  $\mathcal{B}_p$  turn out to be open neighborhoods of  $p$ .

The space  $\Gamma$ , with this topology, is called the “Moore plane.” Notice that  $\Gamma$  is separable and first countable. The subspace topology on the  $x$ -axis is the discrete topology. (*Verify these statements!*)

## B. Base for the topology

**Definition 5.7** A collection of open sets in  $(X, \mathcal{T})$  is called a base for the topology  $\mathcal{T}$  if each  $O \in \mathcal{T}$  is a union of sets from  $\mathcal{B}$ . More precisely,  $\mathcal{B}$  is a base if  $\mathcal{B} \subseteq \mathcal{T}$  and for each  $O \in \mathcal{T}$ , there exists a subfamily  $\mathcal{A} \subseteq \mathcal{B}$  such that  $O = \bigcup \mathcal{A}$ . We also call  $\mathcal{B}$  a base for the open sets and we refer to the open sets in  $\mathcal{B}$  as basic open sets.

If  $\mathcal{B}$  is a base, then it is easy to see that:  $O \in \mathcal{T}$  iff  $\forall x \in O \exists B \in \mathcal{B}$  such that  $x \in B \subseteq O$ . This means that if we were given  $\mathcal{B}$ , we could use it to decide which sets are open and thus “reconstruct”  $\mathcal{T}$ .

Of course, one example of a base is  $\mathcal{B} = \mathcal{T}$ : every topology  $\mathcal{T}$  is a base for itself. But usually there are many ways to choose a base, and the idea is that a simpler base  $\mathcal{B}$  would be easier to work with. For example, the set of all balls is a base for the topology  $\mathcal{T}_d$  in any pseudometric space  $(X, d)$ ; a different base would be the set containing only the balls with positive rational radii. (*Can you describe those topological spaces for which  $\mathcal{T}$  is the only base for  $\mathcal{T}$ ?*)

The following theorem tells us the crucial properties of a base  $\mathcal{B}$  in  $(X, \mathcal{T})$ .

**Theorem 5.8** If  $(X, \mathcal{T})$  is a topological space with a base  $\mathcal{B}$  for  $\mathcal{T}$ , then

$$1) X = \bigcup \{B : B \in \mathcal{B}\}$$

$$2) \text{ if } B_1 \text{ and } B_2 \in \mathcal{B} \text{ and } x \in B_1 \cap B_2, \text{ then there is a set } B_3 \in \mathcal{B} \text{ such that } x \in B_3 \subseteq B_1 \cap B_2.$$

**Proof** 1) Certainly  $\bigcup \mathcal{B} \subseteq X$  and – since  $X$  is open – the definition of base implies that  $X$  is the union of a subfamily of  $\mathcal{B}$ . Therefore  $\bigcup \mathcal{B} = X$ .

2) If  $B_1$  and  $B_2 \in \mathcal{B}$ , then  $B_1 \cap B_2$  is open so  $B_1 \cap B_2$  must be the union of some sets from  $\mathcal{B}$ . Therefore, if  $x \in B_1 \cap B_2$ , there must be a set  $B_3 \in \mathcal{B}$  such that  $x \in B_3 \subseteq B_1 \cap B_2$ . •

The next theorem tells us that if we are given a collection  $\mathcal{B}$  of subsets of a set  $X$  with properties 1) and 2), we can use it to define a topology.

**Theorem 5.9 (The Base Theorem)** Suppose  $X$  is a set and that  $\mathcal{B}$  is a collection of subsets of  $X$  that satisfies conditions 1) and 2) in Theorem 5.8. Define  $\mathcal{T} = \{O \subseteq X : O \text{ is a union of sets from } \mathcal{B}\} = \{O \subseteq X : \forall x \in O \exists B \in \mathcal{B} \text{ such that } x \in B \subseteq O\}$ . Then  $\mathcal{T}$  is a topology on  $X$  and  $\mathcal{B}$  is a base for  $\mathcal{T}$ .

**Proof** First we show that  $\mathcal{T}$  is a topology on  $X$ . Since  $\emptyset$  is the union of the empty subfamily of  $\mathcal{B}$ , we get that  $\emptyset \in \mathcal{T}$ , and condition 1) simply states that  $X \in \mathcal{T}$ .

If  $O_\alpha \in \mathcal{T}$  ( $\alpha \in A$ ), then  $\bigcup\{O_\alpha : \alpha \in A\}$  is a union of sets  $O_\alpha$  each of which is itself a union of sets from  $\mathcal{B}$ . Then clearly  $\bigcup\{O_\alpha : \alpha \in A\}$  is a union of sets from  $\mathcal{B}$ , so  $\bigcup\{O_\alpha : \alpha \in A\} \in \mathcal{T}$ .

Suppose  $O_1$  and  $O_2 \in \mathcal{T}$  and that  $x \in O_1 \cap O_2$ . For each such  $x$  we can use 2) to pick a set  $B_x \in \mathcal{B}$  such that  $x \in B_x \subseteq O_1 \cap O_2$ . Then  $O_1 \cap O_2$  is the union of all the  $B_x$ 's chosen in this way, so  $O_1 \cap O_2 \in \mathcal{T}$ .

Now we know that we have a topology,  $\mathcal{T}$ , on  $X$ . By definition of  $\mathcal{T}$  it is clear that  $\mathcal{B} \subseteq \mathcal{T}$  and that each set in  $\mathcal{T}$  is a union of sets from  $\mathcal{B}$ . Therefore  $\mathcal{B}$  is a base for  $\mathcal{T}$ . •

**Example 5.10** The collections

$$\begin{aligned}\mathcal{B} &= \{B_p(\epsilon) : p \in \mathbb{R}^2, \epsilon > 0\} \\ \mathcal{B}' &= \{B_p\left(\frac{1}{k}\right) : p \in \mathbb{Q}^2, k \in \mathbb{N}\} \text{ and} \\ \mathcal{B}'' &= \{(a, b) \times (c, d) : a, b, c, d \in \mathbb{R}, a < b, c < d\}\end{aligned}$$

each satisfy the conditions 1)-2) in Theorem 5.9, so each collection is the base for a topology on  $\mathbb{R}^2$ . In fact, all three are bases for the same topology on  $\mathbb{R}^2$  – that is, the usual topology (*check this!*) Of the three,  $\mathcal{B}'$  is the simplest choice – it is a countable base for the usual topology.

**Example 5.11** Suppose  $(X, \mathcal{T}')$  and  $(Y, \mathcal{T}'')$  are topological spaces. Let  $\mathcal{B}$  = the set of “open boxes” in  $X \times Y = \{U \times V : U \in \mathcal{T}' \text{ and } V \in \mathcal{T}''\}$ . (*Verify that  $\mathcal{B}$  satisfies conditions 1) and 2) of The Base Theorem.*) The product topology on the set  $X \times Y$  is the topology for which  $\mathcal{B}$  is a base. We always assume that  $X \times Y$  has the product topology unless something else is stated.

Therefore a set  $A \subseteq X \times Y$  is open (in the product topology) iff : for all  $(x, y) \in A$ , there are open sets  $U \subseteq X$  and  $V \subseteq Y$  such that  $(x, y) \in U \times V \subseteq A$ . (*Note that  $A$  itself might not be a “box.”*)

Let  $\pi_1 : X \times Y \rightarrow X$  be the “projection” defined by  $\pi_1(x, y) = x$ . If  $U$  is any open set in  $X$ , then  $\pi_1^{-1}[U] = U \times Y \in \mathcal{B}$ . Therefore  $\pi_1^{-1}[U]$  is open. Similarly, for  $\pi_2 : X \times Y \rightarrow Y$  defined by  $\pi_2(x, y) = y$  : if  $V$  is open in  $Y$ , then  $\pi_2^{-1}[V] = X \times V$  is open in  $X \times Y$ . (*As we see in Section 8, this means that the projection maps are continuous. It is not hard to show that*

the projection maps  $\pi_1$  and  $\pi_2$  are also open maps: that is, the image of open sets in the product is open).

If  $D_1$  is dense in  $X$  and  $D_2$  is dense in  $Y$ , we claim that  $D_1 \times D_2$  is dense in  $X \times Y$ . If  $(x, y) \in A$ , where  $A$  is open, then there are nonempty open sets  $U \subseteq X$  and  $V \subseteq Y$  for which  $(x, y) \in U \times V \subseteq A$ . Since  $x \in \text{cl } D_1$ , we know that  $U \cap D_1 \neq \emptyset$ ; and similarly  $V \cap D_2 \neq \emptyset$ . Therefore  $(U \times V) \cap (D_1 \times D_2) = (U \cap D_1) \times (V \cap D_2) \neq \emptyset$ , so  $A \cap (D_1 \times D_2) \neq \emptyset$ . Therefore  $(x, y) \in \text{cl}(D_1 \times D_2)$ . So  $D_1 \times D_2$  is dense in  $X \times Y$ . In particular, this shows that the product of two separable spaces is separable.

**Example 5.12** The open intervals  $(a, b)$  form a base for the usual topology in  $\mathbb{R}$ , so each set  $(a, b) \times (c, d)$  is in the base  $\mathcal{B}$  for the product topology on  $\mathbb{R} \times \mathbb{R}$ . It is easy to see that every “open box”  $U \times V$  in  $\mathcal{B}$  can be written as a union of “simple open boxes” like  $(a, b) \times (c, d)$ . Therefore  $\mathcal{B}' = \{(a, b) \times (c, d) : a < b, c < d\}$  also is a (simpler) base for the product topology on  $\mathbb{R} \times \mathbb{R}$ . From this, it is clear that the product topology on  $\mathbb{R} \times \mathbb{R}$  is the usual topology on the plane  $\mathbb{R}^2$  (see Example 5.10).

*In general, the open sets  $U$  and  $V$  in the base for the product topology on  $X \times Y$  can be replaced by sets “ $U$  chosen from a base for  $X$ ” and “ $V$  chosen from a base for  $Y$ ,” as in this example. So in the definition of the product topology, it is sufficient to say that basic open sets are of the form  $U \times V$ , where  $U$  and  $V$  are basic open sets from  $X$  and from  $Y$ .*

**Definition 5.13** We say that a space  $(X, \mathcal{T})$  satisfies the second axiom of countability (or, more simply, that  $(X, \mathcal{T})$  is a second countable space) if it is possible to find a countable base  $\mathcal{B}$  for the topology  $\mathcal{T}$ .

For example,  $\mathbb{R}$  is second countable because, for example,  $\mathcal{B} = \{(a, b) : a, b \in \mathbb{Q}\}$  is a countable base. Is  $\mathbb{R}^2$  is second countable (*why or why not*)?

**Example 5.14** The collection  $\mathcal{B} = \{[x, x + \frac{1}{n}] : x \in \mathbb{R}, n \in \mathbb{N}\}$  is a base for the Sorgenfrey topology on  $\mathbb{R}$ . But the collection  $\{[x, x + \frac{1}{n}] : x \in \mathbb{Q}, n \in \mathbb{N}\}$  is not a countable base for the Sorgenfrey topology. Why not?

Since the sets in a base may be simpler than arbitrary open sets, they are often more convenient to work with, and working with the basic open sets is often all that is necessary – not a surprise since all the information about the open sets is contained in the base  $\mathcal{B}$ . For example, you should check that

- 1) If  $\mathcal{B}$  is a base for  $\mathcal{T}$ , then  $x \in \text{cl } A$  iff each basic open set  $B$  containing  $x$  satisfies  $B \cap A \neq \emptyset$ .
- 2) If  $f : (X, d) \rightarrow (Y, d')$  and  $\mathcal{B}$  is a base for the topology  $\mathcal{T}_{d'}$  on  $Y$ , then  $f$  is continuous iff  $f^{-1}[B]$  is open for each  $B \in \mathcal{B}$ . This means that we needn't check the inverse images of all open sets to verify that  $f$  is continuous.

## C. Subbase for the topology

**Definition 5.15** Suppose  $(X, \mathcal{T})$  is a topological space. A family  $\mathfrak{S}$  of open sets is called a subbase for the topology  $\mathcal{T}$  if the collection  $\mathcal{B}$  of all finite intersections of sets from  $\mathfrak{S}$  is a base for  $\mathcal{T}$ . (Clearly, if  $\mathcal{B}$  is a base for  $\mathcal{T}$ , then  $\mathcal{B}$  is automatically a subbase for  $\mathcal{T}$ .)

**Examples** i) The collection  $\mathfrak{S} = \{I : I = (-\infty, b) \text{ or } I = (a, \infty), a, b \in \mathbb{R}\}$  is a subbase for a topology on  $\mathbb{R}$ . All intervals of the form  $(a, b) = (-\infty, b) \cap (a, \infty)$  are in  $\mathcal{B}$ , so  $\mathcal{B}$  is a base for the usual topology on  $\mathbb{R}$ .

ii) The collection  $\mathfrak{S}$  of all sets  $U \times Y$  and  $V \times X$  (for  $U$  open in  $X$  and  $V$  open in  $Y$ ) is a subbase for the product topology on  $X \times Y$ : these sets are open in  $X \times Y$  and the collection  $\mathcal{B}$  of all finite intersections of sets in  $\mathfrak{S}$  includes all the open boxes  
 $U \times V = (U \times Y) \cap (X \times V)$ .

We can define a topology on a set  $X$  by giving a collection  $\mathfrak{S}$  of subsets as the subbase for a topology. Surprisingly, any collection  $\mathfrak{S}$  can be used: no special conditions on  $\mathfrak{S}$  are required.

**Theorem 5.16 (The Subbase Theorem)** Suppose  $X$  is a set and  $\mathfrak{S}$  is any collection of subsets of  $X$ . Let  $\mathcal{B}$  be the collection of all finite intersections of sets from  $\mathfrak{S}$ . Then  $\mathcal{B}$  is a base for a topology  $\mathcal{T}$ , and  $\mathfrak{S}$  is a subbase for  $\mathcal{T}$ .

**Proof** First we show that  $\mathcal{B}$  satisfies conditions 1) and 2) of The Base Theorem.

1)  $X \in \mathcal{B}$  since  $X$  is the intersection of the empty subcollection of  $\mathfrak{S}$  (this follows the convention that the intersection of an empty family of subsets of  $X$  is  $X$  itself. See Example I.4.5.5). Since  $X \in \mathcal{B}$ , certainly  $X = \bigcup\{B : B \in \mathcal{B}\}$ .

2) Suppose  $B_1$  and  $B_2 \in \mathcal{B}$ , and  $x \in B_1 \cap B_2$ . We know that  $B_1 = S_1 \cap \dots \cap S_m$  and  $B_2 = S_{m+1} \cap \dots \cap S_{m+k}$  for some  $S_1, \dots, S_m, \dots, S_{m+k} \in \mathfrak{S}$ , so

$$x \in B_3 = B_1 \cap B_2 = S_1 \cap \dots \cap S_m \cap S_{m+1} \cap \dots \cap S_{m+k} \in \mathcal{B}$$

Therefore  $\mathcal{T} = \{O : O \text{ is a union of sets from } \mathcal{B}\}$  is a topology, and  $\mathcal{B}$  is a base for  $\mathcal{T}$ . By definition of  $\mathcal{B}$  and  $\mathcal{T}$ , we have  $\mathfrak{S} \subseteq \mathcal{B} \subseteq \mathcal{T}$  and each set in  $\mathcal{T}$  is a union of finite intersections of sets from  $\mathfrak{S}$ . Therefore  $\mathfrak{S}$  is a subbase for  $\mathcal{T}$ . •

### Example 5.17

1) Let  $\mathbb{E} = \{2, 4, 6, \dots\} \subseteq \mathbb{N}$  and  $\mathfrak{S} = \{\{1\}, \{2\}, \mathbb{E}\}$ .  $\mathfrak{S}$  is a subbase for a topology on  $\mathbb{N}$ . A base for this topology is the collection  $\mathcal{B}$  of all finite intersections of sets in  $\mathfrak{S}$ :

$$\mathcal{B} = \{\emptyset, \mathbb{N}, \{2\}, \{1\}, \mathbb{E}\}.$$

The not-very-interesting topology  $\mathcal{T}$  generated by the base  $\mathcal{B}$  is collection of all possible unions of sets from  $\mathcal{B}$ :

$$\mathcal{T} = \{\emptyset, \mathbb{N}, \{1\}, \{2\}, \{1, 2\}, \mathbb{E}, \mathbb{E} \cup \{1\}\}$$

2) For each  $n \in \mathbb{N}$ , let  $S_n = \{n, n+1, n+2, \dots\}$ . The collection  $\mathfrak{S} = \{S_n : n \in \mathbb{N}\}$  is a subbase for a topology on  $\mathbb{N}$ . Here,  $S_n \cap S_m = S_k$  where  $k = \max\{m, n\}$ , so the collection of all finite intersections from  $\mathfrak{S}$  is just  $\mathfrak{S}$  itself. So  $\mathcal{B} = \mathfrak{S}$  is actually a base for a topology. The topology is  $\mathcal{T} = \mathfrak{S} \cup \{\emptyset\}$ .

3) Let  $\mathfrak{S} = \{\ell : \ell \text{ is a straight line in } \mathbb{R}^2\}$  in  $\mathbb{R}^2$ . For every point  $p \in \mathbb{R}^2$ ,  $\{p\}$  is the intersection of two sets from  $\mathfrak{S}$ , so  $\{p\} \in \mathcal{B}$ .  $\mathfrak{S}$  is a subbase for the discrete topology on  $\mathbb{R}^2$ .

4) Let  $\mathfrak{S} = \{\ell : \ell \text{ is a vertical line in } \mathbb{R}^2\}$ .  $\mathfrak{S}$  generates a topology on  $\mathbb{R}^2$  for which  $\mathcal{B} = \{\mathbb{R}^2, \emptyset\} \cup \mathfrak{S}$  is a base and  $\mathcal{T} = \{O : O \text{ is a union of vertical lines}\}$ .

5) Let  $\mathfrak{S}$  be a collection of subsets of  $X$  and suppose that  $\mathcal{T}'$  is any topology on  $X$  for which  $\mathfrak{S} \subseteq \mathcal{T}'$ . Since  $\mathcal{T}'$  is a topology, it must contain all finite intersections of sets in  $\mathfrak{S}$ , and therefore must contain all possible unions of such intersections. Therefore  $\mathcal{T}'$  contains the topology  $\mathcal{T}$  for which  $\mathfrak{S}$  is a subbase. To put it another way, the topology for which  $\mathfrak{S}$  is a subbase is in the smallest topology on  $X$  containing the collection  $\mathfrak{S}$ . In fact, as an exercise, you can check that

$$\mathcal{T} = \bigcap \{\mathcal{T}' : \mathcal{T}' \supseteq \mathfrak{S} \text{ and } \mathcal{T}' \text{ is a topology on } X\}.$$

**Caution** We said earlier that for some purposes it is sufficient (and simpler) to work with basic open sets, rather than arbitrary open sets – for example to check whether  $x \in \text{cl } A$ , it is sufficient to check whether  $U \cap A \neq \emptyset$  for every basic open set  $U$  that contains  $x$ . However, it is not always sufficient to work with subbasic open sets. Some caution is necessary.

For example,  $\mathfrak{S} = \{I : I = (-\infty, b) \text{ or } I = (a, \infty), a, b \in \mathbb{R}\}$  is a subbase for the usual topology on  $\mathbb{R}$ . We have  $I \cap \mathbb{Z} \neq \emptyset$  for every subbasic open set  $I$  containing  $\frac{1}{2}$ , but  $\frac{1}{2} \notin \text{cl } \mathbb{Z}$ .

## D. The closure operator

Usually we describe a topology  $\mathcal{T}$  by giving a subbase  $\mathfrak{S}$ , a base  $\mathcal{B}$ , or by giving collections  $\mathcal{B}_x$  to be the basic neighborhoods at each point  $x$ . In the early history of general topology, one other method was sometimes used. We will never actually use it, but we include it here as a curiosity.

Let  $\text{cl}_{\mathcal{T}}$  be the closure operator in  $(X, \mathcal{T})$  (*normally, we would just write “cl” for the closure operator; here we write “ $\text{cl}_{\mathcal{T}}$ ” to emphasize that this closure operator comes from the topology  $\mathcal{T}$  on  $X$* ). It gives us all the information about  $\mathcal{T}$ . That is, using  $\text{cl}_{\mathcal{T}}$ , we can decide whether any set  $A$  is closed (by asking “is  $\text{cl}_{\mathcal{T}} A = A$ ?”) and therefore can decide whether any set  $B$  is open (by asking “is  $X - B$  closed?”). It should be not be a surprise, then, that we can define a topology on a set  $X$  if we are start with an “operator” which “behaves like a closure operator.” How is that? Our first theorem tells us the crucial properties of a closure operator.

**Theorem 5.18** Suppose  $(X, \mathcal{T})$  is a topological space and  $A, B$  are subsets of  $X$ . Then

- 1)  $\text{cl}_{\mathcal{T}} \emptyset = \emptyset$
- 2)  $A \subseteq \text{cl}_{\mathcal{T}} A$
- 3)  $A$  is closed iff  $A = \text{cl}_{\mathcal{T}} A$
- 4)  $\text{cl}_{\mathcal{T}} A = \text{cl}_{\mathcal{T}}(\text{cl}_{\mathcal{T}} A)$
- 5)  $\text{cl}_{\mathcal{T}}(A \cup B) = \text{cl}_{\mathcal{T}} A \cup \text{cl}_{\mathcal{T}} B$

**Proof** 1) Since  $\emptyset$  is closed,  $\text{cl}_{\mathcal{T}} \emptyset = \emptyset$

- 2)  $A \subseteq \bigcap \{F : F \text{ is closed and } A \subseteq F\} = \text{cl}_{\mathcal{T}} A$
- 3)  $\Rightarrow$  : If  $A$  is closed, then  $A$  itself is one of the closed sets  $F$  used in the definition  $\text{cl}_{\mathcal{T}} A = \bigcap \{F : F \text{ is closed and } F \supseteq A\}$ , so  $A = \text{cl}_{\mathcal{T}} A$ .  
 $\Leftarrow$  : If  $A = \text{cl}_{\mathcal{T}} A$ , then  $A$  is closed because  $\text{cl}_{\mathcal{T}} A$  is an intersection of closed sets.
- 4)  $\text{cl}_{\mathcal{T}} A$  is closed, so by 3),  $\text{cl}_{\mathcal{T}} A = \text{cl}_{\mathcal{T}}(\text{cl}_{\mathcal{T}} A)$

5)  $A \cup B \supseteq A$ , so  $\text{cl}_{\mathcal{T}}(A \cup B) \supseteq \text{cl}_{\mathcal{T}} A$ . Similarly,  $\text{cl}_{\mathcal{T}}(A \cup B) \supseteq \text{cl}_{\mathcal{T}} B$ .  
Therefore  $\text{cl}_{\mathcal{T}}(A \cup B) \supseteq \text{cl}_{\mathcal{T}}(A) \cup \text{cl}_{\mathcal{T}}(B)$ .

On the other hand,  $\text{cl}_{\mathcal{T}} A \cup \text{cl}_{\mathcal{T}} B$  is a closed set that contains  $A \cup B$ , and therefore  $\text{cl}_{\mathcal{T}} A \cup \text{cl}_{\mathcal{T}} B \supseteq \text{cl}_{\mathcal{T}}(A \cup B)$ . •

The next theorem tells us that we can use an operator “cl” to create a topology on a set.

**Theorem 5.19 (The Closure Operator Theorem)** Suppose  $X$  is a set and that for each  $A \subseteq X$ , a subset  $\text{cl } A$  is defined (*that is, we have a function  $\text{cl} : \mathcal{P}(X) \rightarrow \mathcal{P}(X)$* ) in such a way that conditions 1), 2), 4) and 5) of Theorem 5.18 are satisfied. Define  $\mathcal{T} = \{O : \text{cl}(X - O) = X - O\}$ . Then  $\mathcal{T}$  is a topology on  $X$ , and  $\text{cl}$  is the closure operator for this topology (*that is,  $\text{cl} = \text{cl}_{\mathcal{T}}$* ).

*Note: i) Such a function  $\text{cl}$  is called a “Kuratowski closure operator,” or just “closure operator” for short.*

*ii) The Closure Operator Theorem does not ask that  $\text{cl}$  satisfy condition 3): initially, there is no topology on the given set  $X$ , so 3) would be meaningless. But 3) motivates the definition of  $\mathcal{T}$  as the collection of sets whose complements are unchanged when “cl” is applied..*

**Proof** (Numbers in parentheses refer to properties of “cl”)

First note that:

$$(*) \quad \text{if } A \subseteq B, \text{ then } \text{cl } A \subseteq \text{cl } A \cup \text{cl } B \stackrel{(5)}{=} \text{cl}(A \cup B) = \text{cl } B$$

$\mathcal{T}$  has the properties required for a topology on the set  $X$  :

- i)  $\stackrel{(2)}{X \subseteq \text{cl } X}$ , so  $X = \text{cl } X$ . Therefore  $\text{cl}(X - \emptyset) = \text{cl } X = X = X - \emptyset$ , so  $\emptyset \in \mathcal{T}$ .  
Also,  $\text{cl}(X - X) = \text{cl } \emptyset \stackrel{(1)}{=} \emptyset = X - X$ , so therefore  $X \in \mathcal{T}$ .

ii) Suppose  $O_\alpha \in \mathcal{T}$  for each  $\alpha \in A$ . For each particular  $\alpha_0 \in A$ , we have

$\text{cl}(X - \bigcup O_\alpha) \stackrel{(*)}{\subseteq} \text{cl}(X - O_{\alpha_0}) = X - O_{\alpha_0}$  because  $O_{\alpha_0} \in \mathcal{T}$ . This is true for every  $\alpha_0 \in A$ , so  $\text{cl}(X - \bigcup O_\alpha) \subseteq \bigcap(X - O_\alpha) = X - \bigcup O_\alpha$ .

But by 2), we know that  $X - \bigcup O_\alpha \subseteq \text{cl}(X - \bigcup O_\alpha)$ .

Therefore  $\text{cl}(X - \bigcup O_\alpha) = X - \bigcup O_\alpha$ , so  $\bigcup O_\alpha \in \mathcal{T}$ .

iii) If  $O_1$  and  $O_2$  are in  $\mathcal{T}$ , then  $\text{cl}(X - (O_1 \cap O_2)) = \text{cl}((X - O_1) \cup (X - O_2))$

$\stackrel{(5)}{=} \text{cl}(X - O_1) \cup \text{cl}(X - O_2) = (X - O_1) \cup (X - O_2)$  (since  $O_1$  and  $O_2 \in \mathcal{T}$ )  
 $= X - (O_1 \cap O_2)$ . Therefore  $O_1 \cap O_2 \in \mathcal{T}$ . Therefore  $\mathcal{T}$  is a topology on  $X$ .

Having this topology  $\mathcal{T}$  now gives us an associated closure operator  $\text{cl}_\mathcal{T}$  and we want to show that  $\text{cl}_\mathcal{T} = \text{cl}$ . First, we observe that for closed sets in  $(X, \mathcal{T})$ :

$$(\text{**}) \quad \text{cl}_\mathcal{T} B = B \text{ iff } B \text{ is closed in } (X, \mathcal{T}) \\ \text{ iff } X - B \in \mathcal{T} \text{ iff } \text{cl}(X - (X - B)) = X - (X - B) \text{ iff } \text{cl } B = B.$$

To finish, we must show that  $\text{cl } A = \text{cl}_\mathcal{T} A$  for every  $A \subseteq X$ .

$A \subseteq \text{cl}_\mathcal{T} A$ , so (\*) gives that  $\text{cl } A \subseteq \text{cl}(\text{cl}_\mathcal{T} A)$ . But  $\text{cl}_\mathcal{T} A$  is closed in  $(X, \mathcal{T})$ , so using  $B = \text{cl}_\mathcal{T} A$  in (\*\*) gives  $\text{cl } \text{cl}_\mathcal{T} A = \text{cl}_\mathcal{T} A$ . Therefore  $\text{cl } A \subseteq \text{cl}_\mathcal{T} A$ .

On the other hand,  $\text{cl } \text{cl } A \stackrel{(4)}{=} \text{cl } A$ , so using  $B = \text{cl } A$  in (\*\*) gives that  $\text{cl } A$  is closed in  $(X, \mathcal{T})$ .

$\stackrel{(2)}{\text{But }} \text{cl } A \supseteq A$ , so  $\text{cl } A$  is one of the closed sets in the intersection that defines  $\text{cl}_\mathcal{T} A$ . Therefore  $\text{cl } A \supseteq \text{cl}_\mathcal{T} A$ . Therefore  $\text{cl } A = \text{cl}_\mathcal{T} A$ . •

### Example 5.20

1) Let  $X$  be a set. For  $A \subseteq X$ , define  $\text{cl } A = \begin{cases} A & \text{if } A \text{ is finite} \\ X & \text{if } A \text{ is infinite} \end{cases}$ .

Then  $\text{cl}$  satisfies the conditions in the Closure Operator Theorem. Since  $\text{cl } A = A$  iff  $A$  is finite or  $A = X$ , the closed sets in the topology generated by  $\text{cl}$  are precisely  $X$  and the finite sets – that is,  $\text{cl}$  generates the cofinite topology on  $X$ .

2) For each subset  $A$  of  $\mathbb{R}$ , define

$$\text{cl } A = \{x \in \mathbb{R} : \text{there is a sequence } (a_n) \text{ in } A \text{ with each } a_n \geq x \text{ and } |a_n - x| \rightarrow 0\}.$$

It is easy to check that  $\text{cl}$  satisfies the hypotheses of the Closure Operator Theorem. Moreover, a set  $A$  is open in the corresponding topology iff  $\forall x \in A \ \exists b > x$  such that  $[x, b) \subseteq A$ . Therefore the topology generated by  $\text{cl}$  is the Sorgenfrey topology on  $\mathbb{R}$ . What happens in this example if “ $\geq$ ” is replaced by “ $>$ ” in the definition of  $\text{cl}$ ?

Since closures, interiors and Frontiers are all related, it shouldn't be surprising that we can also describe a topology by defining an appropriate “int” operator or “Fr” operator on a set  $X$ .

E1. Let  $X = \{0, 1, 2, \dots\} = \{0\} \cup \mathbb{N}$ . For  $O \subseteq X$ , let

$$\phi_O(n) = \text{the number of elements in } O \cap [1, n] = |O \cap [1, n]|. \text{ Then define } T = \{O : 0 \notin O \text{ or } (0 \in O \text{ and } \lim_{n \rightarrow \infty} \frac{\phi_O(n)}{n} = 1)\}.$$

- a) Prove that  $T$  is a topology on  $X$ .
- b) In any topological space: a point  $x$  is called a limit point of the set  $A$  if  $N \cap (A - \{x\}) \neq \emptyset$  for every neighborhood  $N$  of  $x$ . *Informally,  $x$  is a limit point of  $A$  means that there are points of  $A$ , other than  $x$  itself, arbitrarily close to  $x$ .* Prove that in any topological space, a set  $B$  is closed iff  $B$  contains all of its limit points.
- c) For  $(X, T)$  as defined in a): prove that  $x$  is a limit point of  $X$  if and only if  $x = 0$ .

E2. Suppose  $(X, T)$  is a topological space and that  $A_\alpha \subseteq X$  for each  $\alpha \in A$ .

- a) Prove that if  $\bigcup \{\text{cl } A_\alpha : \alpha \in A\}$  is closed, then

$$\bigcup \{\text{cl } A_\alpha : \alpha \in A\} = \text{cl}(\bigcup \{A_\alpha : \alpha \in A\})$$

(Note that “ $\subseteq$ ” is true for any collection of sets  $A_\alpha$ .)

- b) A family  $\{B_\alpha : \alpha \in A\}$  of subsets of  $X$  is called locally finite if each point  $x \in X$  has a neighborhood  $N$  such that  $N \cap B_\alpha \neq \emptyset$  for only finitely many  $\alpha$ 's. Prove that if  $\{B_\alpha : \alpha \in A\}$  is locally finite, then

$$\bigcup \{\text{cl } B_\alpha : \alpha \in A\} = \text{cl}(\bigcup \{B_\alpha : \alpha \in A\})$$

- c) Prove that in  $(X, T)$ , the union of a locally finite family of closed sets is closed.

E3. Suppose  $f : (X, d) \rightarrow (Y, s)$ . Let  $\mathcal{B}$  be a base for the topology  $T_s$  and let  $\mathfrak{S}$  be a subbase for  $T_s$ . Prove or disprove:  $f$  is continuous iff  $f^{-1}[O]$  is open for all  $O \in \mathcal{B}$  iff  $f^{-1}[O]$  is open for all  $O \in \mathfrak{S}$ .

E4. A space  $(X, T)$  is called a  $T_1$ -space if  $\{x\}$  is closed for every  $x \in X$ .

- a) Give an example where  $(X, T)$  is not a  $T_1$ -space and  $T$  is not the trivial topology.
- b) Prove that  $X$  is a  $T_1$ -space if and only if, given any two distinct points  $x, y \in X$ , each point is contained in an open set not containing the other point.
- c) Prove that in a  $T_1$ -space, each set  $\{x\}$  can be written as an intersection of open sets.
- d) Prove that a subspace of a  $T_1$ -space is a  $T_1$ -space.
- e) Prove that if a pseudometric space  $(X, d)$  is a  $T_1$ -space, then  $d$  must in fact be a metric.
- f) Prove that if  $X$  and  $Y$  are  $T_1$ -spaces, so is  $X \times Y$ .

E5.  $(X, \mathcal{T})$  is called a  *$T_2$ -space* ( or *Hausdorff space* ) if whenever  $x, y \in X$  and  $x \neq y$ , then there exist disjoint open sets  $U$  and  $V$  with  $x \in U$  and  $y \in V$ .

- a) Give an example of a space  $(X, \mathcal{T})$  which is a  $T_1$ -space but not a  $T_2$ -space (see E4.) .
- b) Prove that a subspace of a Hausdorff space is Hausdorff.
- c) Prove that if  $X$  and  $Y$  are Hausdorff, then so is  $X \times Y$ .

E6. Prove that every infinite  $T_2$ -space contains an infinite discrete subspace – that is, a subset which is discrete in the subspace topology (see E5).

E7. Suppose that  $(X, \mathcal{T})$  and  $(Y, \mathcal{T}')$  are topological spaces. Recall that the product topology on  $X \times Y$  is the topology for which a base is the collection of “open boxes”

$$\mathcal{B} = \{U \times V : U \in \mathcal{T}, V \in \mathcal{T}'\}$$

Therefore a set  $O \subseteq X \times Y$  is open in the product topology iff for all  $(x, y) \in O$ , there exist open sets  $U$  in  $X$  and  $V$  in  $Y$  such that  $(x, y) \in U \times V \subseteq O$ . (Note that the product topology on  $\mathbb{R} \times \mathbb{R}$  is the usual topology on  $\mathbb{R}^2$ . (We always assume that the topology used on a product of two spaces  $X \times Y$  is the product topology unless something different is explicitly stated.)

- a) Verify that  $\mathcal{B}$  is, in fact, a base for a topology on  $X \times Y$ .
- b) Consider the projection map  $\pi_1 : X \times Y \rightarrow X$ . Prove that if  $O$  is any open set in  $X \times Y$  (not necessarily a “box”), then  $\pi_1[O]$  is open in  $X$ . (We say  $\pi_1$  is an open map. The same is true for the projection  $\pi_2$ .)
- c) Prove that if  $A \subseteq X$  and  $B \subseteq Y$ , then  $\text{cl}_{X \times Y}(A \times B) = \text{cl}_X A \times \text{cl}_Y B$ . Use this to explain why “the product of two closed sets is closed in  $X \times Y$ .”
- d) Show that  $X \times Y$  has a countable base iff each of  $X$  and  $Y$  has a countable base.
- e) Show that there is a countable neighborhood base at  $(x, y) \in X \times Y$  iff there is a countable neighborhood base  $\mathcal{B}_x$  at  $x \in X$  and a countable neighborhood base  $\mathcal{B}_y$  at  $y \in Y$ .
- f) Suppose  $(X, d_1)$  and  $(Y, d_2)$  are pseudometric spaces. Define a pseudometric  $d$  on the set  $X \times Y$  by

$$d((x_1, y_1), (x_2, y_2)) = d_1(x_1, x_2) + d_2(y_1, y_2).$$

Prove that the product topology on  $X \times Y$  is the same as the topology  $\mathcal{T}_d$ .

Note:  $d$  is the analogue of the taxicab metric in  $\mathbb{R}^2$ . There are other equivalent pseudometrics that produce the product topology on  $X \times Y$ , for example

$$d'((x_1, y_1), (x_2, y_2)) = (d_1(x_1, x_2)^2 + d_2(y_1, y_2)^2)^{1/2}, \text{ and}$$

$$d''((x_1, y_1), (x_2, y_2)) = \max\{d_1(x_1, x_2), d_2(y_1, y_2)\}.$$

E8. Suppose  $A \subseteq (X, \mathcal{T})$ . The set  $A$  is called regular open if  $A = \text{int}(\text{cl } A)$  and  $A$  is regular closed if  $A = \text{cl}(\text{int } A)$ .

a) Show that for any subset  $B$

$$\begin{array}{lll} \text{i) } X - \text{cl } B & = & \text{int}(X - B) \\ \text{ii) } X - \text{int } B & = & \text{cl}(X - B) \end{array}$$

b) Give an example of a closed subset of  $\mathbb{R}$  which is not regular closed.

c) Show that the complement of a regular open set in  $(X, \mathcal{T})$  is regular closed and vice-versa.

d) Show that the interior of any closed set in  $(X, \mathcal{T})$  is regular open.

e) Show that the intersection of two regular open sets in  $(X, \mathcal{T})$  is regular open.

f) Give an example of the union of two regular open sets that is not regular open.

E9. In each part, prove the statement or provide a counterexample:

a) For any  $x$  in a topological space  $(X, \mathcal{T})$ ,  $\{x\}$  is equal to the intersection of all open sets containing  $x$ .

b) In a topological space, a finite set must always be closed.

c) Suppose we have topologies  $\mathcal{T}_\alpha$  on  $X$ , one for each  $\alpha \in A$ . Then  $\bigcap\{\mathcal{T}_\alpha : \alpha \in A\}$  is also a topology on  $X$ .

d) If  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are topologies on  $X$ , then there is a unique smallest topology  $\mathcal{T}_3$  on  $X$  such that  $\mathcal{T}_1 \cup \mathcal{T}_2 \subseteq \mathcal{T}_3$ .

e) Suppose, for each  $\alpha \in A$ , that  $\mathcal{T}_\alpha$  is a topology on  $X$ . Then there is a unique smallest topology  $\mathcal{T}$  on  $X$  such that for each  $\alpha$ ,  $\mathcal{T} \supseteq \mathcal{T}_\alpha$ .

E10. Assume that natural number, except 1, can be factored into primes; you shouldn't need any other information about prime numbers. For  $a \in \mathbb{Z}$  and  $d \in \mathbb{N}$ , let

$$B_{a,d} = \{..., a - 2d, a - d, a, a + d, a + 2d, ...\} = \{a + kd : k \in \mathbb{Z}\}$$

and let  $\mathcal{B} = \{B_{a,d} : a \in \mathbb{Z}, d \in \mathbb{N}\}$

a) Prove that  $\mathcal{B}$  is a base for a topology  $\mathcal{T}$  on  $\mathbb{Z}$

b) Show that each set  $B_{a,d}$  is closed in  $(\mathbb{Z}, \mathcal{T})$ ?

c) What is the set  $\bigcup\{B_{0,p} : p \text{ is a prime number}\}$ ?

d) Part c) tells you what famous fact about the set of prime numbers?

## 6. Countability Properties of Spaces

Countable sets are often easier to work with than uncountable sets, so it is not surprising that spaces with certain “countability properties” are viewed as desirable. Most of these properties have already been defined, but the definitions are collected together here for convenience.

**Definition 6.1**  $(X, \mathcal{T})$  is called

first countable if we can choose a countable neighborhood base  $\mathcal{B}_x$  at every point  $x \in X$ .  
(We also say that  $X$  satisfies the first axiom of countability.)

second countable if there is a countable base  $\mathcal{B}$  for the topology  $\mathcal{T}$ . (We also say that  $X$  satisfies the second axiom of countability.)

separable if there is a countable dense set  $D$  in  $X$

Lindelöf if whenever  $\mathcal{U}$  is a collection of open sets for which  $\bigcup \mathcal{U} = X$ , then there is a countable subcollection  $\mathcal{U}' = \{U_1, U_2, \dots, U_n, \dots\} \subseteq \mathcal{U}$  for which  $\bigcup \mathcal{U}' = X$ .

$\mathcal{U}$  is called an open cover of  $X$ , and  $\mathcal{U}'$  is called a subcover from  $\mathcal{U}$ . Thus,  $X$  is Lindelöf if “every open cover has a countable subcover.”)

### Example 6.2

1) A countable discrete space  $(X, \mathcal{T})$  is second countable because  $\mathcal{B} = \{\{x\} : x \in X\}$  is a countable base. Is  $X$  first countable? separable? Lindelöf?

2)  $\mathbb{R}$  is second countable because  $\mathcal{B} = \{(a, b) : a, b \in \mathbb{Q}\}$  is a countable base. Similarly,  $\mathbb{R}^n$  is second countable since the collection  $\mathcal{B}$  of boxes  $(x_1, y_1) \times (x_2, y_2) \times \dots \times (x_n, y_n)$  with rational endpoints is a countable base (check!).

3) Let  $X$  be a countable set with the cofinite topology  $\mathcal{T}$ .  $X$  has only countably many finite subsets (see Theorem I.11.1) so there are only countably many sets in  $\mathcal{T}$ .  $(X, \mathcal{T})$  is second countable because we could choose  $\mathcal{B} = \mathcal{T}$  as a countable base.

The following theorem implies that each of the spaces in Example 6.2 is also first countable, separable, and Lindelöf. (However, it is really worthwhile to try to verify these properties, in each example, directly from the definitions.)

**Theorem 6.3** A second countable topological space  $(X, \mathcal{T})$  is also separable, first countable, and Lindelöf.

**Proof** Let  $\mathcal{B} = \{O_1, O_2, \dots, O_n, \dots\}$  be a countable base for  $\mathcal{T}$ .

i) For each  $n$ , pick a point  $x_n \in O_n$  and let  $D = \{x_n : n \in \mathbb{N}\}$ . The countable set  $D$  is dense. To see this, notice that if  $U$  is any nonempty open set in  $X$ , then for some  $n$ ,  $x_n \in O_n \subseteq U$  so  $U \cap D \neq \emptyset$ . Therefore  $X$  is separable.

ii) For each  $x \in X$ , let  $\mathcal{B}_x = \{O \in \mathcal{B} : x \in O\}$ . Clearly  $\mathcal{B}_x$  is a neighborhood base at  $x$ , so  $X$  is first countable.

iii) Let  $\mathcal{U}$  be any open cover of  $X$ . If  $x \in X$ , then  $x \in$  some set  $U_x \in \mathcal{U}$ . For each  $x$ , we can then pick a basic open set  $O_x \in \mathcal{B}$  such that  $x \in O_x \subseteq U_x$ . Let  $\mathcal{V} = \{O_x : x \in X\}$ . Since each  $O_x \in \mathcal{B}$ , there can be only countably many different sets  $O_x$ : that is,  $\mathcal{V}$  may contain “repeats.” Eliminate any “repeats” and list only the different sets in  $\mathcal{V}$ , so  $\mathcal{V} = \{O_{x_1}, O_{x_2}, \dots, O_{x_n}, \dots\}$  where  $O_{x_n} \subseteq U_{x_n} \in \mathcal{U}$ . Every  $x$  is in one of the sets  $O_{x_n}$ , so  $\mathcal{U}' = \{U_{x_1}, U_{x_2}, \dots, U_{x_n}, \dots\}$  is a countable subcover from  $\mathcal{U}$ . Therefore  $X$  is Lindelöf. •

The following examples show that no other implications exist among the countability properties in Theorem 6.3.

#### Example 6.4

1) Suppose  $X$  is uncountable and let  $\mathcal{T}$  be the cofinite topology on  $X$ .

$(X, \mathcal{T})$  is separable since any infinite set is dense.

$(X, \mathcal{T})$  is Lindelöf. To see this, let  $\mathcal{U}$  be an open cover of  $X$ . Pick any one nonempty set  $U \in \mathcal{U}$ . Then  $X - U$  is finite, say  $X - U = \{x_1, \dots, x_n\}$ . For each  $x_i$ , pick a set  $U_i \in \mathcal{U}$  with  $x_i \in U_i$ . Then  $\mathcal{U}' = \{U\} \cup \{U_i : i = 1, \dots, n\}$  is a countable (actually, finite) subcover chosen from  $\mathcal{U}$ .

However  $X$  is not first countable (Example 4.5.3), and therefore, by Theorem 6.3,  $X$  is also not second countable.

2) Suppose  $X$  is uncountable. Define  $\mathcal{T} = \{O \subseteq X : O = \emptyset \text{ or } X - O \text{ is countable}\}$ .

$\mathcal{T}$  is a topology on  $X$  (*check!*) called the cocountable topology. A set  $C \subseteq X$  is closed iff  $C = X$  or  $C$  is countable. (*This is an “upscale” analogue of the cofinite topology.*)

An argument very similar to the one in the preceding example shows that  $(X, \mathcal{T})$  is Lindelöf. But  $(X, \mathcal{T})$  is not separable — every countable subset is closed and therefore not dense. By Theorem 6.3,  $(X, \mathcal{T})$  also cannot be second countable.

3) Suppose  $X$  is uncountable set and choose a particular point  $p \in X$ . Define  $\mathcal{T} = \{O \subseteq X : O = \emptyset \text{ or } p \in O\}$ . (*Check that  $\mathcal{T}$  is a topology.*)

$(X, \mathcal{T})$  is separable because  $\{p\}$  is dense.

$(X, \mathcal{T})$  is not Lindelöf — because the cover  $\mathcal{U} = \{\{x, p\} : x \in X\}$  has no countable subcover.

Is  $(X, \mathcal{T})$  first countable?

4) Suppose  $X$  is uncountable and let  $\mathcal{T}$  be the discrete topology on  $X$ . Then  $(X, \mathcal{T})$  is

<u>first countable</u>	because $\mathcal{B}_x = \{\{x\}\}$ is a neighborhood base at $x$
<u>not second countable</u>	because each open set $\{x\}$ would have to be in a base $\mathcal{B}$
<u>not separable</u>	because $\text{cl } D = D \neq X$ for any countable set $D$
<u>not Lindelöf</u>	because the cover $\mathcal{U} = \{\{x\} : x \in X\}$ has no countable subcover. (In fact, not a single set can be omitted from $\mathcal{U}$ : $\mathcal{U}$ has no proper subcover.)

For “special” topological spaces – pseudometrizable ones, for example – it turns out that things are better behaved. For example, we noted earlier that every pseudometric space  $(X, d)$  is first countable (*Example 4.3*). The following theorem shows that in  $(X, d)$  the other three countability properties are equivalent to each other: that is, either all of them are true in  $(X, d)$  or none are true.

**Theorem 6.5** Any pseudometric space  $(X, d)$  is first countable.  $(X, d)$  is second countable iff  $(X, d)$  is separable iff  $(X, d)$  is Lindelöf.

**Proof** i) Second countable  $\Rightarrow$  Lindelöf: by Theorem 6.3, this implication is true in any topological space.

ii) Lindelöf  $\Rightarrow$  separable: suppose  $(X, d)$  is Lindelöf. For each  $n \in \mathbb{N}$ , let  $\mathcal{U}_n = \{B_{\frac{1}{n}}(x) : n \in \mathbb{N}, x \in X\}$ . For each  $n$ ,  $\mathcal{U}_n$  is an open cover so  $\mathcal{U}_n$  has a countable subcover – that is, for each  $n$  we can find countably many  $\frac{1}{n}$ -balls that cover  $X$ : say  $X = \bigcup_{k=1}^{\infty} B_{\frac{1}{n}}(x_{n,k})$ . Let  $D$  be the set of centers of all these balls:  $D = \{x_{n,k} : n, k \in \mathbb{N}\}$ . For any  $x \in X$  and every  $n$ , we have  $x \in B_{\frac{1}{n}}(x_{n,k})$  for some  $k$ , so  $d(x, x_{n,k}) < \frac{1}{n}$ . Therefore, for every  $n$ ,  $B_{\frac{1}{n}}(x) \cap D \neq \emptyset$  – in other words,  $x$  can be approximated arbitrarily closely by points from  $D$ . Therefore  $D$  is dense, so  $(X, d)$  is separable.

iii) Separable  $\Rightarrow$  second countable: suppose  $(X, d)$  is separable and that  $D = \{x_1, x_2, \dots, x_k, \dots\}$  is a countable dense set. Let  $\mathcal{B} = \{B_{\frac{1}{n}}(x_k) : n, k \in \mathbb{N}\}$ .  $\mathcal{B}$  is a countable collection of open balls and we claim  $\mathcal{B}$  is a base for the topology  $\mathcal{T}_d$ .

Suppose  $y \in O \in \mathcal{T}_d$ . By the definition of “open,” there is an  $\epsilon > 0$  for which  $B_\epsilon(y) \subseteq O$ . Pick  $n$  so that  $\frac{1}{n} < \frac{\epsilon}{2}$ , and pick  $x_k \in D$  so that  $d(x_k, y) < \frac{1}{n}$ . Then  $y \in B_{\frac{1}{n}}(x_k) \subseteq B_\epsilon(y) \subseteq O$  (because  $z \in B_{\frac{1}{n}}(x_k)$   
 $\Rightarrow d(z, y) \leq d(z, x_k) + d(x_k, y) < \frac{1}{n} + \frac{1}{n} < 2(\frac{\epsilon}{2}) = \epsilon$ ). •

*It's customary to call a metric space that has these three equivalent properties a “separable metric space” rather than a “second countable metric space” or “Lindelöf metric space.”*

Theorem 6.5 implies that the spaces in parts 1), 2), 3) of Example 6.4 are not pseudometrizable. In general, to show a space  $(X, \mathcal{T})$  is not pseudometrizable we can i) show that it fails to have some property shared by all pseudometric spaces (for example, first countability), or ii) show that it has one but not all of the properties “second countable,” “Lindelöf,” or “separable.”

## Exercises

E11. Define  $\mathcal{T} = \{U \cup V : U \text{ is open in the usual topology on } \mathbb{R} \text{ and } V \subseteq \mathbb{P}\}$ .

a) Show that  $\mathcal{T}$  is a topology on  $\mathbb{R}$ . If  $x$  is irrational, describe an “efficient” neighborhood base at  $x$ . Do the same if  $x$  is rational.

b) Is  $(\mathbb{R}, \mathcal{T})$  first countable? second countable? Lindelöf? separable?

*The space  $(\mathbb{R}, \mathcal{T})$  is called the “scattered line.” We could change the definition of  $\mathcal{T}$  by replacing  $\mathbb{P}$  with some other set  $A \subseteq \mathbb{R}$ . This creates a space in which the set  $A$  is “scattered.”*

*Hint: See Example I.7.9.6. It is possible to find open intervals  $I_n$  such that  $\bigcup_{n=1}^{\infty} I_n \supseteq \mathbb{Q}$*

*and for which  $\sum_{n=1}^{\infty} \text{length}(I_n) < 1$ .*

E12. A point  $x \in (X, \mathcal{T})$  is called a condensation point if every neighborhood of  $x$  is uncountable.

a) Let  $C$  be the set of all condensation points in  $X$ . Prove that  $C$  is closed.

b) Prove that if  $X$  is second countable, then  $X - C$  is countable.

E13. Suppose  $(X, \mathcal{T})$  is a second countable space and let  $\mathcal{B}$  be a countable base for the topology. Suppose  $\mathcal{B}'$  is another base (not necessarily countable) for  $\mathcal{T}$  containing open sets all of which have some property  $P$ . (For example, “ $P$ ” could be “clopen” or “separable.”) Show that there is a countable base  $\mathcal{B}''$  consisting of open sets with property  $P$ .

*Hint: think about the Lindelöf property.*

E14. A space  $(X, \mathcal{T})$  is called *hereditarily Lindelöf* if every subspace of  $X$  is Lindelöf.

a) Prove that a second countable space is hereditarily Lindelöf.

*In any space, a point  $x$  is called a limit point of the set  $A$  if  $N \cap (A - \{x\}) \neq \emptyset$  for every neighborhood  $N$  of  $x$ . Informally,  $x$  is a limit point of  $A$  if there are points in  $A$  different from  $x$  and arbitrarily close to  $x$ .*

b) Suppose  $X$  is hereditarily Lindelöf. Prove that the set  $A = \{x \in X : x \text{ is not a limit point of } X\}$  is countable.

E15. A space  $(X, \mathcal{T})$  is said to satisfy the *countable chain condition* ( $= CCC$ ) if every family of disjoint open sets must be countable.

- a) Prove that a separable space  $(X, \mathcal{T})$  satisfies the CCC.
- b) Give an example of a space that satisfies the CCC but that is not separable. (*It is not necessary to do so, but can you find an example which is a metric space?*)

E16. Suppose  $(X, \mathcal{T})$  is a topological space and that  $\mathcal{P}$  and  $\mathcal{B}$  are two bases for the topology  $\mathcal{T}$ , and that  $\mathcal{P}$  and  $\mathcal{B}$  are infinite.

a) Prove that there is a subfamily  $\mathcal{B}' \subseteq \mathcal{B}$  such that  $\mathcal{B}'$  is also a base and  $|\mathcal{B}'| \leq |\mathcal{P}|$ .  
*(Hint: For each pair  $t = (U, V) \in \mathcal{P} \times \mathcal{P}$ , pick, if possible, a set  $W_t \in \mathcal{B}$  such that  $U \subseteq W \subseteq V$ ; otherwise set  $W_t = \emptyset$ .)*

b) Use part a) to prove that the Sorgenfrey line is not second countable.  
*(Hint: Show that otherwise there would be a countable base of sets of the form  $[a, b)$ , but that this is impossible.)*

## 7. More About Subspaces

Suppose  $(X, \mathcal{T})$  is a topological space. In Definition 3.1, we defined the subspace topology  $\mathcal{T}_A$  on  $A \subseteq X$  :  $\mathcal{T}_A = \{A \cap O : O \in \mathcal{T}\}$ . In this section we explore some simple but important properties of subspaces.

If  $A \subseteq B \subseteq X$ , there are two ways to put a topology on  $A$  :

- 1) we can give  $A$  the subspace topology  $\mathcal{T}_A$  from  $X$ , or
- 2) we can give  $B$  the subspace topology  $\mathcal{T}_B$ , and then give  $A$  the subspace topology from the space  $(B, \mathcal{T}_B)$  – that is, we can give  $A$  the topology  $(\mathcal{T}_B)_A$ .

In other words, we can think of  $A$  as a subspace of  $X$  or as a subspace of the subspace  $B$ . Fortunately, the next theorem says that these two topologies are the same. More informally, Theorem 7.1 says that in a space  $X$  “a subspace of a subspace is a subspace.”

**Theorem 7.1** If  $A \subseteq B \subseteq X$ , and  $\mathcal{T}$  is a topology on  $X$ , then  $\mathcal{T}_A = (\mathcal{T}_B)_A$ .

**Proof**  $U \in \mathcal{T}_A$  iff  $U = O \cap A$  for some  $O \in \mathcal{T}$  iff  $U = O \cap (B \cap A)$  iff  $U = (O \cap B) \cap A$ . But  $O \cap B \in \mathcal{T}_B$ , so the last equation holds iff  $U \in (\mathcal{T}_B)_A$ . •

We always assume a subset  $A$  has the subspace topology (unless something else is explicitly stated). The notation  $A \subseteq (X, \mathcal{T})$  emphasizes that  $A$  is considered a subspace, not merely a subset.

By definition, a set is open in the subspace topology on  $A$  iff it is the intersection with  $A$  of an open set in  $X$ . The same is also true for closed sets.

**Theorem 7.2** Suppose  $F \subseteq A \subseteq (X, \mathcal{T})$ .  $F$  is closed in  $A$  iff  $F = A \cap C$  where  $C$  is closed in  $X$ .

**Proof**  $F$  is closed in  $A$  iff  $A - F$  is open in  $A$  iff  $A - F = O \cap A$  (for some open set  $O$  in  $X$ ) iff  $F = A - (O \cap A) = (X - O) \cap A = C \cap A$  (where  $C = X - O$  is a closed set in  $X$ ). •

**Theorem 7.3** Suppose  $A \subseteq (X, \mathcal{T})$ .

- 1) Let  $a \in A$ . If  $\mathcal{B}_a$  is a neighborhood base at  $a$  in  $X$ , then  $\{B \cap A : B \in \mathcal{B}_a\}$  is a neighborhood base at  $a$  in  $A$ .
- 2) If  $\mathcal{B}$  is a base for  $\mathcal{T}$ , then  $\{B \cap A : B \in \mathcal{B}\}$  is a base for  $\mathcal{T}_A$ .

*With a slight abuse of notation, we can informally write these collections as  $\mathcal{B}_a \cap A$  and  $\mathcal{B} \cap A$ . Why is this an “abuse?” What do  $\mathcal{B}_a \cap A$  and  $\mathcal{B} \cap A$  mean if taken literally?*

**Proof** 1) Suppose  $a \in A$  and that  $N$  is a neighborhood of  $a$  in  $A$ . Then  $a \in \text{int}_A N \subseteq N$ , so there is an open set  $O$  in  $X$  such that  $a \in \text{int}_A N = O \cap A \subseteq O$ .


Since  $\mathcal{B}_a$  is a neighborhood base at  $a$  in  $X$ , there is a neighborhood  $B \in \mathcal{B}_a$  such that  $a \in B \subseteq O$ . Then  $a \in (\text{int}_X B) \cap A \subseteq B \cap A \in \mathcal{B}_a \cap A$ . Since  $(\text{int}_X B) \cap A$  is open in  $A$ , we see that  $B \cap A$  is a neighborhood of  $a$  in  $A$ . And since  $a \in B \cap A \subseteq O \cap A = \text{int}_A N \subseteq N$ , we see that  $\mathcal{B}_a \cap A$  is a neighborhood base at  $a$  in  $A$ . •

## 2) Exercise

Theorem 7.3 tells us that in the subspace  $A$  we can get a neighborhood base at a point  $a$  by choosing a neighborhood base at  $a$  in  $X$  and then restricting all its sets to  $A$ ; and that the same applies to a base for the subspace topology.

**Corollary 7.4** Every subspace of a first countable (or second countable) space  $(X, \mathcal{T})$  is first countable (or second countable).

**Example 7.5** Suppose  $S^1$  is a circle in  $\mathbb{R}^2$  and that  $p \in S^1 \subseteq \mathbb{R}^2$ .  $\mathcal{B}_p = \{B_\epsilon(p) : \epsilon > 0\}$  is a neighborhood base at  $p$  in  $\mathbb{R}^2$ , and therefore  $\mathcal{B}_p \cap S^1$  is a neighborhood base at  $p$  in the subspace  $S^1$ . The sets in  $\mathcal{B}_p \cap S^1$  are “open arcs on  $S^1$  containing  $p$ .” (See the figure,)



The following theorem relates closure in a subspace to closure in the larger space. It turns out to be a very useful technical observation.

**Theorem 7.6** Suppose  $A \subseteq B \subseteq (X, \mathcal{T})$ , then  $\text{cl}_B A = B \cap \text{cl}_X A$ .

**Proof**  $B \cap \text{cl}_X A$  is a closed set in  $B$  that contains  $A$ , so  $B \cap \text{cl}_X A \supseteq \text{cl}_B A$ .

On the other hand, suppose  $b \in B \cap \text{cl}_X A$ . To show that  $b \in \text{cl}_B A$ , pick an open set  $U$  in  $B$  that contains  $b$ . We need to show  $U \cap A \neq \emptyset$ . There is an open set  $O$  in  $X$  such that  $O \cap B = U$ . Since  $b \in \text{cl}_X A$ , we have that  $\emptyset \neq O \cap A = O \cap (B \cap A) = (O \cap B) \cap A = U \cap A$ . •

**Example 7.7** 1)  $\mathbb{Q} = \text{cl}_{\mathbb{Q}} \mathbb{Q} = \text{cl}_{\mathbb{R}} \mathbb{Q} \cap \mathbb{Q}$

$$2) \text{cl}_{(0,2]}(0,1) = \text{cl}_{\mathbb{R}}(0,1) \cap (0,2] = [0,1] \cap (0,2] = (0,1]$$

3) The analogous calculations are not necessarily not true for interiors and boundaries. For example:

$$\mathbb{Q} = \text{int}_{\mathbb{Q}} \mathbb{Q} \neq \text{int}_{\mathbb{R}} \mathbb{Q} \cap \mathbb{Q} = \emptyset, \text{ and}$$

$$\emptyset = \text{Fr}_{\mathbb{Q}} \mathbb{Q} \neq \text{Fr}_{\mathbb{R}} \mathbb{Q} \cap \mathbb{Q} = \mathbb{Q}.$$

Why does “cl” have a privileged position here? Is there a “reason” why you would expect a better connection between closures in  $A$  and closures in  $X$  than you would expect between interiors in  $A$  and interiors in  $X$ ?

**Definition 7.8** A property  $P$  of topological spaces is called hereditary if whenever a space  $X$  has property  $P$ , then every subspace  $A$  also has property  $P$ .

For example, Corollary 7.4 tells us that first and second countability are hereditary properties. Other hereditary properties include “finite cardinality” and “pseudometrizability.” On the other hand, “infinite cardinality” is not a hereditary property.

**Example 7.9**

1) Separability is not a hereditary property. For example, consider the Sorgenfrey plane  $X$  (see Example 5.4).  $X$  is separable because  $\mathbb{Q}^2$  is dense.

Consider the subspace  $D = \{(x, y) : x + y = 1\}$ . The set  $U = [a, a + 1) \times [b, b + 1)$  is open in  $X$  so if  $(a, b) \in D$ , then  $U \cap D = \{(a, b)\}$  is open in the subspace  $D$ . Therefore  $D$  is a discrete subspace of  $X$ , and an uncountable discrete space is not separable.

Similarly, the Moore plane  $\Gamma$  is separable (see Example 5.6); the  $x$ -axis in  $\Gamma$  is an uncountable discrete subspace which is not separable.

2) The Lindelöf property is not hereditary. Let  $A$  be an uncountable set and let  $X = A \cup \{p\}$ , where  $p$  is any additional point not in  $A$ . Put a topology on  $X$  by giving a neighborhood base at each point.

$$\begin{cases} \mathcal{B}_a = \{\{a\}\} \text{ for } a \in A \\ \mathcal{B}_p = \{B : p \in B \text{ and } X - B \text{ is finite}\} \end{cases}$$

(Check that the collections  $\mathcal{B}_x$  satisfy the hypotheses of the Neighborhood Base Theorem 5.2.)

If  $\mathcal{V}$  is an open cover of  $X$ , then  $p \in V$  for some  $V \in \mathcal{V}$ . By ii), every neighborhood of  $p$  in  $X$  has a finite complement, so  $X - V$  is finite. For each  $y$  in the finite set  $X - V$ , we can choose a set  $V_y \in \mathcal{V}$  with  $y \in V_y$ . Then  $\mathcal{V}' = \{V\} \cup \{V_y : y \in X - V\}$  is a countable (in fact, finite) subcover from  $\mathcal{V}$ , so  $X$  is Lindelöf.

The definition of  $\mathcal{B}_a$  implies that each point of  $A$  is isolated in  $A$ ; that is,  $A$  is an uncountable discrete subspace. Then  $\mathcal{U} = \{\{a\} : a \in A\}$  is an open cover of  $A$  that has no countable subcover. Therefore the subspace  $A$  is not Lindelöf.

Even when a property is not hereditary, it is sometimes “inherited” by certain subspaces – perhaps, for example, by closed subspaces, or by open subspaces. The next theorem illustrates this.

**Theorem 7.10** A closed subspace of a Lindelöf space is Lindelöf (*so we say that the Lindelöf property is “closed hereditary”*).

**Proof** Suppose  $K$  is a closed subspace of the Lindelöf space  $X$ . Let  $\mathcal{U} = \{U_\alpha : \alpha \in A\}$  be a cover of  $K$  by sets  $U_\alpha$  that are open in  $K$ . For each  $\alpha$ , there is an open set  $V_\alpha$  in  $X$  such that  $V_\alpha \cap K = U_\alpha$ .

Since  $K$  is closed,  $X - K$  is open and  $\mathcal{V} = \{\{X - K\}\} \cup \{V_\alpha : \alpha \in A\}$  is an open cover of  $X$ . But  $X$  is Lindelöf, so  $\mathcal{V}$  has a countable subcover from  $\mathcal{V}$ , say  $\mathcal{V}' = \{X - K, V_1, \dots, V_n, \dots\}$ . (*The set  $X - K$  might not be needed in  $\mathcal{V}'$ , but it can't hurt to include it.*) Clearly then, the collection  $\{U_1, \dots, U_n, \dots\}$  is a countable subcover of  $K$  from  $\mathcal{U}$ . •

*Note:*

1. *A little reflection on the proof shows that to prove  $K$  is Lindelöf, it would be equivalent to show that every cover of  $K$  by sets open in  $X$  has a countable subcover.*
2. *A space  $X$  with the property that every open cover has a finite subcover is called compact. See, for example, the space in Example 6.4.1.*

*An obvious “tweak” to the proof for Theorem 7.10 shows that a closed subspace of a compact space is compact. We will look at the properties of compact spaces in much more detail in Chapter 4 and beyond.*

## 8. Continuity

We first defined continuous functions between pseudometric spaces by using the distance function  $d$  to mimic the definition of continuity as given in calculus. But then we saw that our definition could be restated in other equivalent ways in terms of open sets, closed sets, or neighborhoods. For topological spaces, we do not have distance functions available to use in a definition of continuity. But we can still make a definition using neighborhoods (or open sets, or closed sets) since the neighborhoods of  $a$  describe “nearness” to  $a$ . Of course, the definition parallels the way neighborhoods describe continuity in pseudometric spaces.

**Definition 8.1** A function  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{T}')$  is continuous at  $a \in X$  if whenever  $N$  is a neighborhood of  $f(a)$ , then  $f^{-1}[N]$  is a neighborhood of  $a$ . We say  $f$  is continuous if  $f$  is continuous at each point of  $X$ .

The statement that  $f$  is continuous at  $a$  is clearly equivalent to each of the following statements: :

- i) for each neighborhood  $N$  of  $f(a)$  there is a neighborhood  $W$  of  $a$  such that  $f[W] \subseteq N$
- ii) for each open set  $V$  containing  $f(a)$  there is an open set  $U$  containing  $a$  such that  $f[U] \subseteq V$
- iii) for each basic open set  $V$  containing  $f(a)$  there is a basic open set  $U$  containing  $a$  such that  $f[U] \subseteq V$ .

In the following theorem, the conditions i) - iii) for continuity are the same as those in Theorem II.5.6 for pseudometric spaces. Condition iv) was not mentioned in Chapter II, but it is sometimes handy.

**Theorem 8.2** Suppose  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{T}')$ . The following are equivalent.

- i)  $f$  is continuous
- ii) if  $O \in \mathcal{T}'$ , then  $f^{-1}[O] \in \mathcal{T}$  (*the inverse image of an open set is open*)
- iii) if  $F$  is closed in  $Y$ , then  $f^{-1}[F]$  is closed in  $X$  (*the inverse image of a closed set is closed*)
- iv) for every  $A \subseteq X$  :  $f[\text{cl}_X(A)] \subseteq \text{cl}_Y(f[A])$ .

**Proof** The proof that i) - iii) are equivalent is identical to the proof for Theorem II.5.6 for pseudometric spaces. That proof was deliberately worded in terms of open sets, closed sets, and neighborhoods so that it would carry over to this new situation.

iii)  $\Rightarrow$  iv)  $A \subseteq f^{-1}[f[A]] \subseteq f^{-1}[\text{cl}_Y(f[A])]$ . Since  $\text{cl}_Y(f[A])$  is closed in  $Y$ , iii) tells us that  $f^{-1}[\text{cl}_Y(f[A])]$  is a closed set in  $X$  that contains  $A$ . Therefore  $\text{cl}_X(A) \subseteq f^{-1}[\text{cl}_Y(f[A])]$ , so  $f[\text{cl}_X(A)] \subseteq [\text{cl}_Y(f[A])]$ .

iv)  $\Rightarrow$  i) Suppose  $a \in X$  and that  $N$  is a neighborhood of  $f(a)$ . Let  $K = X - f^{-1}[N]$  and  $U = X - \text{cl}_X K \subseteq f^{-1}[N]$ .  $U$  is open, and we claim that  $a \in U$  – which will show that  $f^{-1}[N]$  is a neighborhood of  $a$ , completing the proof. So we need to show that  $a \notin \text{cl}_X K$ . But this is clear because: if  $a \in \text{cl}_X K$ , then using iv) would give us that  $f(a) \in f[\text{cl}_X K] \subseteq \text{cl}_Y f[K]$ , which is impossible since  $N \cap f[K] = \emptyset$ .  $\bullet$

**Example 8.3** Sometimes we want to know whether a certain property is “preserved by continuous functions” – that is, if  $X$  has property  $P$  and  $f : X \rightarrow Y$  is continuous and onto, must the image  $Y$  also have the property  $P$ ?

For example, condition iv) in Theorem 8.2 implies that continuous maps preserve separability. Suppose  $D$  is a countable dense set in  $X$ . Then  $f[D]$  is countable and  $f[D]$  is dense in  $Y$  because  $Y = f[X] = f[\text{cl } D] \subseteq \text{cl } f[D]$ .

By contrast, continuous maps do not preserve first countability: for example, let  $(Y, \mathcal{T})$  be any topological space. Let  $\mathcal{T}'$  be the discrete topology on  $Y$ .  $(Y, \mathcal{T}')$  is first countable and the identity map  $i : (Y, \mathcal{T}') \rightarrow (Y, \mathcal{T})$  is continuous and onto. Thus, every space  $(Y, \mathcal{T})$  is the continuous image of a first countable space.

Do continuous maps preserve other properties that we have studied – such as Lindelöf, second countable, or metrizable?

The following theorem makes a few simple and useful observations about continuity.

**Theorem 8.4** Suppose  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{T}')$ .

- 1) Let  $B = \text{ran}(f) \subseteq Y$ . Then  $f$  is continuous iff  $f : (X, \mathcal{T}) \rightarrow (B, \mathcal{T}'_B)$  is continuous. *In other words,  $B = \text{the range of } f$  (a subspace of the codomain  $Y$ ) is what matters for the continuity of  $f$ ; points of  $Y$  not in  $B$  (if any) are irrelevant. For example, the function  $\sin : \mathbb{R} \rightarrow \mathbb{R}$  is continuous iff the function  $\sin : \mathbb{R} \rightarrow [-1, 1]$  is continuous.*
- 2) Let  $A \subseteq X$ . If  $f$  is continuous, then  $f|A = g : A \rightarrow Y$  is continuous. *That is, the restriction of a continuous function to a subspace is continuous.* *For example,  $\sin : \mathbb{R} \rightarrow \mathbb{R}$  is continuous, so  $\sin : \mathbb{Q} \rightarrow [-1, 1]$  is continuous.* *(The second function is really  $\sin|_{\mathbb{Q}}$ , but it's abbreviated here to just “sin”.*
- 3) If  $\mathcal{S}$  is a subbase for  $\mathcal{T}'$  (in particular, if  $\mathcal{S}$  is a base), then  $f$  is continuous iff  $f^{-1}[U]$  is open whenever  $U \in \mathcal{S}$ . In other words, *to check continuity, it is sufficient to show that the inverse image of every subbasic open set is open.*

**Proof** 1) Exercise: the crucial observation is that if  $O \subseteq Y$ , then  $f^{-1}[O] = f^{-1}[B \cap O]$ .

2) If  $O$  is open in  $Y$ , then  $g^{-1}[O] = f^{-1}[O] \cap A$  – which is an open set in  $A$ .

3) Exercise: the proof depends only on the definition of a subbase and set theory:

$$\begin{aligned} f^{-1}[\bigcup U_\alpha : \alpha \in A] &= \bigcup \{f^{-1}[U_\alpha] : \alpha \in A\} \text{ and} \\ f^{-1}[\bigcap U_\alpha : \alpha \in A] &= \bigcap \{f^{-1}[U_\alpha] : \alpha \in A\} \quad \bullet \end{aligned}$$


**Example 8.5**

- 1) For any topological spaces  $X$  and  $Y$ , every constant function  $f : X \rightarrow Y$  must be continuous. (Suppose  $f(x) = y_0$  for all  $x$ . If  $O$  is open in  $Y$ , then  $f^{-1}[O] = ?$ )  
If  $X$  has the discrete topology and  $Y$  is any topological space, then every function  $f : X \rightarrow Y$  is continuous.

2) Suppose  $X$  has the trivial topology and that  $f : X \rightarrow \mathbb{R}$ . If  $f$  is not constant, then there are points  $a, b \in X$  for which  $f(a) \neq f(b)$ . Let  $I$  be an open set in  $\mathbb{R}$  containing  $f(a)$  but not  $f(b)$ . Then  $f^{-1}[I]$  is not open in  $X$  so  $f$  is not continuous. We conclude that  $f$  is continuous iff  $f$  is constant.

*In this example, we could replace  $\mathbb{R}$  by any metric space  $(Y, d)$ ; or, for that matter, by any topological space  $(Y, \mathcal{T})$  that has what property?*

3) Let  $X$  be a rectangle inscribed inside a circle  $Y$  centered at  $P$ . For  $a \in X$ , let  $f(a)$  be the point where the ray from  $P$  through  $a$  intersects  $Y$ . (*The function  $f$  is called a “central projection.”*) Then both  $f : X \rightarrow Y$  and  $f^{-1} : Y \rightarrow X$  are continuous bijections.



**Example 8.6 (Weak topologies)** Suppose  $X$  is a set. Let  $\mathcal{F} = \{f_\alpha : \alpha \in A\}$  be a collection of functions where each  $f_\alpha : X \rightarrow \mathbb{R}$ . If we put the discrete topology on  $X$ , then all of the functions  $f_\alpha$  will be continuous. But a topology on  $X$  smaller than the discrete topology might also make all the  $f_\alpha$ 's continuous. The smallest topology on  $X$  that makes all the given  $f_\alpha$ 's continuous is called the weak topology  $\mathcal{T}$  on  $X$  generated by the collection  $\mathcal{F}$ .

How can we describe that topology more directly?  $\mathcal{T}$  makes all the  $f_\alpha$ 's continuous iff for each open  $O \subseteq \mathbb{R}$  and each  $\alpha \in A$ , the set  $f_\alpha^{-1}[O]$  is in  $\mathcal{T}$ . Therefore the weak topology generated by  $\mathcal{F}$  is the smallest topology that contains all these sets. According to Example 5.17.5, this means that weak topology  $\mathcal{T}$  is the one for which the collection  $\mathcal{S} = \{f_\alpha^{-1}[O] : O \text{ open in } \mathbb{R}, \alpha \in A\}$  is a subbase. (*It is clearly sufficient here to use only basic open sets  $O$  from  $\mathbb{R}$  – that is, open intervals  $(a, b)$  : why? Would using all open sets  $O$  put any additional sets into  $\mathcal{T}$ ?*)

For example, suppose  $X = \mathbb{R}^2$  and that  $\mathcal{F} = \{\pi_1, \pi_2\}$  contains the two projection maps  $\pi_1(x, y) = x$  and  $\pi_2(x, y) = y$ . For an open interval  $U = (a, b) \subseteq \mathbb{R}$ ,  $\pi_1^{-1}[U]$  is the “open vertical strip”  $U \times \mathbb{R}$ ; and  $\pi_2^{-1}[V]$  is the “open horizontal strip”  $\mathbb{R} \times V$ . Therefore a subbase for the weak topology on  $\mathbb{R}^2$  generated by  $\mathcal{F}$  consists of all such open horizontal or vertical strips. Two such strips intersect in an “open box”  $(a, b) \times (c, d)$  in  $\mathbb{R}^2$ , so it is easy to see that the weak topology is the product topology on  $\mathbb{R} \times \mathbb{R}$ , that is, the usual topology of  $\mathbb{R}^2$ .

Suppose  $A \subseteq \mathbb{R}$  and that  $i : A \rightarrow \mathbb{R}$  is the identity function  $i(x) = x$ . What is the weak topology on the domain  $A$  generated by the collection  $\mathcal{F} = \{i\}$ ?

**Definition 8.7** A mapping  $f: (X, \mathcal{T}) \rightarrow (Y, \mathcal{T}')$  is called

open if whenever  $O$  is open in  $X$ , then  $f[O]$  is open in  $Y$ , and  
closed if whenever  $F$  is closed in  $X$ , then  $f[F]$  is closed in  $Y$ .

Suppose  $|X| > 1$ . Let  $\mathcal{T}$  be the discrete topology on  $X$  and let  $\mathcal{T}'$  be the trivial topology on  $X$ . The identity map  $i: (X, \mathcal{T}) \rightarrow (X, \mathcal{T}')$  is continuous but neither open nor closed, and  $i: (X, \mathcal{T}') \rightarrow (X, \mathcal{T})$  is both open and closed but not continuous. Open and closed maps are quite different from continuous maps – even when the mapping is a bijection! Here are some examples that are more interesting.

**Example 8.8**

1)  $f: [0, 2\pi) \rightarrow S^1 = \{(x, y) \in \mathbb{R}^2 : x^2 + y^2 = 1\}$  given by  $f(\theta) = (\cos \theta, \sin \theta)$ .

It is easy to check that  $f$  is continuous, one-to-one, and onto. The set  $F = [\pi, 2\pi)$  is closed in  $[0, 2\pi)$  but  $f[[\pi, 2\pi))$  is not closed in  $S^1$ . Also,  $[0, \pi)$  is open in  $[0, 2\pi)$  but  $f[[0, \pi))$  is not open in  $S^1$ . A continuous, one-to-one, onto mapping does not need to be open or closed!

2) Suppose  $X$  and  $Y$  are topological spaces and that  $f: X \rightarrow Y$  is a bijection. Then there is an inverse function  $g = f^{-1}: Y \rightarrow X$ , and  $f^{-1}$  is continuous iff  $f$  is open. To check this, consider an open set  $O$  in  $X$ . Then  $y \in g^{-1}[O]$  iff  $g(y) \in O$  iff  $y = f(g(y)) \in f[O]$ , so  $f[O] = g^{-1}[O]$ . So  $f[O]$  is open iff  $g^{-1}[O]$  is open. So, for a bijection  $f$ ,  $f$  is open iff  $f^{-1}$  is continuous.

If  $O$  is replaced in the argument by a closed set  $F \subseteq X$ , then similar reasoning shows that a bijection  $f$  is closed iff  $f^{-1}$  is continuous.

In part 1), the bijection  $f$  is not open and therefore  $f^{-1}: S^1 \rightarrow [0, 2\pi)$  is not continuous. (*Explain directly, without part 2), why  $f^{-1}$  is not continuous.*)

**Definition 8.9** A mapping  $f: (X, \mathcal{T}) \rightarrow (Y, \mathcal{T}')$  is called a homeomorphism if  $f$  is a bijection and  $f$  and  $f^{-1}$  are both continuous. If a homeomorphism  $f$  exists, we say that  $X$  and  $Y$  are homeomorphic and write  $X \simeq Y$ .

*Note: The term is “homeomorphism,” not “homomorphism” (a term from algebra). The etymologies are closely related: “-morphism” comes from the Greek word ( $\mu\sigma\phi\eta'$ ) for “shape” or “form.” The prefixes “homo” and “homeo” come from Greek words meaning “same” and “similar” respectively. There was a major dispute in western religious history, mostly during the 4<sup>th</sup> century AD, that hinged on the distinction between “homeo” and “homo.”*

As noted in the preceding example, we could also describe a homeomorphism as a “continuous open bijection” or a “continuous closed bijection.”

It is obvious that among topological spaces, homeomorphism is an equivalence relation, that is, for topological spaces  $X, Y$ , and  $Z$  :

- i)  $X \simeq X$
- ii) if  $X \simeq Y$ , then  $Y \simeq X$
- iii) if  $X \simeq Y$  and  $Y \simeq Z$ , then  $X \simeq Z$ .

### Example 8.10

- 1) The function  $f : [0, 2\pi) \rightarrow S^1$  given by  $f(\theta) = (\cos \theta, \sin \theta)$  is not a homeomorphism even though  $f$  is continuous, one-to-one, and onto.
- 2) The “central projection” from the rectangle to the circle (*Example 8.5.3*) is a homeomorphism.
- 3) It is easy to see that any two open intervals  $(a, b)$  in  $\mathbb{R}$  are homeomorphic (just use a linear map of one interval onto the other).  
The mapping  $\tan : (-\frac{\pi}{2}, \frac{\pi}{2}) \rightarrow \mathbb{R}$  is a homeomorphism, so that each nonempty open interval in  $\mathbb{R}$  is actually homeomorphic to  $\mathbb{R}$  itself.
- 4) If  $f : (X, d) \rightarrow (Y, s)$  is an isometry (onto) between metric spaces, then both  $f$  and  $f^{-1}$  are continuous, so  $f$  is a homeomorphism.

In general, a homeomorphism between metric spaces need not be an isometry. But, of course, an isometry is automatically a homeomorphism


- 5) The function  $f : \{\frac{1}{n} : n \in \mathbb{N}\} \rightarrow \mathbb{N}$  given by  $f(\frac{1}{n}) = n$  is a homeomorphism (*both spaces have the discrete topology!*) Topologically, these spaces are the identical: both are just countable infinite sets with the discrete topology.  $f$  is not an isometry

In general, two discrete spaces  $X$  and  $Y$  are homeomorphic iff they have the same cardinality: any bijection between them is a homeomorphism. Roughly speaking, “size is the only possible topological difference between two discrete spaces.”

6) Let  $P$  denote the “north pole” of the sphere

$$S^2 = \{(x, y, z) : x^2 + y^2 + z^2 = 1\} \subseteq \mathbb{R}^3.$$

The function  $f$  illustrated below is a “stereographic projection”. The arrow starts at  $P$ , runs for a while inside the sphere and then exits through the surface of the sphere at a point  $(x, y, z)$ . Let  $f(x, y, z)$  be the point where the tip of the arrow hits the  $xy$ -plane  $\mathbb{R}^2$ . In this way  $f$  maps each point in  $S^2 - \{P\}$  to a point in  $\mathbb{R}^2$ . The function  $f$  is a homeomorphism. (See the figure below. Consider the images or inverse images of open sets.)



In general, what is the significance of a homeomorphism  $f : X \rightarrow Y$ ?

i)  $f$  is a bijection so it sets up a perfect one-to-one correspondence between the points in  $X$  and  $Y$ :  $x \leftrightarrow y = f(x)$ . We can imagine that  $f$  just “renames” the points in  $X$ . There is also a perfect one-to-one correspondence between the subsets  $X_\alpha$  and  $Y_\alpha$  of  $X$  and  $Y$ :  $X_\alpha \leftrightarrow Y_\alpha = f[X_\alpha]$ . Because  $f$  is a bijection, each subset  $Y_\alpha \subseteq Y$  corresponds in this way to one and only one subset  $X_\alpha \subseteq X$ .

ii)  $f$  is a bijection, so  $f$  treats unions, intersections and complements “nicely”:

- a)  $f^{-1}[\bigcup Y_\alpha : \alpha \in A] = \bigcup\{f^{-1}[Y_\alpha] : \alpha \in A\}$
- b)  $f^{-1}[\bigcap Y_\alpha : \alpha \in A] = \bigcap\{f^{-1}[Y_\alpha] : \alpha \in A\}$ , and
- c)  $f[\bigcup X_\alpha : \alpha \in A] = \bigcup\{f[X_\alpha] : \alpha \in A\}$
- d)  $f[\bigcap X_\alpha : \alpha \in A] = \bigcap\{f[X_\alpha] : \alpha \in A\}$ , and
- e)  $f[X - C] = f[X] - f[C] = Y - f[C]$

(Actually a), b), c) are true for any function; but d) and e) depend on  $f$  being a bijection.)

These properties say that this correspondence between subsets preserves unions: if each  $X_\alpha \leftrightarrow Y_\alpha$ , then  $\bigcup X_\alpha \leftrightarrow \bigcup Y_\alpha = \bigcup f[X_\alpha] = f[\bigcup X_\alpha]$ . Similarly,  $f$  preserves intersection and complements.

iii) Finally if  $f$  and  $f^{-1}$  are continuous, then open (closed) sets in  $X$  correspond to open (closed) sets in  $Y$  and vice-versa.

The total effect is that all the “topological structure” in  $X$  is exactly “duplicated” in  $Y$  and vice versa: we can think of points, subsets, open sets and closed sets in  $Y$  are just “renamed” copies of their counterparts in  $X$ . Moreover  $f$  preserves unions, intersections and complements, so  $f$  also preserves all properties of  $X$  that can be defined by using unions, intersections and complements of open sets. For example, we can check that if  $f$  is a homeomorphism and  $A \subseteq X$ , then  $f[\text{int}_X A] = \text{int}_Y f[A]$ , that  $f[\text{cl}_X A] = \text{cl}_Y f[A]$ , and that  $f[\text{Fr}_X A] = \text{Fr}_Y f[A]$ . That is,  $f$  takes interiors to interiors, closures to closures, and boundaries to boundaries.

**Definition 8.11** A property  $P$  of topological spaces is called a topological property if, whenever a space  $X$  has property  $P$  and  $Y \simeq X$ , then the space  $Y$  also has property  $P$ .

If  $X$  and  $Y$  are homeomorphic, then the very definition of “topological property” says that  $X$  and  $Y$  have the same topological properties. Conversely, if two topological spaces  $X$  and  $Y$  have the same topological properties, then  $X$  and  $Y$  must be homeomorphic. (*Why? Let  $P$  be the property “is homeomorphic to  $X$ .”  $P$  is a topological property because if  $Y$  has  $P$  (that is, if  $Y \simeq X$ ) and  $Z \simeq Y$ , then  $Z$  also has  $P$ . Moreover,  $X$  has this property  $P$ , because  $X \simeq X$ . So if we assume that  $Y$  has the same topological properties as  $X$ , then  $Y$  has the property  $P$ , that is,  $Y$  is homeomorphic to  $X$ .*)

So we think of two homeomorphic spaces as “topologically identical” — they are homeomorphic iff they have exactly the same topological properties. We can show that two spaces are not homeomorphic by naming a topological property of one space that the other space doesn't possess.

**Example 8.12** Let  $P$  be the property that “every continuous real-valued function achieves a maximum value.” Suppose a space  $X$  has property  $P$  and that  $h : X \rightarrow Y$  is a homeomorphism. We claim that  $Y$  also has property  $P$ .

Let  $f$  be any continuous real-valued function defined on  $Y$ .

Then  $f \circ h : X \rightarrow \mathbb{R}$  is continuous :

$$\begin{array}{ccc} & \mathbb{R} & \\ g \nearrow & \uparrow f & \\ X \xrightarrow{h} & Y & \\ & h & \end{array}$$

By assumption,  $g = f \circ h$  achieves a maximum value at some point  $a \in X$ , and we claim that  $f$  must achieve a maximum value at the point  $b = h(a) \in Y$ . If not, then there is a point  $y \in Y$  where  $f(y) > f(b)$ . Let  $x = h^{-1}(y)$ . Then

$$g(x) = f(h(x)) = f(h(h^{-1}(y))) = f(y) > f(b) = f(h(a)) = g(a),$$

which contradicts the fact that  $g$  achieves a maximum value at  $a$ .

Therefore  $P$  is a topological property.

For example, the closed interval  $[0, 1]$  has property  $P$  discussed in Example 8.12 (*this is a well-known fact from elementary analysis – which we will prove later*). But  $(0, 1)$  and  $[0, 1]$  do not have this property  $P$  (*why?*). So we can conclude that  $[0, 1]$  is not homeomorphic to either  $(0, 1)$  or  $[0, 1]$ .

Some other simple examples of topological properties are cardinality, first and second countability, Lindelöf, separability, and (pseudo)metrizability. In the case of metrizability, for example:

If  $(X, d)$  is a metric space and  $f : (X, d) \rightarrow (Y, \mathcal{T})$  is a homeomorphism, then we can define a metric  $d'$  on  $Y$  as  $d'(a, b) = d(f^{-1}(a), f^{-1}(b))$  for  $a, b \in Y$ . You then need to check that  $\mathcal{T}_{d'} = \mathcal{T}$  (using the properties of a homeomorphism and the definition of  $d'$ ). This shows that  $(Y, \mathcal{T})$  is metrizable. Be sure you can do this!

## 9. Sequences

In Chapter II we saw that sequences are a useful tool for working with pseudometric spaces. In fact, sequences are sufficient to describe the topology in a pseudometric space – because the convergent sequences in  $(X, d)$  determine the closure of a set.

We can easily define convergent sequences in any topological space  $(X, \mathcal{T})$ . But, as we will see, sequences need to be used with more care in spaces that are not pseudometrizable. Whether or not a sequence  $(x_n)$  converges to a particular point  $x$  is a “local” question – it depends on “the  $x_n$ ’s approaching nearer and nearer to  $x$ ” and, in the absence of a distance function, we use the neighborhoods of  $x$  to determine “nearness to  $x$ .” If the neighborhood system  $\mathcal{N}_x$  is “too large” or “too complicated,” then it may be impossible for a sequence to “get arbitrarily close” to  $x$ . Soon we will see a specific example where such a difficulty actually occurs. But first, we look at some of the things that do work out just as nicely for topological spaces as they do in pseudometric spaces.

**Definition 9.1** Suppose  $(x_n)$  is a sequence in  $(X, \mathcal{T})$ . We say that  $(x_n)$  converges to  $x$  if, for every neighborhood  $W$  of  $x$ ,  $\exists k \in \mathbb{N}$  such that  $x_n \in W$  when  $n \geq k$ . In this case we write  $(x_n) \rightarrow x$ . More informally, we can say that  $(x_n) \rightarrow x$  if  $(x_n)$  is eventually in every neighborhood  $W$  of  $x$ .

*Clearly, we can replace “every neighborhood  $W$  of  $x$ ” in the definition with “every basic neighborhood  $B$  of  $x$ ” or “every open set  $O$  containing  $x$ .” Be sure you are convinced of this.*

In a pseudometric space a sequence can converge to more than one point, but we proved that in a metric space limits of convergent sequences must be unique. A similar distinction holds in topological spaces: the important issue is whether we can “separate points by open sets.”

**Definition 9.2**  $(X, \mathcal{T})$  is a  $T_1$ -space if whenever  $x \neq y \in X$  there exist open sets  $U$  and  $V$  such that  $x \in U$ ,  $y \notin U$  and  $y \in V$ ,  $x \notin V$  (that is, each point is in an open set that does not contain the other point).

$(X, \mathcal{T})$  is a  $T_2$ -space (or Hausdorff space) if whenever  $x \neq y \in X$  there exist disjoint open sets  $U$  and  $V$  such that  $x \in U$  and  $y \in V$ .

It is easy to check that

- i)  $X$  is a  $T_1$ -space iff for every  $x \in X$ ,  $\{x\}$  is closed, that
- ii) every  $T_2$ -space is a  $T_1$ -space
- iii) every metric space  $(X, d)$  is a  $T_2$ -space (Hausdorff space)

*There is a hierarchy of ever stronger “separation axioms” called  $T_0, T_1, T_2, T_3$ , and  $T_4$  that a topological space might satisfy. Eventually we will look at all of them.*

*Each condition is stronger than the preceding ones in the list (for example,  $T_2 \Rightarrow T_1$ ).*

*The letter “T” is used here because in the early (German) literature, the word for “separation axioms” was “Trennungsaxiome.”*

**Theorem 9.3** In a Hausdorff space  $(X, \mathcal{T})$ , a sequence can converge to at most one point.

**Proof** Suppose  $x \neq y \in X$ . Choose disjoint open sets  $U$  and  $V$  with  $x \in U$  and  $y \in V$ . If  $(x_n) \rightarrow x$ , then  $(x_n)$  is eventually in  $U$ , so  $(x_n)$  is not eventually in  $V$ . Therefore  $(x_n)$  does not also converge to  $y$ . •

When we try to generalize results from pseudometric spaces to topological spaces, we often get a better insight about where the heart of a proof lies. For example, to prove that limits of sequences are unique it is the Hausdorff property that is important, not the presence of a metric. Here is another example: for a pseudometric space  $(X, d)$  we proved that  $x \in \text{cl } A$  iff  $x$  is a limit of a sequence  $(a_n)$  in  $A$ . That proof (see Theorem II.5.18) used the fact that there was a countable neighborhood base  $\{B_{\frac{1}{n}}(x) : n \in \mathbb{N}\}$  at each point  $x$ . We can see now that the countable neighborhood base was the crucial fact – because we can prove the same result in any first countable topological space  $(X, \mathcal{T})$ .

But first, two technical lemmas are helpful.

**Lemma 9.4** Suppose  $\{N_1, N_2, \dots, N_k, \dots\}$  is a countable neighborhood base at  $a \in X$ . Define  $U_k = \text{int}(N_1 \cap \dots \cap N_k)$ . Then  $\{U_1, U_2, \dots, U_k, \dots\}$  is also a neighborhood base at  $a$ .

**Proof**  $N_1 \cap \dots \cap N_k$  is a neighborhood of  $a$ , so  $a \in \text{int}(N_1 \cap \dots \cap N_k) = U_k$ . Therefore  $U_k$  is an open neighborhood of  $a$ . If  $N$  is any neighborhood of  $a$ , then  $a \in N_k \subseteq N$  for some  $k$ , so then  $a \in U_k \subseteq N_k \subseteq N$ . Therefore the  $U_k$ 's are a neighborhood base at  $a$ . •

The exact formula for the  $U_k$ 's in Lemma 9.4 doesn't matter; the important thing is that we get a “much improved” neighborhood base  $\{U_1, U_2, \dots, U_k, \dots\}$  – one in which the  $U_k$ 's are open and  $U_1 \supseteq U_2 \supseteq \dots \supseteq U_k \supseteq \dots$ . This new neighborhood base at  $a$  plays a role analogous to the neighborhood base  $B_1(a) \supseteq B_{\frac{1}{2}}(a) \supseteq \dots \supseteq B_{\frac{1}{k}}(a) \supseteq \dots$  in a pseudometric space. We call  $\{U_1, U_2, \dots, U_k, \dots\}$  an open, shrinking neighborhood base at  $a$ .

**Lemma 9.5** Suppose  $\{U_1, U_2, \dots, U_k, \dots\}$  is a shrinking neighborhood base at  $x$  and that  $a_n \in U_n$  for each  $n$ . Then  $(a_n) \rightarrow x$ .

**Proof** If  $W$  is any neighborhood of  $x$ , then there is a  $k$  such that  $x \in U_k \subseteq W$ . Since the  $U_k$ 's are a shrinking neighborhood base, we have that for any  $n \geq k$ ,  $a_n \in U_n \subseteq U_k \subseteq W$ . So  $(a_n) \rightarrow x$ . •

**Theorem 9.6** Suppose  $(X, \mathcal{T})$  is first countable and  $A \subseteq X$ . Then  $x \in \text{cl } A$  iff there is a sequence  $(a_n)$  in  $A$  such that  $(a_n) \rightarrow x$ . (More informally, “sequences are sufficient” to describe the topology in a first countable topological space.)

**Proof** ( $\Leftarrow$ ) Suppose  $(a_n)$  is a sequence in  $A$  and that  $(a_n) \rightarrow x$ . For each neighborhood  $W$  of  $x$ ,  $(a_n)$  is eventually in  $W$ . Therefore  $W \cap A \neq \emptyset$ , so  $x \in \text{cl } A$ . (This half of the proof works in any topological space: it does not depend on first countability.)

( $\Rightarrow$ ) Suppose  $x \in \text{cl } A$ . Using Lemma 9.4, choose a countable shrinking neighborhood base  $\{U_1, \dots, U_n, \dots\}$  at  $x$ . Since  $x \in \text{cl } A$ , we can choose a point  $a_n \in U_n \cap A$  for each  $n$ . By Lemma 9.5,  $(a_n) \rightarrow x$ . •

We can use Theorem 9.6 to get an upper bound on the size of certain topological spaces, analogous to what we did for pseudometric spaces. This result is not very important, but it illustrates that in Theorem II.5.21 the properties that are really important are “first countability” and “Hausdorff,” not the actual presence of a metric  $d$ .

**Corollary 9.7** If  $D$  is a dense subset in a first countable Hausdorff space  $(X, \mathcal{T})$ , then  $|X| \leq |D|^{\aleph_0}$ . In particular, If  $X$  is a separable, first countable Hausdorff space, then  $|X| \leq \aleph_0^{\aleph_0} = c$ .

**Proof**  $X$  is first countable, so for each  $x \in X$  we can pick a sequence  $(d_n)$  in  $D$  such that  $(d_n) \rightarrow x$ ; formally, this sequence is a function  $f_x : \mathbb{N} \rightarrow D$ , so  $f_x \in D^{\mathbb{N}}$ . Since  $X$  is Hausdorff, a sequence cannot converge to two different points: so if  $x \neq y \in X$ , then  $f_x \neq f_y$ . Therefore the function  $\Phi : X \rightarrow D^{\mathbb{N}}$  given by  $\Phi(x) = f_x$  is one-to-one, so  $|X| \leq |D^{\mathbb{N}}| = |D|^{\aleph_0}$ . •

The conclusion in Theorem 9.6 may not be true if  $X$  is not first countable: sequences are not always “sufficient to describe the topology” of  $X$  – that is, convergent sequences cannot always determine the closure of a set.

### Example 9.8 (the space $L$ )

Let  $L = \{(m, n) : m, n \in \mathbb{Z}, m, n \geq 0\}$ , and let  $C_j$  be “the  $j^{\text{th}}$  column of  $L$ ,” that is  $C_j = \{(j, n) \in L : n = 0, 1, \dots\}$ . We put a topology on  $L$  by giving a neighborhood base at each point  $p$ :

$$\mathcal{B}_p = \begin{cases} \{(m, n)\} & \text{if } p = (m, n) \neq (0, 0) \\ \{B : (0, 0) \in B \text{ and } C_j - B \text{ is finite for all but finitely many } j\} & \text{if } p = (0, 0) \end{cases}$$

(Check that this definition satisfies the conditions in the Neighborhood Base Theorem 5.2 and therefore does describe a topology for  $L$ .)

If  $p \neq (0, 0)$  then  $p$  is isolated in  $L$ . A basic neighborhood of  $(0, 0)$  is a set which contains  $(0, 0)$  and which, we could say, contains “most of the points from most of the columns.” With this topology,  $L$  is a Hausdorff space.

Certainly  $(0, 0) \in \text{cl}(L - \{(0, 0)\})$ , but no sequence from  $L - \{(0, 0)\}$  converges to  $(0, 0)$ . To see this, consider any sequence  $(a_n)$  in  $L - \{(0, 0)\}$ :

- i) if there is a column  $C_{j_0}$  that contains infinitely many of the terms  $a_n$ , then  $N = (L - C_{j_0}) \cup \{(0, 0)\}$  is a neighborhood of  $(0, 0)$  and  $(a_n)$  is not eventually in  $N$ .
- ii) if every column  $C_j$  contains only finitely many  $a_n$ 's, then  $N = L - \{a_n : n \in \mathbb{N}\}$  is a neighborhood of  $(0, 0)$  and  $(a_n)$  is not eventually in  $N$  (in fact, the sequence is never in  $N$ ).

In  $L$ , sequences are not sufficient to describe the topology: convergent sequences can't show us that  $(0, 0) \in \text{cl}(L - \{(0, 0)\})$ . According to Theorem 9.6, this means that the countable space  $L$  cannot be first countable – there is a countable neighborhood base at each point  $p \neq (0, 0)$  but not at  $(0, 0)$ .

The neighborhood system at  $(0, 0)$  “measures nearness to”  $(0, 0)$  but the ordering relationship ( $\supseteq$ ) among the basic neighborhoods at  $(0, 0)$  is very complicated – much more complicated than the neat, simple nested chain of neighborhoods  $U_1 \supseteq U_2 \supseteq \dots \supseteq U_n \supseteq \dots$  that could form a base at  $x$  in a first countable space. Roughly, the complexity of the neighborhood system is the reason why the terms of a sequence can't get “arbitrarily close” to  $(0, 0)$ .

Since sequences do suffice to describe the topology in a first countable space, it is not surprising that we can use sequences to determine the continuity of a function defined on a first countable space  $X$ .

**Theorem 9.9** Suppose  $(X, \mathcal{T})$  is first countable and  $f : (X, \mathcal{T}) \rightarrow (Y, \mathcal{T}')$ . Then  $f$  is continuous at  $a \in X$  iff whenever  $(x_n) \rightarrow a$ , then  $(f(x_n)) \rightarrow f(a)$ .

**Proof** ( $\Rightarrow$ ) If  $f$  is continuous at  $a$  and  $W$  is a neighborhood of  $f(a)$ , then  $f^{-1}[W]$  is a neighborhood of  $a$ . Therefore  $(x_n)$  is eventually in  $f^{-1}[W]$ , so  $f((x_n))$  is eventually in  $W$ . (This half of the proof is valid for any topological space  $X$ : continuous functions always “preserve convergent sequences.”)

( $\Leftarrow$ ) Let  $\{U_1, \dots, U_n, \dots\}$  be a shrinking neighborhood base at  $a$ . If  $f$  is not continuous at  $a$ , then there is a neighborhood  $W$  of  $f(a)$  such that for every  $n$ ,  $f[U_n] \not\subseteq W$ . For each  $n$ , choose a point  $x_n \in U_n - f^{-1}[W]$ . Then (since the  $U_n$ 's are shrinking) we have  $(x_n) \rightarrow a$  but  $(f(x_n))$  fails to converge to  $f(a)$  because  $f(x_n)$  is never in  $W$ . •  
(Compare this to the proof of Theorem II.5.22.)

## 10. Subsequences

**Definition 10.1** Suppose  $f : \mathbb{N} \rightarrow X$  is a sequence in  $X$  and that  $\phi : \mathbb{N} \rightarrow \mathbb{N}$  is strictly increasing. The composition  $f \circ \phi : \mathbb{N} \rightarrow X$  is called a subsequence of  $f$ .

$$\begin{array}{ccc} & f & \\ \mathbb{N} & \longrightarrow & X \\ \phi \uparrow & \nearrow & f \circ \phi \\ & \mathbb{N} & \end{array}$$

If we write  $f(n) = x_n$  and  $\phi(k) = n_k$ , then  $(f \circ \phi)(k) = f(n_k) = x_{n_k}$ . We write the sequence  $f$  informally as  $(x_n)$  and the subsequence  $f \circ \phi$  as  $(x_{n_k})$ . Since  $\phi$  is increasing, we have that  $n_k \rightarrow \infty$  as  $k \rightarrow \infty$ .

For example, if  $\phi(k) = n_k = 2k$ , then  $f \circ \phi$  is the subsequence written informally as  $(x_{n_k}) = (x_{2k})$ , that is, the subsequence  $(x_2, x_4, x_6, \dots, x_{2k}, \dots)$ . But if  $\phi(n) = 1$  for all  $n$ , then  $f \circ \phi$  is not a subsequence: informally,  $(x_1, x_1, x_1, \dots, x_1, \dots)$  is not a subsequence of  $(x_1, x_2, x_3, \dots, x_n, \dots)$ . Every sequence  $f$  is a subsequence of itself: just let  $\phi(n) = n$ .

**Theorem 10.2** Suppose  $x \in (X, \mathcal{T})$ . Then  $(x_n) \rightarrow x$  iff every subsequence  $(x_{n_k}) \rightarrow x$ .

**Proof** ( $\Leftarrow$ ) This is clear because  $(x_n)$  is a subsequence of itself.

( $\Rightarrow$ ) Suppose  $(x_n) \rightarrow x$  and that  $(x_{n_k})$  is a subsequence. If  $W$  is any neighborhood of  $x$ , then  $x_n \in W$  for all  $n >$  some  $n_0$ . Since the  $n_k$ 's are strictly increasing,  $n_k > n_0$  for all  $k >$  some  $k_0$ . Therefore  $(x_{n_k})$  is eventually in  $W$ , so  $(x_{n_k}) \rightarrow x$ . •

**Definition 10.3** Suppose  $x \in (X, \mathcal{T})$ . We say that  $x$  is a cluster point of the sequence  $(x_n)$  if for each neighborhood  $W$  of  $x$  and for each  $k \in \mathbb{N}$ , there is an  $n > k$  for which  $x_n \in W$ . More informally, we say that  $x$  is a cluster point of  $(x_n)$  if the sequence is frequently in every neighborhood  $W$  of  $x$ .

**Definition 10.4** Suppose  $x \in (X, \mathcal{T})$  and  $A \subseteq X$ . We say that  $x$  is a limit point of  $A$  if  $N \cap (A - \{x\}) \neq \emptyset$  for every neighborhood  $N$  of  $x$  – that is, every neighborhood of  $x$  contains points of arbitrarily close to  $x$  but different from  $x$ .

**Example 10.5.**

- 1) Suppose  $X = [0, 1] \cup \{2\}$  and  $A = \{2\}$ . Then  $W \cap A \neq \emptyset$  for every neighborhood  $W$  of 2, but 2 is not a limit point of  $A$  – because  $W = \{2\}$  is a neighborhood of 2 in  $X$  and  $W \cap (A - \{2\}) = \emptyset$ . Each  $x \in [0, 1]$  is a limit point of  $[0, 1]$  and also a limit point of  $X$ . Since  $\{2\}$  is open in  $X$ , 2 is also not a limit point of  $X$ .
- 2) Every point  $r$  in  $\mathbb{R}$  is a limit point of  $\mathbb{Q}$ . If  $A \subseteq \mathbb{N}$ , then  $A$  has no limit points (in  $\mathbb{N}$  or in  $\mathbb{R}$ ).
- 3) If  $(x_n) \rightarrow x$ , then  $x$  is a cluster point of  $(x_n)$ . More generally, if  $(x_n)$  has a subsequence  $(x_{n_k})$  that converges to  $x$ , then  $x$  is a cluster point of  $(x_n)$ . (Why?)

4) A sequence can have many cluster points. For example, if the sequence  $(q_n)$  lists all the elements of  $\mathbb{Q}$ , then every  $r \in \mathbb{R}$  is a cluster point of  $(q_n)$ .

5) In  $\mathbb{R}$ , the sequence  $(x_n) = ((-1)^n)$  has exactly two cluster points:  $-1$  and  $1$ . But the set  $\{x_n : n \in \mathbb{N}\} = \{-1, 1\}$  has no limit points in  $\mathbb{R}$ . The set of cluster points of a sequence is not always the same as the set of limit points of the set of terms in the sequence! (*Is one of these sets always a subset of the other?*)

**Theorem 10.6** Suppose  $a$  is a cluster point of  $(x_n)$  in a first countable space  $(X, \mathcal{T})$ . Then there is a subsequence  $(x_{n_k}) \rightarrow a$ .

**Proof** Let  $\{U_1, U_2, \dots, U_n, \dots\}$  be a countable shrinking neighborhood base at  $a$ . Since  $(x_n)$  is frequently in  $U_1$ , we can pick  $n_1$  so that  $x_{n_1} \in U_1$ . Since  $(x_n)$  is frequently in  $U_2$ , we can pick an  $n_2 > n_1$  so that  $x_{n_2} \in U_2 \subseteq U_1$ . Continue inductively: having chosen  $n_1 < \dots < n_k$  so that  $x_{n_k} \in U_k \subseteq \dots \subseteq U_1$ , we can then choose  $n_{k+1} > n_k$  so that  $x_{n_{k+1}} \in U_{k+1} \subseteq U_k$ . Then  $(x_{n_k})$  is a subsequence of  $(x_n)$  and  $(x_{n_k}) \rightarrow a$ . •

### Example 10.7 (the space $L$ , revisited)

Let  $L$  be the space in Example 9.8 and let  $(x_n)$  be a sequence which lists all the elements of  $L - \{(0, 0)\}$ .

Every basic neighborhood  $B$  of  $(0, 0)$  is infinite, so  $B$  must contain terms  $x_n$  for arbitrarily large  $n$ . This means that  $(x_n)$  is frequently in  $B$ , so  $(0, 0)$  is a cluster point of  $(x_n)$ .

But no subsequence of  $(x_n)$  can converge to  $(0, 0)$  – because we showed in Example 9.8 that no sequence whatsoever from  $L - \{(0, 0)\}$  can converge to  $(0, 0)$ . Therefore Theorem 10.6 may not be true if the space  $X$  is not first countable.

Consider any sequence  $(x_n) \rightarrow (0, 0)$  in  $L$ . If there were infinitely many  $x_n \neq (0, 0)$ , then we could form the subsequence that contains those terms, and that subsequence would be a sequence in  $L - \{(0, 0)\}$  that converges to  $(0, 0)$  – which is impossible. Therefore we conclude that eventually  $x_n = (0, 0)$ .

Suppose now that  $f : L \rightarrow \mathbb{N}$  is any bijection, and let  $k = f(0, 0)$ .

- whenever a sequence  $(x_n) \rightarrow (0, 0)$  in  $L$ , then  $(f(x_n)) \rightarrow f(0, 0) = k$  in  $\mathbb{N}$  (*because, by the preceding paragraph,  $f(x_n) = f(0, 0) = k$  eventually*)
- the topology on  $\mathbb{N}$  is discrete so  $\{k\}$  is a neighborhood of  $f(0, 0)$ , but  $f^{-1}[\{k\}] = \{(0, 0)\}$  is not a neighborhood of  $(0, 0)$ . Therefore  $f$  is not continuous at  $(0, 0)$

Theorem 9.9 does not apply to  $L$ : if a space is not first countable, sequences may be inadequate to check whether a function  $f$  is continuous at a point.

## Exercises

E17. Suppose  $f, g : (X, \mathcal{T}_1) \rightarrow (Y, \mathcal{T}_2)$  are continuous functions, that  $D$  is dense in  $X$ , and that  $f|D = g|D$ . Prove that if  $Y$  is Hausdorff, then  $f = g$ . (This generalizes the result in Chapter 2, Theorem 5.12.)

E18. A function  $f : (X, \mathcal{T}) \rightarrow \mathbb{R}$  is called lower semicontinuous if

$$f^{-1}[(b, \infty)] = \{x : f(x) > b\} \text{ is open for every } b \in \mathbb{R},$$

and  $f$  is called upper semicontinuous if

$$f^{-1}[(-\infty, b)] = \{x : f(x) < b\} \text{ is open for each } b \in \mathbb{R}.$$

- a) Show that  $f$  is continuous iff  $f$  is both upper and lower semicontinuous.
- b) Give an example of a lower semicontinuous  $f : \mathbb{R} \rightarrow \mathbb{R}$  which is not continuous. Do the same for upper semicontinuous.
- c) Suppose  $A \subseteq X$ . Prove that the characteristic function  $\chi_A$  is lower semicontinuous if  $A$  is open in  $X$  and upper semicontinuous if  $A$  is closed in  $X$ .

E19. Suppose  $X$  is an infinite set with the cofinite topology, and that  $Y$  has the property that every singleton  $\{y\}$  is a closed set. (You might want to check: this is equivalent to saying that  $Y$  is a  $T_1$ -space: see Definition 9.2). Prove that if  $f : X \rightarrow Y$  is continuous and onto, then either  $f$  is constant or  $X$  is homeomorphic to  $Y$ .

- 1) Note: the problem does not say that if  $f$  is not constant, then  $f$  is a homeomorphism.
- 2) Hint: Prove first that if  $f$  is not constant, then  $|X| = |Y|$ . Then examine the topology of  $Y$ .

E20. Suppose  $X$  is a countable set with the cofinite topology. State and prove a theorem that completely answers the question: “what sequences in  $(X, \mathcal{T})$  converge to what points?”

E21. Suppose that  $(X, \mathcal{T})$  and  $(Y, \mathcal{T}')$  are topological spaces. Recall that the product topology on  $X \times Y$  is the topology for which the collection of “open boxes”

$$\mathcal{B} = \{U \times V : U \in \mathcal{T}, V \in \mathcal{T}'\} \text{ is a base.}$$

- a) The “projection maps”  $\pi_1 : X \times Y \rightarrow X$  and  $\pi_2 : X \times Y \rightarrow Y$  are defined by

$$\pi_1(x, y) = x \text{ and } \pi_2(x, y) = y.$$

We showed in Example 5.11 that  $\pi_1$  and  $\pi_2$  are continuous. Prove that  $\pi_1$  and  $\pi_2$  are open maps. Give examples to show that  $\pi_1$  and  $\pi_2$  might not be closed.

b) Suppose that  $(Z, \mathcal{T}')$  is a topological space and that  $f : Z \rightarrow X \times Y$ . Prove that  $f$  is continuous iff both compositions  $\pi_1 \circ f : Z \rightarrow X$  and  $\pi_2 \circ f : Z \rightarrow Y$  are continuous. (Informally: a mapping into a product is continuous iff its composition with each projection is continuous.)

c) Prove that  $((x_n, y_n)) \rightarrow (x, y) \in X \times Y$  iff  $(x_n) \rightarrow x$  in  $X$  and  $(y_n) \rightarrow y$  in  $Y$ . (For this reason, the product topology is sometimes called the “topology of coordinatewise convergence.”)

d) Prove that  $X \times Y$  is homeomorphic to  $Y \times X$ . (Topological products are commutative.)

e) Prove that  $(X \times Y) \times Z$  is homeomorphic to  $X \times (Y \times Z)$  (Topological products are associative.)

E22. Let  $(X, \mathcal{T})$  and  $(Y, \mathcal{S})$  be topological spaces. Suppose  $f : X \rightarrow Y$ . Let

$$\Gamma(f) = \{(x, y) \in X \times Y : y = f(x)\} = \text{“the graph of } f\text{.”}$$

Prove that the map  $h : X \rightarrow \Gamma(f)$  defined by  $h(x) = (x, f(x))$  is a homeomorphism if and only if  $f$  is continuous.

*Note: if we think of  $f$  as a set of ordered pairs, the “graph of  $f$ ” is  $f$ . More informally, however, the problem states a function is continuous iff its graph is homeomorphic to its domain.*

E23. In  $(X, \mathcal{T})$ , a family of sets  $\mathcal{F} = \{B_\alpha : \alpha \in \Lambda\}$  is called locally finite if each point  $x \in X$  has a neighborhood  $N$  such that  $N \cap B_\alpha \neq \emptyset$  for only finitely many  $\alpha$ 's. (Part b) was also in Exercise E2.)

a) Suppose  $(X, d)$  is a metric space and that  $\mathcal{F}$  is a family of closed sets. Suppose there is an  $\epsilon > 0$  such that  $d(B_1, B_2) \geq \epsilon$  for all  $B_1, B_2 \in \mathcal{F}$ . Prove that  $\mathcal{F}$  is locally finite.

b) Prove that if  $\mathcal{F}$  is a locally finite family of sets in  $(X, \mathcal{T})$ , then  $\text{cl}(\bigcup_{\alpha \in A} B_\alpha) = \bigcup_{\alpha \in A} \text{cl}(B_\alpha)$ . Explain why this implies that if all the  $B_\alpha$ 's are closed, then  $\bigcup_{\alpha \in A} B_\alpha$  is closed. (This would apply, for example, to the sets in part a). See also Exercise E2.)

c) (The Pasting Lemmas: compare Exercise II.E24) Let  $(X, \mathcal{T})$  and  $(Y, \mathcal{T}')$  be topological spaces. For each  $\alpha \in A$ , suppose  $B_\alpha \subseteq X$ , that  $f_\alpha : B_\alpha \rightarrow Y$  is a continuous function, and that  $f_\alpha|_{(B_\alpha \cap B_\beta)} = f_\beta|_{(B_\alpha \cap B_\beta)}$  for all  $\alpha, \beta \in A$ . Then  $\bigcup_{\alpha \in A} f_\alpha = f$  is a function and  $f : \bigcup_{\alpha \in A} B_\alpha \rightarrow Y$ . (Informally: each pair of functions  $f_\alpha$  and  $f_\beta$  agree wherever their domains overlap; this allows use to define  $f$  by “pasting together” all the “function pieces”)

i) Show that if all the  $B_\alpha$ 's are open, then  $f$  is continuous.


ii) Show that if there are only finitely many  $B_\alpha$ 's and they are all closed, then  $f$  is continuous. (Hint: use a characterization of continuity in terms of closed sets.)

iii) Give an example to show that  $f$  might not be continuous when there are infinitely many  $B_\alpha$ 's all of which are closed.

iv) Show that if  $\mathcal{F}$  is a locally finite family of closed sets, then  $f$  is continuous. (Of course, iv)  $\Rightarrow$  ii) ).

*Note: the most common use of the Pasting Lemma is when the index set  $A$  is finite. For example, suppose*

$H_1 : [0, 1] \times [0, \frac{1}{2}] \rightarrow (X, d)$  is continuous, and  
 $H_2 : [0, 1] \times [\frac{1}{2}, 1] \rightarrow (X, d)$  is continuous, and  
 $H_1(t, \frac{1}{2}) = H_2(t, \frac{1}{2})$  for all  $t \in [0, 1]$



$H_1$  is defined on the lower closed half of the box  $[0, 1]^2$ ,  $H_2$  is defined on the upper closed half, and they agree on the “overlap” – that is, on the horizontal line segment  $[0, 1] \times \{\frac{1}{2}\}$ . Part b) (or part c) ) says that the two functions can be pieced together into a continuous function  $H : [0, 1]^2 \rightarrow (X, d)$ , where  $H = H_1 \cup H_2$ , that is

$$H(t, y) = \begin{cases} H_1(t, y) & \text{if } t \leq \frac{1}{2} \\ H_2(t, y) & \text{if } t \geq \frac{1}{2} \end{cases}$$

## Chapter III Review

Explain why each statement is true, or provide a counterexample. If nothing else is mentioned,  $X$  and  $Y$  are topological spaces with no other properties assumed.

1. For every possible topology  $\mathcal{T}$ , the space  $(\{0, 1, 2\}, \mathcal{T})$  is pseudometrizable.
2. A convergent sequence in a first countable topological space has at most one limit.
3. A one point set  $\{x\}$  in a pseudometric space  $(X, d)$  is closed.
4. Suppose  $\mathcal{T}$  and  $\mathcal{T}'$  are topologies on  $X$  and that for every subset  $A$  of  $X$ ,  $\text{cl}_{\mathcal{T}}(A) = \text{cl}_{\mathcal{T}'}(A)$ . Then  $\mathcal{T} = \mathcal{T}'$ .
5. Suppose  $f: X \rightarrow Y$  and  $A \subseteq X$ . If  $f|A: A \rightarrow Y$  is continuous, then  $f$  is continuous at each point of  $A$ .
6. Suppose  $\mathcal{S}$  is a subbase for the topology on  $X$  and that  $D \subseteq X$ . If  $U \cap D \neq \emptyset$  for every  $U \in \mathcal{S}$ , then  $D$  is dense in  $X$ .
7. If  $D$  is dense in  $(X, \mathcal{T})$  and  $\mathcal{T}^*$  is another topology on  $X$  with  $\mathcal{T} \subseteq \mathcal{T}^*$ , then  $D$  is dense in  $(X, \mathcal{T}^*)$ .
8. If  $f: X \rightarrow Y$  is both continuous and open, then  $f$  is also closed.
9. Every space is a continuous image of a first countable space.
10. Let  $X = \{0, 1\}$  with topology  $\mathcal{T} = \{\emptyset, X, \{1\}\}$ . There are exactly 3 continuous functions  $f: (X, \mathcal{T}) \rightarrow (X, \mathcal{T})$ .
11. If  $A$  and  $B$  are subspaces of  $(X, \mathcal{T})$  and both  $A$  and  $B$  are discrete in the subspace topology, then  $A \cup B$  is discrete in the subspace topology.
12. If  $A \subseteq (X, \mathcal{T})$  and  $X$  is separable, then  $A$  is separable.

13. If  $\mathcal{T}$  is the cofinite topology on  $X$ . Every bijection  $f: (X, \mathcal{T}) \rightarrow (X, \mathcal{T})$  is a homeomorphism.

14. If  $X$  has the cofinite topology, then the closure of any open set in  $X$  is open.

15. A continuous bijection from  $\mathbb{R}$  to  $\mathbb{R}$  must be a homeomorphism.

16. For every cardinal  $m$ , there is a separable topological space  $(X, \mathcal{T})$  with  $|X| = m$ .

17. If every family of disjoint open sets in  $(X, \mathcal{T})$  is countable, then  $(X, \mathcal{T})$  is separable.

18. For  $a \in \mathbb{R}^2$  and  $\epsilon > 0$ , let  $C_\epsilon(a) = \{x \in \mathbb{R}^2: d(x, a) = \epsilon\}$ . Let  $\mathcal{T}$  be the topology on  $\mathbb{R}^2$  for which the collection  $\{C_\epsilon(a): \epsilon > 0, a \in \mathbb{R}^2\}$  is a subbasis. Let  $\mathcal{U}$  be the usual topology on  $\mathbb{R}^2$ . Then the function  $f: (\mathbb{R}^2, \mathcal{T}) \rightarrow (\mathbb{R}^2, \mathcal{U})$  given by  $f(x, y) = (\sin x, \sin y)$  is continuous.

19. If  $D \subseteq \mathbb{R}$  and each point of  $D$  is isolated in  $D$ , then  $\text{cl } D$  must be countable.

20. Consider the separation property  $S$  : every minimal nonempty closed set  $F$  is a singleton.  
 $(F$  is a minimal nonempty closed set means: if  $A$  is a nonempty closed set and  $A \subseteq F$ , then  $A = F$ ). If  $X$  is a  $T_1$ -space, then  $X$  has property  $S$ .

21. A one-to-one, continuous, onto map  $f: (X, \mathcal{T}) \rightarrow (Y, \mathcal{T}')$  must be a homeomorphism.

22. An uncountable closed set in  $\mathbb{R}$  must contain an interval of positive length.

23. A countable metric space has a base consisting of clopen sets.

24. Suppose  $D \subseteq X$  and that  $D$  is dense in  $(X, \mathcal{T})$ . If  $\mathcal{T}^*$  is topology on  $X$  with  $\mathcal{T}^* \subseteq \mathcal{T}$ , then  $D$  is dense in  $(X, \mathcal{T}^*)$ .

25. If  $D \subseteq \mathbb{R}$  and  $D$  is discrete in the subspace topology, then  $D$  is countable.

26. The Sorgenfrey plane has a subspace homeomorphic to  $\mathbb{R}$  (with its usual topology).

27. In  $\mathbb{N}$ , let  $\mathcal{B}_n = \{B \subseteq \mathbb{N} : n \in B \text{ and } B \subseteq \{1, 2, \dots, n\}\}$ . At each point  $n$ , the  $\mathcal{B}_n$ 's satisfy the conditions in the Neighborhood Base Theorem and therefore describe a topology on  $\mathbb{N}$ .

28. At each point  $n \in \mathbb{N}$ , let  $\mathcal{B}_n = \{B : B \supseteq \{n, n+1, n+2, \dots\}\}$ . At each point  $n$ , the  $\mathcal{B}_n$ 's satisfy the conditions in the Neighborhood Base Theorem and therefore describe a topology on  $\mathbb{N}$ .

29. Suppose  $(X, T)$  has a base  $\mathcal{B}$  with  $|\mathcal{B}| = c$ . Then  $X$  has a dense set  $D$  with  $|D| \leq c$ .

30. Suppose  $(X, T)$  has a base  $\mathcal{B}$  with  $|\mathcal{B}| = c$ . Then at each point  $x \in X$ , there is a neighborhood base with  $|\mathcal{B}_x| \leq c$ .

31. Suppose  $X$  is an infinite set. Let  $T_1$  be the cofinite topology on  $X$  and let  $T_2$  be the discrete topology on  $X$ . If a function  $f : \mathbb{R} \rightarrow (X, T_1)$  is continuous, then  $f : \mathbb{R} \rightarrow (X, T_2)$  is also continuous.

32. Let  $T$  be the “right-ray topology” on  $\mathbb{R}$ , that is,  $T = \{(a, \infty) : a \in \mathbb{R}\} \cup \{\emptyset, \mathbb{R}\}$ . The space  $(\mathbb{R}, T)$  is first countable.