

Lecture 14: Determinants (concl.)

Determinants and Volume

Let $\vec{v}_1, \dots, \vec{v}_n \in \mathbb{R}^n$. What is the n -volume of

$$P = P(\vec{v}_1, \dots, \vec{v}_n) := \{ a_1 \vec{v}_1 + \dots + a_n \vec{v}_n \mid 0 \leq a_i \leq 1 \text{ (for each } i) \} ?$$

A basic observation is that if you swap \vec{v}_i & \vec{v}_j it doesn't change the volume; while if you scale (multiply) the length of \vec{v}_i by μ , it multiplies the volume by $|\mu|$.

Finally, if you replace \vec{v}_j by $\vec{v}_j + a \vec{v}_i$, this causes a shear of the parallelepiped P , and shears don't affect volume. But these are precisely the effects that these operations have on $|\det(A)|$ (the absolute value of $\det(A)$), where $A = \begin{pmatrix} \vec{v}_1 \rightarrow \\ \vdots \\ \vec{v}_n \rightarrow \end{pmatrix}$.

More precisely, if a sequence of row operations gets you from $\vec{e}_1, \dots, \vec{e}_n$ to $\vec{v}_1, \dots, \vec{v}_n$ (as rows of a matrix), and involves scaling by μ_1, \dots, μ_n and N swaps, then $\det A = (-1)^N \prod_{i=1}^n \mu_i$. Now $\text{vol}(P(\vec{e}_1, \dots, \vec{e}_n)) = 1$, so $\text{vol}(P(\vec{v}_1, \dots, \vec{v}_n)) = 1 \cdot \prod_{i=1}^n |\mu_i|$ by the above observations, and this is $|\det A|$.

Theorem 1: $\text{Vol} \{P(\vec{v}_1, \dots, \vec{v}_n)\} = \left| \det \begin{pmatrix} \leftarrow \vec{v}_1 \rightarrow \\ \vdots \\ \leftarrow \vec{v}_n \rightarrow \end{pmatrix} \right|$

$$= \left| \det \begin{pmatrix} \uparrow \vec{v}_1 \downarrow & \dots & \uparrow \vec{v}_n \downarrow \\ \vdots & & \vdots \end{pmatrix} \right|$$

transpose

Now let $T: \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a linear transformation with matrix M .

Corollary: If $\Omega \subset \mathbb{R}^n$ is any bounded subset,

$$\frac{\text{Vol} \{T(\Omega)\}}{\text{Vol} \{\Omega\}} = |\det(M)|.$$

"dilation factor"

Proof: Covering Ω with little parallelepipeds and taking a limit as their size $\rightarrow 0$, we see it suffices to check the result for such parallelepipeds:

$$\frac{\text{Vol} \{T(P(\vec{v}_1, \dots, \vec{v}_n))\}}{\text{Vol} \{P(\vec{v}_1, \dots, \vec{v}_n)\}} = \frac{\text{Vol} \{P(T\vec{v}_1, \dots, T\vec{v}_n)\}}{\text{Vol} \{P(\vec{v}_1, \dots, \vec{v}_n)\}}$$

$$= \left| \frac{\det \begin{pmatrix} \uparrow M\vec{v}_1 \downarrow & \dots & \uparrow M\vec{v}_n \downarrow \\ \vdots & & \vdots \end{pmatrix}}{\det \begin{pmatrix} \uparrow \vec{v}_1 \downarrow & \dots & \uparrow \vec{v}_n \downarrow \\ \vdots & & \vdots \end{pmatrix}} \right| = \left| \frac{\det \left(M \cdot \begin{pmatrix} \uparrow \vec{v}_1 \downarrow & \dots & \uparrow \vec{v}_n \downarrow \\ \vdots & & \vdots \end{pmatrix} \right)}{\det \begin{pmatrix} \uparrow \vec{v}_1 \downarrow & \dots & \uparrow \vec{v}_n \downarrow \\ \vdots & & \vdots \end{pmatrix}} \right|$$

$$= \left| \frac{\det M \cdot \cancel{\det(\vec{v}_1 \dots \vec{v}_n)}}{\cancel{\det(\vec{v}_1 \dots \vec{v}_n)}} \right| = |\det M|. \quad \square$$

Ex $\vee \Omega = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 \leq 1\}$

$M = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}$ matrix of $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$

$\Rightarrow T(\Omega) = \{(x, y) \in \mathbb{R}^2 \mid \frac{x^2}{a^2} + \frac{y^2}{b^2} \leq 1\}$

$\text{area}(T(\Omega)) = \text{area}(\Omega) \cdot |\det M| = \pi ab.$ //


Remark: In calculus, this shows up in the change-of-variables formula for multiple integrals. Given a nonlinear smooth change, $(x_1, \dots, x_n) \mapsto (y_1(x_1, \dots, x_n), \dots, y_n(x_1, \dots, x_n))$, the Jacobian matrix is $J = \begin{pmatrix} \partial y_1 / \partial x_1 & \dots & \partial y_1 / \partial x_n \\ \vdots & \ddots & \vdots \\ \partial y_n / \partial x_1 & \dots & \partial y_n / \partial x_n \end{pmatrix}$, and

$|\det J|$ is the "infinitesimal dilation factor" giving

the ratio $\frac{dy_1 \wedge \dots \wedge dy_n}{dx_1 \wedge \dots \wedge dx_n}$ of volumes of infinitesimal

parallelepipeds.]

Ex 2 / Find the area of the parallelogram with vertices $(-1, -2)$, $(1, 3)$, $(2, 4)$, and $(0, -1)$.



$$\vec{v}_1 = \begin{pmatrix} 0 \\ -1 \end{pmatrix} - \begin{pmatrix} -1 \\ -2 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \quad \vec{v}_2 = \begin{pmatrix} 1 \\ 3 \end{pmatrix} - \begin{pmatrix} -1 \\ -2 \end{pmatrix} = \begin{pmatrix} 2 \\ 5 \end{pmatrix}$$

$$\text{Area} \{ P(\vec{v}_1, \vec{v}_2) \} = \left| \det \begin{pmatrix} 1 & 2 \\ 1 & 5 \end{pmatrix} \right| = 3.$$

Cramer's Rule

For $A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$, $A^{-1} = \frac{1}{\det A} \begin{pmatrix} a_{22} & -a_{12} \\ -a_{21} & a_{11} \end{pmatrix}$.

So the solution to $A \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \end{pmatrix}$ is

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = A^{-1} \begin{pmatrix} b_1 \\ b_2 \end{pmatrix} = \frac{1}{\det A} \begin{pmatrix} a_{22}b_1 - a_{12}b_2 \\ a_{11}b_2 - a_{21}b_1 \end{pmatrix}$$

$$\leadsto x_1 = \frac{\det \begin{pmatrix} b_1 & a_{12} \\ b_2 & a_{22} \end{pmatrix}}{\det A}, \quad x_2 = \frac{\det \begin{pmatrix} a_{11} & b_1 \\ a_{21} & b_2 \end{pmatrix}}{\det A}$$

Theorem 2: For A $n \times n$ invertible with columns \vec{c}_j , the unique solution of $A\vec{x} = \vec{b}$ is given componentwise by

$$x_i = \frac{\det(A_i(\vec{b}))}{\det(A)} \quad (i=1, \dots, n)$$

where $A_i(\vec{b})$ is the $n \times n$ matrix obtained by replacing column \vec{c}_i by \vec{b} .

Proof: $\det A_i(\vec{b}) = \det \begin{pmatrix} \uparrow & \uparrow & \uparrow \\ \vec{c}_1 & \vec{b} & \vec{c}_n \\ \downarrow & \downarrow & \downarrow \end{pmatrix} = \det \begin{pmatrix} \uparrow & \uparrow & \uparrow \\ \vec{c}_1 & x_1 \vec{c}_i & \vec{c}_n \\ \downarrow & \downarrow & \downarrow \end{pmatrix}$

$= \det \begin{pmatrix} \uparrow & \uparrow & \uparrow \\ \vec{c}_1 & (\sum_{j=1}^n x_j \vec{c}_j) & \vec{c}_n \\ \downarrow & \downarrow & \downarrow \end{pmatrix} = \sum_{j=1}^n x_j \det \begin{pmatrix} \uparrow & \uparrow & \uparrow \\ \vec{c}_1 & \vec{c}_j & \vec{c}_n \\ \downarrow & \downarrow & \downarrow \end{pmatrix}$

$= x_i \det \begin{pmatrix} \uparrow & \uparrow & \uparrow \\ \vec{c}_1 & \vec{c}_i & \vec{c}_n \\ \downarrow & \downarrow & \downarrow \end{pmatrix} = x_i \det A.$

if $j \neq i$, then \vec{c}_j appears twice $\Rightarrow \det = 0$

Ex 3 / Solve $\begin{cases} sx - ty = 1 \\ sy - tx = a \\ sz - tz = 0 \end{cases}$ for arbitrary $s, t, a \in \mathbb{R}$ (when possible).

$\begin{pmatrix} s & -t & 0 \\ 0 & s & -t \\ -t & 0 & s \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ a \\ 0 \end{pmatrix}, \quad \det A = s \begin{vmatrix} s & -t \\ 0 & s \end{vmatrix} - t \begin{vmatrix} -t & 0 \\ s & -t \end{vmatrix} = s^3 - t^3.$

$A \quad \vec{x} \quad \vec{b}$

So assume $s^3 \neq t^3$ (or just $s \neq t$, equivalently), so the system is solvable. By Cramer,

$x = \frac{\begin{vmatrix} 1 & -t & 0 \\ a & s & -t \\ 0 & 0 & s \end{vmatrix}}{\det A} = \frac{\begin{vmatrix} 1 & -t \\ a & s \end{vmatrix}}{s^3 - t^3} = \frac{s^2 + ast}{s^3 - t^3}$

$$y = \frac{\begin{vmatrix} s & 1 & 0 \\ 0 & a & -c \\ -c & 0 & s \end{vmatrix}}{s^3 - c^3} = \frac{as^2 + c^2}{s^3 - c^3}, \quad z = \frac{\begin{vmatrix} s & -c & 1 \\ 0 & s & a \\ -c & 0 & 0 \end{vmatrix}}{s^3 - c^3} = \frac{ac^2 + s^2c}{s^3 - c^3}.$$


Clearly not something you'd want to do by row-reduction. //

The adjugate matrix

Let $A =$ invertible $n \times n$. Writing $A^{-1} = \begin{pmatrix} \vec{\alpha}_1 & \dots & \vec{\alpha}_n \\ \vdots & & \vdots \end{pmatrix} = \begin{pmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{n1} & \dots & a_{nn} \end{pmatrix}$

$$A \cdot A^{-1} = I_n \Rightarrow A \cdot \vec{\alpha}_j = \vec{e}_j \text{ for each } j=1, \dots, n.$$

By Cramer's rule,

$$d_{ij} = i^{\text{th}} \text{ entry of } \vec{\alpha}_j = \frac{\det(A_i^{\wedge}(\vec{e}_j))}{\det(A)}$$


$$= \frac{(-1)^{j+i} \det(A_{ji}^{\wedge})}{\det A} = \frac{C_{ji}}{\det(A)}$$

Laplace
(ith column)

Set $\text{adj}(A) := n \times n$ matrix with $(i, j)^{\text{th}}$ entry C_{ji} .

Theorem 3: $A^{-1} = \frac{1}{\det(A)} \cdot \text{adj}(A).$

This generalises the 2×2 formula, though is generally not preferred to the row-reduction approach unless there

are variables (or nasty numbers) in the entries of A .

Ex 4/Find A^{-1} in Example 3.

$$A = \begin{pmatrix} s & -t & 0 \\ 0 & s & -t \\ -t & 0 & s \end{pmatrix} \Rightarrow \det A = s^3 - t^3 \text{ and}$$

$$C_{11} = \begin{vmatrix} s & -t \\ 0 & s \end{vmatrix} = s^2, \quad C_{21} = -\begin{vmatrix} -t & 0 \\ 0 & s \end{vmatrix} = st, \quad C_{31} = \begin{vmatrix} -t & 0 \\ s & -t \end{vmatrix} = t^2$$

$$C_{12} = -\begin{vmatrix} 0 & -t \\ -t & s \end{vmatrix} = t^2, \quad C_{22} = \begin{vmatrix} s & 0 \\ -t & s \end{vmatrix} = s^2, \quad C_{32} = -\begin{vmatrix} s & 0 \\ 0 & -t \end{vmatrix} = st$$

$$C_{13} = \begin{vmatrix} 0 & s \\ -t & 0 \end{vmatrix} = st, \quad C_{23} = -\begin{vmatrix} s & -t \\ -t & 0 \end{vmatrix} = t^2, \quad C_{33} = \begin{vmatrix} s & -t \\ 0 & s \end{vmatrix} = s^2$$

$$\Rightarrow A^{-1} = \frac{1}{s^3 - t^3} \begin{pmatrix} s^2 & st & t^2 \\ t^2 & s^2 & st \\ st & t^2 & s^2 \end{pmatrix}.$$