

Lecture 21: Change of Basis

A few lectures back we discussed writing vectors

$\vec{v} \in V$ with respect to a basis $\mathcal{B} = \{\vec{b}_1, \dots, \vec{b}_n\}$ for V :

$$[\vec{x}]_{\mathcal{B}} = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} \iff \vec{x} = \alpha_1 \vec{b}_1 + \dots + \alpha_n \vec{b}_n.$$

When $V = \mathbb{R}^n$, we had the change-of-basis matrix

$$P_{\mathcal{B}} = \begin{pmatrix} \uparrow & & \uparrow \\ \vec{b}_1 & \cdots & \vec{b}_n \\ \downarrow & & \downarrow \end{pmatrix}$$

and the relation

$$P_{\mathcal{B}} [\vec{x}]_{\mathcal{B}} = \vec{x}.$$

Ex 1 / $V = \mathbb{R}^2$, $\vec{x} = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$. Find coordinates of \vec{x} with respect to the rotated axes $\mathcal{B} = \left\{ \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \right\}$



Write $P_{\mathcal{B}} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$, $\Rightarrow P_{\mathcal{B}}^{-1} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix}$ (why?).

$$\text{Then } [\vec{x}]_{\mathcal{B}} = P_{\mathcal{B}}^{-1} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} x_1 \cos \theta + x_2 \sin \theta \\ -x_1 \sin \theta + x_2 \cos \theta \end{pmatrix}.$$

Ex 2 / Let V be some 2-dimensional vector space with bases

$$B = \{\vec{b}_1, \vec{b}_2\} \text{ and } C = \{\vec{c}_1, \vec{c}_2\}.$$

Suppose we're given $\vec{b}_1 = -\vec{c}_1 + 4\vec{c}_2$ & $\vec{b}_2 = 3\vec{c}_1 - 6\vec{c}_2$.

If $\vec{x} = 2\vec{b}_1 + 3\vec{b}_2$, what is $[\vec{x}]_C$?

$$\begin{aligned} \text{Write } \vec{x} &= 2(-\vec{c}_1 + 4\vec{c}_2) + 3(3\vec{c}_1 - 6\vec{c}_2) \\ &= (-1 \cdot 2 + 3 \cdot 3)\vec{c}_1 + (4 \cdot 2 - 6 \cdot 3)\vec{c}_2 = 7\vec{c}_1 - 10\vec{c}_2. \end{aligned}$$

Notice that this also can be rewritten as...

$$[\vec{x}]_C = \begin{pmatrix} -1 & 3 \\ 4 & -6 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} \uparrow & \uparrow \\ [\vec{b}_1]_C & [\vec{b}_2]_C \\ \downarrow & \downarrow \end{pmatrix} \begin{pmatrix} \uparrow \\ [\vec{x}]_B \\ \downarrow \end{pmatrix} =: \underset{C \leftarrow B}{P} [\vec{x}]_B$$

More generally, if $C = \{\vec{c}_1, \dots, \vec{c}_n\}$ and $B = \{\vec{b}_1, \dots, \vec{b}_n\}$

are bases for V , set

$$P_{C \leftarrow B} := \begin{pmatrix} \uparrow & \uparrow \\ [\vec{b}_1]_C & \dots & [\vec{b}_n]_C \\ \downarrow & & \downarrow \end{pmatrix}$$

so that if $[\vec{x}]_B = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix}$, then

$$P_{C \leftarrow B} [\vec{x}]_B = \alpha_1 [\vec{b}_1]_C + \dots + \alpha_n [\vec{b}_n]_C = [\alpha_1 \vec{b}_1 + \dots + \alpha_n \vec{b}_n]_C = [\vec{x}]_C.$$

Properties: Since $P_{C \leftarrow B} [\vec{x}]_B = [\vec{x}]_C$, $P_{C \leftarrow B}^{-1} [\vec{x}]_C = [\vec{x}]_B = P_{B \leftarrow C} [\vec{x}]_C$

for all \vec{x} , and so

$$(i) \quad P_{C \leftarrow B}^{-1} = P_{B \leftarrow C}.$$

Given another basis A , we have also

$$(ii) \quad P_{C \leftarrow B} \cdot P_{B \leftarrow A} = P_{C \leftarrow A}.$$

Ex 3 / Rewrite $f(t) = 5 - 6t + 3t^2 + t^3$ in powers of $(t-1)$.

In P_3 , write $B = \{1, t, t^2, t^3\}$ and $C = \{1, t-1, (t-1)^2, (t-1)^3\}$,

so that $[f]_B = \begin{pmatrix} 5 \\ -6 \\ 3 \\ 1 \end{pmatrix}$. Now $P_{B \leftarrow C} = \begin{pmatrix} \uparrow & & & \\ [c_1]_B & \dots & [c_4]_B & \\ \downarrow & & & \end{pmatrix} = \begin{pmatrix} 1 & -1 & 1 & -1 \\ 0 & 1 & -2 & 3 \\ 0 & 0 & 1 & -3 \\ 0 & 0 & 0 & 1 \end{pmatrix}$

\implies take inverse using RREF $P_{C \leftarrow B} = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{pmatrix}$, and thus $[f]_C = P_{C \leftarrow B} [f]_B = \begin{pmatrix} 3 \\ 3 \\ 6 \\ 1 \end{pmatrix}$.

That is, $f(t) = 3 + 3(t-1) + 6(t-1)^2 + (t-1)^3$.

Now let $V = \mathbb{R}^n$, $E = \{\vec{e}_1, \dots, \vec{e}_n\}$ (standard basis), and B & C other bases.

Properties: Since $P_B [\vec{x}]_B = \vec{x} = [\vec{x}]_E$, really

- $P_B = E \leftarrow B$, and $P_B^{-1} = B \leftarrow E$.

Combining this with (ii) above,

- $P_C^{-1} P_B = C \leftarrow E \cdot E \leftarrow B = C \leftarrow B$.

Think: P_B sends $[\vec{x}]_B$ to \vec{x} , then P_C^{-1} sends \vec{x} to $[\vec{x}]_C$.

Ex 4 / In \mathbb{R}^2 , find the change-of-basis matrix from

$$C = \left\{ \begin{pmatrix} 7 \\ 5 \end{pmatrix}, \begin{pmatrix} -3 \\ 1 \end{pmatrix} \right\} \text{ to } B = \left\{ \begin{pmatrix} 1 \\ -5 \end{pmatrix}, \begin{pmatrix} -2 \\ 2 \end{pmatrix} \right\}.$$

$$P_B = \begin{pmatrix} 1 & -2 \\ -5 & 2 \end{pmatrix} \implies P_B^{-1} = \frac{1}{1 \cdot 2 - (-2) \cdot (-5)} \begin{pmatrix} 2 & 2 \\ 5 & 1 \end{pmatrix} = -\frac{1}{8} \begin{pmatrix} 2 & 2 \\ 5 & 1 \end{pmatrix}$$

$$\implies P_{B \leftarrow C} = P_B^{-1} P_C = -\frac{1}{8} \begin{pmatrix} 2 & 2 \\ 5 & 1 \end{pmatrix} \begin{pmatrix} 7 & -3 \\ 5 & 1 \end{pmatrix} = \begin{pmatrix} -3 & \frac{1}{2} \\ -5 & \frac{7}{4} \end{pmatrix}$$

$[c_1]_B \quad [c_2]_B$

Here is another way to proceed. Consider that we want to find the coordinate vectors of \vec{c}_1, \vec{c}_2 wrt \mathcal{B} . That is, we want to solve

$$P_{\mathcal{B}} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \vec{c}_1, \quad P_{\mathcal{B}} \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = \vec{c}_2,$$

for which one would ordinarily row-reduce $[P_{\mathcal{B}} | \vec{c}_1], [P_{\mathcal{B}} | \vec{c}_2]$.

An even better way to think of this is: adjoining both \vec{c}_1 & \vec{c}_2 to $P_{\mathcal{B}}$, we get

$$[P_{\mathcal{B}} | P_{\mathcal{C}}] = [\vec{b}_1 \ \vec{b}_2 | \vec{c}_1 \ \vec{c}_2] \xrightarrow{\text{RREF}} \left[\mathbb{I}_2 \mid \begin{matrix} \text{the desired } \begin{pmatrix} x_1 & y_1 \\ x_2 & y_2 \end{pmatrix} \\ M \end{matrix} \right].$$

To see what M is, notice that the row-reduction has to multiply $P_{\mathcal{B}}$ by $P_{\mathcal{B}}^{-1}$ to get \mathbb{I}_2 . So

$$M = P_{\mathcal{B}}^{-1} P_{\mathcal{C}} = P_{\mathcal{B} \leftarrow \mathcal{C}}.$$

Ex 4
(cont'd) /

$$\left[\begin{array}{cc|cc} 1 & -2 & 7 & -3 \\ -5 & 2 & 5 & 1 \end{array} \right] \rightarrow \left[\begin{array}{cc|cc} 1 & -2 & 7 & -3 \\ 0 & -8 & 40 & -14 \end{array} \right]$$

$$\begin{matrix} \vec{b}_1 & \vec{b}_2 & \vec{c}_1 & \vec{c}_2 \end{matrix} \rightarrow \left[\begin{array}{cc|cc} 1 & -2 & 7 & -3 \\ 0 & 1 & -5 & 7/4 \end{array} \right]$$

$$\rightarrow \left[\begin{array}{cc|cc} 1 & 0 & -3 & 1/2 \\ 0 & 1 & -5 & 7/4 \end{array} \right]$$

